

1 **Factors associated with positive urine cultures in cats with subcutaneous ureteral**
2 **bypass (SUB) system implantation.**

3 Catrina E Pennington, Department of Surgery, University of Liverpool, United Kingdom

4 Zoe Halfacree, Department of Surgery, Royal Veterinary College, London, United Kingdom

5 Charlotte Colville-Hyde, Corner House Veterinary Surgery in Exmouth, Devon.

6 Rebecca F Geddes, MA VetMB MVetMed PhD DipACVIM MRCVS, Department of Internal
7 Medicine, Royal Veterinary College, London, United Kingdom (**Corresponding author** – Email
8 address: rgeddes@rvc.ac.uk)

9

10

11

12

13

14

15

16

17 **Abstract:**

18 OBJECTIVES

19 To report the postoperative incidence of SUB-associated bacteriuria and risk factors in a large
20 population of UK cats, to identify the commonly implicated isolates in these cases and to
21 report associations of positive post-operative urine cultures with device occlusion or need for
22 further surgery.

23 METHODS

24 Electronic clinical records were reviewed to identify cats with ureteral obstruction who
25 underwent unilateral or bilateral SUB implantation between September 2011 and September
26 2019. One hundred and eighteen client-owned cats were included in the study population.
27 Information recorded included signalment, history, surgical and biochemical factors, urinalysis
28 and culture results. Multivariable logistic regression was performed to identify variables
29 associated with a positive post-operative culture.

30 RESULTS

31 In total, 8.5% of cats had a positive post-operative culture within one-month post-surgery and
32 41.2% within one year post-surgery.

33 Cats with a positive pre-operative culture were significantly more likely to have a positive
34 culture within six months post-operatively ($p=0.026$ OR 0.245 CI 0.071-0.848). Of the 14 cats
35 with a positive pre-operative culture, six (42.9%) returned a positive culture within one year
36 post-operatively and in four cases (66.7%) the same isolate was identified.

37 Cats with higher end-anaesthetic rectal temperatures were significantly less likely to return a
38 positive culture within three months ($p=0.006$ OR 0.398 CI 0.205-0.772) post-surgery.
39 Cats culturing positive for Escherichia coli at any time point ($p=0.008$ or 4.542 CI 1.485 -13.89)
40 were significantly more likely to have their implant removed or replaced.

41 CONCLUSIONS AND CLINICAL RELEVANCE

42 Peri-operative hypothermia and pre-operative positive culture were independent predictors of
43 a post-operative positive culture and this should be taken into consideration when managing
44 these cases. Positive post-operative culture rates were higher than have previously been
45 reported.

46 **Keywords:** subcutaneous ureteral bypass, SUB, bacteriuria, ureterolithiasis

47

48

49 **Factors associated with positive urine cultures in cats with subcutaneous ureteral**
50 **bypass (SUB) system implantation.**

51

52 Introduction

53 Intraluminal ureteral obstruction is an occurrence of increasing incidence in feline
54 medicine; with ureterolithiasis implicated in the majority of cases (1). Ureteral
55 obstruction may also be diagnosed secondary to stricture, stenosis, iatrogenic injury,
56 thrombi and neoplasia (2).

57 Successful management of ureteral obstruction depends on prompt renal
58 decompression to prevent permanent loss of glomerular filtration capacity (3).
59 Historically where medical treatment failed, **nephrectomy** was indicated; however in
60 recent years many institutions have adopted ureteral bypass using the SUB system (4)
61 placement as the treatment of choice.

62 The SUB system consists of two catheters, placed in the renal pelvis and bladder
63 lumen, connected to a shunting port sited in the subcutaneous tissue (4), allowing
64 urine to bypass the affected ureter. SUB placement mortality rates compare
65 favourably to more traditional surgical techniques, with 0% - 13% perioperative
66 mortality reported (5, 6) compared to 18% perioperative mortality for both traditional
67 interventions (1) and ureteral stent placement (6). Intraoperative complications are
68 reported to occur in 7% of cases (5) and perioperative complications necessitating
69 surgical intervention reported in 7% and 9% of cases respectively (5, 6); with device
70 occlusion, device leakage and urethral obstruction most commonly observed.

71 Bacteriuria following SUB placement is reported in 8-21% of cats (5, 7). Therefore
72 identifying factors predictive of device colonisation and the clinical impact of this may
73 help guide client decision-making.

74 Risk factors for positive urine cultures in cats with SUB system implantation have
75 previously been described in a small population of 43 cats within an American
76 institution (8), this study population included cats that underwent SUB system or
77 ureteral stent placement. The authors reported that of all variables assessed only use
78 of post-operative antibiotics was associated with likelihood of positive urine culture.
79 Berent et al (2018) reported that both positive pre-operative culture and post-
80 operative use of an indwelling catheter were associated with positive post-operative
81 culture, however other factors were not assessed.

82 The microbial profile of SUB-system-associated bacteriuria has been reported twice
83 previously (8, 5), both studies from American cat populations. The clinical implications
84 of SUB-associated bacteriuria, in terms of association with device obstruction and
85 requirement for surgical intervention, have not been previously reported.

86 The objectives of this study were thus threefold. Firstly to report the postoperative
87 incidence of SUB-associated bacteriuria and risk factors in a large population of UK
88 cats; secondly to identify the commonly implicated isolates in these cases; and thirdly

89 to report associations of positive post-operative urine cultures with device occlusion or
90 need for further surgery.

91 Materials and methods

92 Clinical records of all cats with implantation of a SUB system at the Queen Mother
93 Hospital for Small Animal, London between September 2011 and September 2019
94 were reviewed. Cases were excluded if they had not survived to discharge. Only cases
95 with a minimum of one urine culture performed at least two weeks post-surgery were
96 included. Information recorded included signalment, history, surgical and biochemical
97 factors in addition to urinalysis and culture results. Initial diagnosis of ureteral
98 obstruction was made via abdominal ultrasound performed by board-certified
99 radiologists or residents in training, with pyelography performed to confirm aetiology
100 as required.

101 Surgery was performed by board-certified surgeons or residents in training, with
102 general anaesthesia supervised by board-certified anaesthetists or residents. A SUB
103 device (Norfolk Vet Products) was implanted as described by the manufacturer (4)
104 under fluoroscopic guidance.

105 In cats with an active urine sediment, intravenous antibiotic administration was
106 initiated preoperatively; surgery was delayed up to 48 hours during antibiotic
107 treatment if clinical condition allowed. Cats with an active sediment were continued
108 on a therapeutic course of antibiotics for 4-6 weeks post-operatively based upon
109 culture and sensitivity results of urine obtained pre-operatively or from the renal pelvis
110 at surgery.

111 Cats that survived to discharge were scheduled to revisit the hospital approximately
112 one month post-operatively, at which point a urine sample was collected via aspiration
113 of the subcutaneous port via a sterile technique. Follow-up appointments were then
114 recommended every three months for the first two years and every six months
115 thereafter, when the same technique was performed. From February 2019 standard
116 hospital protocol was revised to include flushing of the systems with 2.5ml of sterile
117 Tetrasodium Ethylenediaminetetraacetic acid (TetraEDTA) per port post sample
118 collection.

119 Cats with positive urine cultures were generally administered a 3-8 weeks course of
120 appropriate antibiotics based on culture and sensitivity results with culture performed
121 during and at least one week post-treatment. Where empirical antibiotics was required
122 pending culture a 4-6 week course of oral potentiated amoxicillin was prescribed with

123 change of agent as required on return of culture results. Antibiosis was often stopped
124 or not re-prescribed for cats with recurrent or persistent positive urine cultures
125 without lower urinary tract signs

126 All samples were submitted to the same laboratory. Samples were streaked with an
127 inoculation loop onto MacConkey agar and Columbia agar supplemented with 5%
128 sheep blood. Samples were incubated for up to 24 hours in 5% carbon dioxide at 37 °C.

129 Bacterial identification was based on Gram-staining, colony type and morphology in
130 addition to routine biochemical testing. Isolate sensitivity was established with the
131 Kirby-Bauer disc diffusion method performed in accordance with Clinical and
132 Laboratory Standards Institute guidelines (9).

133 Statistical analysis

134 Descriptive analysis was used for signalment, treatment and sample data. Data were
135 analysed with SPSS version 24 (IBM, Armonk, New York) with significance set at
136 $p < 0.05$. Variables were tested for normality using a Shapiro-Wilk test. Normally
137 distributed data is reported as mean, standard deviation, non-Gaussian data is
138 reported as median, range.

139 A chi-square test was used to compare likelihood of device explantation in bacteriuric
140 cats, between those with clinical signs and those without.

141 Univariable logistic regression analysis was initially used to assess predictive factors for
142 the following six models; first positive culture results within one (0-30 days), three (0-
143 90 days), six (0-180 days) and 12 (0-365 days) months respectively, for device occlusion
144 at any time point and for device removal at any time-point. Variables for which $p < 0.1$
145 were then further explored using a multivariable analysis, with backward elimination
146 to identify variables for which $p < 0.05$.

147 Results

148 One hundred and twenty-four cats had a SUB implanted between the 16th April 2012
149 and the 6th September 2019, but six cats did not survive to discharge, therefore 118
150 cats met the inclusion criteria.

151 Mean age at the time of device placement was 7.17 years (SD: 3.35). Of the 118 cats
152 included in the study, 70 (59.3%) were neutered females, 47 (39.8%) were neutered
153 males and one was an entire male. Breeds representing over 2% of the population are
154 summarised in table 1.

155 Median body weight at the time of surgery was 4kg (range 2.1-9.7kg), median body
156 condition score was 4/9, (range 1-9).

157 In 93 cats (78.8%) a SUB system was placed due to ureterolithiasis, four cats (3.4%)
158 presented due to iatrogenic injury (three due to inadvertent ureteral ligation during
159 ovariohysterectomy and one due to stricture formation post ureterotomy), one cat
160 (0.8%) was diagnosed with a ureteral stricture in the absence of ureteroliths or known
161 prior trauma and in 20 cats (16.9%) the underlying pathology was not confirmed.

162 Forty six (39%) cases had a bilateral system placed and 72 (61%) a unilateral system. Of
163 the unilateral systems 41 (56.9%) were left-sided and 31 (43.1%) right-sided. Median
164 surgical and anaesthesia time were 105 minutes (range 50-275) and 200 minutes
165 (range 125-420) respectively. 64 cats (54.2%) received potentiated amoxicillin peri-
166 operatively, 39 cats (33.1%) received cefuroxime, in 15 cats (12.7%) perioperative
167 antibiotic choice was not recorded.

168 Cats were hospitalised for a median of 6 days (range 2- 23) post-surgery.

169 Pre-operative and post-operative (obtained between one and three months post-
170 surgery) clinicopathologic data is summarised in table 2.

171 One hundred and ten cats had a pre-operative culture performed, of these 14 (12.7%)
172 were positive. Thirteen cultures identified a single isolate and one culture identified
173 two isolates. The most frequent isolates were *E. coli* (42.9%) and *Enterococcus faecalis*
174 (28.6%). Of the 14 cats with positive pre-operative cultures six (42.9%) had a positive
175 post-operative culture, all between day 20 and day 176 post-surgery. Of these six
176 cultures four (66.7%) identified the same isolate both pre and post operatively. Of
177 these four cultures, two isolated *E. coli*, one *E. faecalis* and one *Staphylococcus*
178 *pseudintermedius*.

179 At least one month follow-up was available for all 118 cats. Of these, 10 (8.5%) had a
180 first positive post-operative culture within one month post-surgery. The most common
181 organisms isolated were *E. coli* (40%) and *Pseudomonas aeruginosa* (30%). Only one of
182 these 10 cats had had a positive pre-operative urine culture, and the bacteria cultured
183 was different (*E.coli* pre-operatively and *S. pseudintermedius* post-operatively).

184 A minimum of three months follow-up was available for 112 cats, of these 15 (13.4%)
185 had a first positive culture within three months post-surgery. A minimum of six
186 months follow-up was available for 95 cats, of these 25 (26.3%) had a first positive
187 culture within six months post-surgery. A minimum of one year follow-up was available
188 for 68 cats. Of these 28 (41.2%) had a first positive culture within one year post

189 surgery. Of the first positive cultures returned within the first year post surgery, 23
190 (82.1%) were single isolates and 5 (17.9%) were mixed cultures of two isolates. The
191 isolates identified in single organism cultures are summarised in Table 3.

192 Cats with a positive pre-operative culture were significantly more likely to have a
193 positive post-operative culture within six months post-operatively ($p=0.026$ OR 4.09 CI
194 1.18-14.18) than those with a negative pre-operative culture in multivariable analyses.
195 This was not statistically significant within one ($p=0.787$), three ($p=0.935$) or 12 ($p=0.328$)
196 months post-surgery. Of the 14 cats with a positive pre-operative culture six (42.9%)
197 returned a positive culture within one year post-operatively, in four cases (66.7%) the
198 same isolate was identified.

199 In multivariable analyses cats with higher end-anaesthetic rectal temperatures were
200 significantly less likely to return a positive culture by one month ($p=0.010$ OR 0.404 CI
201 0.203-0.803) and three months ($p=0.006$ OR 0.398 CI 0.205-0.772) post-surgery, but
202 not by six months ($p=0.121$) or 12 months ($p=0.555$) post-surgery.

203 Age, sex, breed, weight, condition score, perioperative antibiotic choice, reason for
204 device placement, outdoor access, pre-operative urea, creatinine, urine pH and
205 specific gravity, length of hospitalisation, anaesthetic and surgery were not
206 significantly correlated with likelihood of returning a positive culture at any time-point

207 when investigated with univariable analysis (all $p>0.1$) and were not included in the
208 multivariable model.

209 Use of TetraEDTA as part of a maintenance protocol from one-month post-surgery was
210 not significantly correlated with likelihood of returning a positive culture within three
211 months post-operatively and was not included in this multivariable model.

212 Site of implant and sex were included in the multivariable regression analysis but were
213 not statistically associated with likelihood of returning a positive culture at any time-
214 point ($p>0.05$).

215 Of the 28 cats returning positive urine cultures within 12 months post-surgery, 10
216 (35.7%) were asymptomatic with no noted device complications, two (7.1%) were
217 asymptomatic with device obstruction diagnosed on imaging, 11 (39.3%) had lower
218 urinary tract signs and/or secondary pyelonephritis and five (17.9%) had transient
219 lower urinary tract signs, which were at times absent in the presence of bacteriuria.

220 11 cats with a positive culture had their implants removed or replaced, of these seven
221 (63.6%) had either transient or perpetual clinical signs and four (36.4%) were
222 asymptomatic. 17 cats with a positive culture did not have their devices removed, of
223 these nine (52.9%) had transient or perpetual clinical signs and eight (47.1%) were

224 asymptomatic. There was no significant association between presence of clinical signs
225 and likelihood of device removal in cats with bacteriuria ($p=0.539$).

226 In total 26 devices became obstructed over the period studied in 22 cats (18.6%): 18
227 (15.3%) due to device mineralisation, seven (5.9%) due to a catheter kink, and one
228 (0.8%) due to a blood clot. Of the 18 devices (17 cats) with mineralised obstructions
229 nine were revised or replaced, three cats were euthanased and five declined surgical
230 intervention due to minimal or absent clinical signs.

231 Nineteen (16.1%) of the devices were removed or replaced, nine (47.4%) due to device
232 obstruction, 10 (52.6%) due to persistent infection that could not be eliminated with
233 antibiotics and one (5.3%) due to a combination of infection and obstruction.

234 Cats culturing positive for *E.coli* at any time point ($p=0.008$ OR 4.542 CI 1.485 -13.89)
235 were significantly more likely to have their implant removed or replaced when
236 investigated with multivariable analysis. This was not significant for cats culturing
237 positive for *P. aeruginosa* ($p=0.09$) or *E. faecalis* ($p=0.059$) when investigated with
238 multivariable analysis, or cats with positive cultures pre-operatively or within any time
239 point post-operatively when investigated with univariable analysis ($p>0.1$).

240 Cats with a positive urine culture pre-operatively or within any time-point post-
241 operatively were not significantly more likely to develop a device occlusion than cats
242 without a positive culture at any time-point when investigated with univariable
243 analysis ($p < 0.1$). However cats with *P. aeruginosa* cultured at any time-point post-
244 operatively were significantly more likely to develop an implant obstruction ($p = 0.033$
245 OR 5.0 CI 1.138-21.98), this was not significant for *E. coli* ($p = 0.583$) or *E. faecalis*
246 ($p = 0.532$) positive cultures when investigated with multivariable analysis.

247 Discussion

248 This study presents the results from a large number of urine cultures collected from
249 cats with SUB systems placed in a UK referral centre over an eight year period. Positive
250 cultures were obtained from 41.2% of the cats at some time point within twelve-
251 months post-operatively, with 8.5% returning a positive culture within one-month
252 post-operatively.

253 A previous paper by Kopecny et al (2019) reported 25% of cats returned a positive
254 post-operative urine culture, however it is difficult to compare these results with those
255 reported here as the population studied included cats with ureteral stent placement,
256 the time points assessed were different and only six of 48 samples were collected via
257 sterile subcutaneous port aspiration. It is also interesting to note that this paper

258 reported only 2.1% of cats returned a positive pre-operative culture compared to
259 12.7% of the cases reviewed here, 0% of the cases reported by Wolff et al (2016) and
260 25% of the cases reported by Berent et al (2018). The latter papers reported a post-
261 operative positive culture rate of 21% within ten days post operatively (7) and 24% at
262 any time point (5), however again direct comparison is challenging due to the
263 differences in data handling and proportion of cases lost to follow-up. Discrepancies
264 may also reflect difference in sampling methods or previous management of the cats
265 presented for surgery.

266 The most commonly cultured isolate in this study, both pre-operatively and at any time
267 point post-surgery, was *E.coli*. Berent et al (2018) also reported *E.coli* as the
268 predominant isolate cultured from pre-operative urine collection, however, in both
269 that paper and Kopecny et al (2019) *E. faecalis* was reported as the most common
270 isolate cultured from post-operative samples. *E.coli* and *E. faecalis* are both
271 commensals of the feline gastrointestinal tract (10) and thus the most likely
272 mechanism of urinary tract infection is ascending colonisation by pre-existing enteric
273 microflora. In this study, *P. aeruginosa* was the second most commonly cultured
274 organism at one month and joint second most commonly cultured organism (in
275 addition to *E. faecalis*) at 12-months post-surgery. *P. aeruginosa* is a ubiquitous
276 environmental organism, commonly implicated in opportunistic nosocomial infections

277 (11), in this study culturing *P. aeruginosa* at any time point was significantly associated
278 with risk of device obstruction, although not significantly associated with device
279 removal. This discrepancy is likely due to clients declining removal due to financial
280 constraints, or because the native ureter(s) had regained patency and the infection
281 was subclinical or resulting in only mild clinical signs.

282 The only factors identified as predictive of post-operative positive culture were
283 positive pre-operative culture and lower end-anaesthetic rectal temperature. In this
284 cohort four of the 14 cats with a positive pre-surgical urine culture cultured positive for
285 the same isolate within 176 days post-surgery. Berent et al (2018) also demonstrated
286 that pre-operative bacteriuria was significantly associated with post-operative
287 bacteriuria, however in this study it is not clear what proportion of cats returned the
288 same isolate at both cultures. Perioperative hypothermia has not previously been
289 investigated as a risk factor for post-operative implant infection, Beal et al (2000)
290 assessed the effect of hypothermia on surgical site infection rates in dogs and cats, and
291 found no significant relationship, however this paper reported incision infections,
292 rather than implant associated infections.

293 In human patients perioperative hypothermia has been shown to be associated with
294 surgical wound infections. It is thought that this occurs due to induction of peripheral

295 vasoconstriction leading to reduced tissue oxygenation and subsequent impaired
296 chemotaxis, phagocytosis, and antibody production (13). In one study of human
297 patients undergoing colorectal surgery, core temperature at end of surgery was highly
298 correlated with wound infection up to two weeks post-surgery, with 19% incidence in
299 the hypothermic group compared to 6% in the normothermic group (14). In our study
300 hypothermia was associated with increased risk of bacteriuria up to three months
301 post-surgery, it is possible that later positive cultures represent initial false negative
302 cultures or subclinical cases who missed the one month recheck.

303 In this study *E.coli* was associated with need for device removal/replacement. This
304 finding highlights the clinical significance of bacterial colonisation in these cases. In the
305 cases reviewed here the most common reason for implant removal/replacement was
306 bacterial infection. This is in contrast to previous work which reported mineralisation
307 to be the most frequent reason for device exchange, causing occlusion in 24.2% of
308 devices (5). In our study population occlusion due to device mineralisation occurred in
309 only 15.3% of cases and led to device removal or replacement in only 8.5% of cats. The
310 discrepancy in these figures may be attributable to multiple factors, such as
311 differences in diet and water mineralisation levels.

312 In this population clinical signs were seen in 57.2% of cats with positive urine cultures,
313 but only 39.3% of cats showed persistent lower urinary tract signs or pyelonephritis,
314 suggesting that many cats with positive urine cultures after device placement are not
315 clinically affected, i.e. have subclinical bacteriuria. This reflects previous work by
316 Berent et al (2018) (5) in which only 62.5% of persistently affected cats had clinical
317 signs suggestive of a urinary tract infection. The optimal way to manage cats with SUB
318 implants and positive urine cultures without lower urinary tract signs remains a topic
319 of debate. In this study the presence of any lower urinary tract signs was not
320 associated with an increased likelihood of device removal, but for some of these cats
321 the clinical signs were only transient and overall case numbers for cats with positive
322 urine cultures were small. Further work to investigate the optimal management
323 strategies for cats with SUB implants and positive urine cultures, both symptomatic
324 and subclinical, are warranted.

325 Our post-operative maintenance protocol was changed in February 2019 to include
326 instillation of TetraEDTA into the devices following sample collection. In this
327 population there was no significant difference in likelihood of returning a positive
328 culture at three months post-operatively for cats receiving routine TetraEDTA flushing
329 compared to those which did not. Correlation with positive culture at later time-points
330 post-surgery could not be investigated in this population as timing of data collection

331 meant follow-up length was limited in cats receiving the updated protocol; however,
332 this is an area which warrants further investigation.

333 The main limitations of this study are attributable to its retrospective design, with
334 available follow-up and pre-operative details variable. Although revisits were advised
335 at standardised intervals many patients did not revisit when advised. Additionally
336 cultures reported here may represent either clinical urinary tract infections or
337 subclinical bacteriuria. Although an attempt has been made to retrospectively classify
338 the cats as asymptomatic or otherwise, this distinction was made by reviewing
339 historical data which may not have been complete. Equally categories were not always
340 clearly defined, e.g. some cases showed transient clinical signs which waxed and
341 waned independent of treatment whereas others had persistent signs.

342 In conclusion post-operative bacteriuria occurred at least once within 12 months post-
343 operatively in 41.2% of cats and was a risk factor for device removal/replacement.
344 Both peri-operative hypothermia and post-operative positive culture were predictive
345 of post-operative positive culture and this should be taken into consideration when
346 managing these cases.

347 Statements

348 The authors would like to thank Dr. Yu-Mei Chang for her assistance with this project.

349 The authors have no conflict of interest to declare.

350 The authors received no financial support for the research, authorship, and/or
351 publication of this article.

352 This work involved the use of non-experimental animals only (including owned or
353 unowned animals and data from prospective or retrospective studies). Established
354 internationally recognised high standards ('best practice') of individual veterinary
355 clinical patient care were followed. Ethical approval was granted by the RVC ethical
356 approval board, submission reference SR2017-1364

357 Informed consent (either verbal or written) was obtained from the owner or legal
358 custodian of all animal(s) described in this work (either experimental or non-
359 experimental animals) for the procedure(s) undertaken

360 References

361 1 Kyles AE, Hardie EM, Wooden BG, et al. Clinical, clinicopathologic, radiographic and
362 ultrasonographic abnormalities in cats with ureteral calculi: 163 cases (1984–2002).
363 JAVMA 2005; 226: 932–936.

364 2 McLouglin MA, Bjorling DE. Ureters. In: Slatter D, editor. Textbook of small animal
365 surgery. 3rd edition. Philadelphia: WB Saunders; 2003. p. 1619–28.

366 3 Wen, J., Frøkiær, J., Jørgensen, T. and Djurhuus, J., 1999. Obstructive nephropathy:
367 an update of the experimental research. *Urological Research*, 27(1), pp.29-39.

368 4 Berent A, Weisse C. [Internet]. *Norfolkvetproducts.com*. 2020 [cited 24 January
369 2020]. Available from:
370 https://norfolkvetproducts.com/PDF/SUB/SUB2_Surgical_Guide_2018-03-email.pdf

371 5 Berent A, Weisse C, Bagley D, Lamb K. Use of a subcutaneous ureteral bypass device
372 for treatment of benign ureteral obstruction in cats: 174 ureters in 134 cats (2009–
373 2015). *Journal of the American Veterinary Medical Association*. 2018;253(10):1309-
374 1327.

375 6 Deroy C, Rossetti D, Ragetly G, Hernandez J, Poncet C. Comparison between double-
376 pigtail ureteral stents and ureteral bypass devices for treatment of ureterolithiasis in
377 cats. *Journal of the American Veterinary Medical Association*. 2017;251(4):429-437.

378 7 Wolff EDS, Dorsch R, Knebel J, et al. Initial outcomes and complications of the
379 subcutaneous ureteral bypass procedure at two university hospitals (2012-2015).
380 Denver (CO): Abstract ACVIM 2016

381 8 Kopecny L, Palm C, Drobatz K, Balsa I, Culp W. Risk factors for positive urine cultures
382 in cats with subcutaneous ureteral bypass and ureteral stents (2010-2016). Journal of
383 Veterinary Internal Medicine. 2018;33(1):178-183.

384 9 CLSI. Performance standards for antimicrobial disk and dilution susceptibility tests for
385 bacteria isolated from animals; approved standard-third edition. CLSI document M31-
386 A3. Wayne, PA, Clinical Laboratory Standards Institute 2008

387 10 KuKanich K, Lubbers B. Review of Enterococci Isolated from Canine and Feline Urine
388 Specimens from 2006 to 2011. Journal of the American Animal Hospital Association.
389 2015;51(3):148-154.

390 11 Grosso-Becerra M, Santos-Medellín C, González-Valdez A, Méndez J, Delgado G,
391 Morales-Espinosa R et al. Pseudomonas aeruginosa clinical and environmental isolates
392 constitute a single population with high phenotypic diversity. BMC Genomics.
393 2014;15(1):318.

394 12 Beal M, Brown D, Shofer F. The Effects of Perioperative Hypothermia and the
395 Duration of Anesthesia on Postoperative Wound Infection Rate in Clean Wounds: A
396 Retrospective Study. Veterinary Surgery. 2000;29(2):123-127.

397 13 Van Oss, C., Absolom, D., Moore, L., Park, B., Humbert, J. Effect of temperature on

398 the chemotaxis, phagocytic engulfment, digestion and O₂ consumption of human
399 polymorphonuclear leukocytes. *J Reticuloendothel Soc.*1980; 27:561-565.

400 14 Kurz, A., Sessler, D., Lenhardt, R. Perioperative normothermia to reduce the
401 incidence of surgical-wound infection and shorten hospitalization. Study of Wound
402 Infection and Temperature Group. *N Engl J Med.* 1996; 334:1209-1215.