



Biomarker changes with systolic anterior motion of the mitral valve in cats with hypertrophic cardiomyopathy

Joonbum Seo¹ | Jessie Rosie Payne² | Jose Novo Matos^{1,3} |
Wesley Wynne Fong¹ | David J. Connolly¹ | Virginia Luis Fuentes¹

¹Clinical Science and Services, Royal Veterinary College, Hertfordshire, United Kingdom

²Langford Vets Small Animal Referral Hospital, University of Bristol, Bristol, United Kingdom

³Department of Veterinary Medicine, University of Cambridge, Cambridge, United Kingdom

Correspondence

Joonbum Seo, Clinical Science and Services, Royal Veterinary College, Hawkshead Lane, North Mymms, Hertfordshire, United Kingdom.
Email: jseo7@rvc.ac.uk

Abstract

Background: N-terminal pro B-type natriuretic peptide (NT-proBNP) and cardiac troponin-I (cTnI) are biomarkers commonly evaluated in cats with suspected heart disease. Many cats with hypertrophic cardiomyopathy (HCM) have systolic anterior motion of the mitral valve (SAM), but its influence on circulating NT-proBNP or cTnI concentrations is currently unknown.

Hypothesis/Objectives: Cats with HCM and SAM (HCM^{SAM+}) have higher NT-proBNP and cTnI concentrations than do cats with HCM but without SAM (HCM^{SAM-}).

Animals: One hundred forty cats with HCM: 70 with SAM and 70 without SAM.

Methods: Retrospective case-to-case study. Cats were recruited if diagnosed with HCM by echocardiography and results were available for NT-proBNP or cTnI concentrations or both. Cats with SAM were matched to those without SAM for clinical presentation, left atrial (LA) size and left ventricular (LV) fractional shortening.

Results: A total of 119 NT-proBNP and 123 cTnI results were available. The HCM^{SAM+} cats had higher median concentrations than did HCM^{SAM-} cats for NT-proBNP (729 pmol/L; interquartile range [IQR], 275-1467 versus 65 pmol/L; IQR, 25-271; $P < .001$) and cTnI (0.27 ng/mL; IQR, 0.10-0.81 versus 0.07 ng/mL; IQR, 0.01-0.43; $P = .002$). In general linear models for both NT-proBNP and cTnI, the independent explanatory variables were SAM, congestive heart failure, maximal LV wall thickness, and LA size.

Conclusions and Clinical Importance: For cats with HCM and equivalent LA size and LV systolic function, those with SAM had higher NT-proBNP and cTnI concentrations than did those without SAM. Presence of SAM should be considered when interpreting biomarker concentrations in cats with HCM.

Abbreviations: ATE, aortic thromboembolism; CHF, congestive heart failure; cTnI, cardiac troponin-I; DLVOTO, dynamic left ventricular outflow tract obstruction; FS%, left ventricular fractional shortening percentage; HCM, hypertrophic cardiomyopathy; HCM^{SAM+}, cats with hypertrophic cardiomyopathy and systolic anterior motion of the mitral valve; HCM^{SAM-}, cats with hypertrophic cardiomyopathy but without systolic anterior motion of the mitral valve; LA : Ao, left atrium-to-aortic ratio; LAD Max, maximal left atrial diameter; LVIDd, left ventricular internal diameter in end-diastole; LVIDs, left ventricular internal diameter in end-systole; LVOT Vmax, maximal left ventricular outflow tract velocity; LVWT Max, maximal end-diastolic left ventricular wall thickness; NT-proBNP, N-terminal pro B-type natriuretic hormone; POC, point-of-care; QMHA, Queen Mother Hospital for Animals; SABP, systolic arterial blood pressure; SAM, systolic anterior motion of the mitral valve.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

© 2020 The Authors. *Journal of Veterinary Internal Medicine* published by Wiley Periodicals, Inc. on behalf of the American College of Veterinary Internal Medicine.

KEYWORDS

cardiology, cat, natriuretic peptide, obstructive cardiomyopathy, troponin

1 | INTRODUCTION

Cardiac biomarkers increasingly have been used in cats with heart disease. N-terminal pro B-type natriuretic hormone (NT-proBNP) is a biomarker of myocardial stress and stretch. Its application as a cardiac biomarker includes screening for preclinical cardiomyopathy,¹⁻⁴ assessing disease severity,⁵ and differentiating cardiogenic and non-cardiogenic causes in cats with dyspnea.⁴⁻⁸ It has been proposed as being particularly useful for general practitioners without access to echocardiography.⁹ Cardiac troponin-I (cTnI) is a biomarker of myocardial injury. Its use as a cardiac biomarker includes assessing cardiomyopathy severity,¹⁰ differentiating cardiogenic and noncardiac causes in cats with dyspnea,¹¹ and providing additional prognostic information independent of left atrial size and function, and left ventricular systolic function.^{12,13}

Hypertrophic cardiomyopathy (HCM) is a disease with considerable morphological and functional variation. For example, there are differences in segmental versus global concentric hypertrophy,¹⁴ presence or absence of systolic anterior motion of the mitral valve (SAM)^{15,16} and mid-ventricular obstruction,¹⁷ and papillary muscle morphology varies.¹⁵ Of these, SAM is encountered frequently, with a reported prevalence of approximately 30% in cats with HCM phenotype.^{18,19} Systolic anterior motion of the mitral valve is associated with more severe left ventricular concentric hypertrophy in cats, but whether SAM is an effect or a cause of the hypertrophy is unknown.¹⁵ Cardiac magnetic resonance imaging studies suggest that abnormalities of both left ventricular geometry and the mitral valve apparatus likely contribute to the origin of SAM in people.²⁰ Similar abnormalities likely occur in affected cats.¹⁵

Although the primary stimulus for release of NT-proBNP is myocardial stretch, other conditions that result in increased wall stress or ventricular hypertrophy have been found to increase plasma concentrations of NT-proBNP.²¹⁻²⁵ Dynamic left ventricular outflow tract obstruction (DLVOTO) would be expected to increase myocardial wall stress, and therefore increase NT-proBNP.^{26,27} Furthermore, DLVOTO has been associated with an increased risk of myocardial ischemia in people with HCM, which would be expected to result in increased cTnI concentrations.^{26,28} Many studies have shown that NT-proBNP and cTnI concentrations are increased in cats with HCM phenotype,^{2,5,10,12,29} but whether or not the presence of SAM further increases these biomarker concentrations currently is unknown.

2 | AIMS/OBJECTIVES

The first aim of our study was to compare circulating concentrations of NT-proBNP and cTnI in cats with HCM phenotype and SAM as compared to cats with HCM phenotype but without SAM. The second aim was to create multivariable models to identify factors associated

with variation in NT-proBNP and cTnI concentrations. We hypothesized that cats with SAM would have higher NT-proBNP and cTnI concentrations than cats at a similar stage of HCM without SAM, and that this relationship would be independent of other factors when assessed in a multivariable model.

3 | MATERIALS AND METHODS

3.1 | Study design

This study is a retrospective case-to-case study design. Ethical approval was provided by the Royal Veterinary College (URN: SR2019-0245). Electronic medical records between December 2011 and May 2018 from the Queen Mother Hospital for Small Animals (QMHA) and a cardiac screening study at rehoming centers for cats (the CatScan study¹⁸) were reviewed for cats diagnosed with HCM and with NT-proBNP, cTnI, or both measured. The diagnosis of HCM was defined as left ventricular hypertrophy at any region measuring ≥ 6 mm on 2-dimensional (2D) imaging.¹⁶

For each case, medical records corresponding with the time of biomarker measurement were reviewed. The signalment, medical history, physical examination findings, systemic arterial blood pressure (SABP) measurements using Doppler technique, presence of congestive heart failure (CHF), arrhythmias or aortic thromboembolism (ATE), and current medications were recorded. Where the intensity of a murmur varied, the highest murmur grade was recorded. Cats were excluded if they were diagnosed with dehydration, hyperthyroidism, SABP ≥ 180 mm Hg,³⁰ diabetes mellitus, cardiomyopathies with phenotype other than HCM, cardiac neoplasia, congenital heart disease, anemia (packed cell volume $< 20\%$), or had incomplete clinical records. Cats with serum creatinine concentrations ≥ 2.9 mg/dL (≥ 251 $\mu\text{mol/L}$) or ≥ 1.6 mg/dL (≥ 140 $\mu\text{mol/L}$) without SABP measurement also were excluded.³⁰

Echocardiograms matching the dates of the biomarker measurement were reviewed by a single observer (J.S.). All echocardiographic measurements were taken over 3 different cardiac cycles and averaged. After confirming the diagnosis, the presence of SAM was assessed with frame-by-frame review of the 2D imaging of a right parasternal 5-chamber view, optimized for the left ventricular outflow tract (LVOT).¹⁵ Systolic anterior motion of the mitral valve was defined as anterior motion of either septal or both mitral valve leaflets during systole toward the LVOT on review of the 2D cine loop.³¹ Other echocardiographic variables were measured by the same investigator (J.S.) as described in Table 1. These included: left atrium to aortic ratio (LA : Ao),³² maximal left atrial diameter (LAD Max),^{33,34} maximal left ventricular wall thickness (LVWT Max),³⁴ left ventricular internal diameter in end-diastole (LVIdd) and end-systole (LVIdS),³⁵ left ventricular fractional shortening (FS%),³⁵ and maximal LVOT velocity (LVOT Vmax).¹⁶

Dynamic left ventricular outflow tract obstruction was defined as LVOT Vmax ≥ 2.5 m/s in the absence of echocardiographic evidence of fixed aortic stenosis.^{15,16} Regional wall hypokinesis was defined as asynchronous motion or minimal excursion of a myocardial segment compared with the adjacent segment.^{12,36}

3.2 | Matching process

The case-to-case groups (HCM^{SAM+} and HCM^{SAM-}) were created starting with all cats without SAM (HCM^{SAM-}), and calculating the proportion of cats with CHF, ATE and arrhythmias, and the mean or

median of the echocardiographic variables (LAD Max, LA : Ao, LVIDd, LVIDs, and FS%). Cats with SAM were introduced 1 cat at a time to a HCM^{SAM+} group, to create a group with characteristics matching the HCM^{SAM-} group. As described in the introduction, LVWT Max was expected to be higher in the HCM^{SAM+} group, and no attempt was made to match this variable between the groups.¹⁵

3.3 | NT-proBNP and cTnl measurements

An EDTA blood sample was collected for a second-generation NT-proBNP assay. The NT-proBNP samples were processed by

TABLE 1 Echocardiographic variables of interest

Echocardiographic variable	View and timing	Measurement
LA : Ao ³²	View: RPSAX at the level of the aortic valve Imaging modality: 2D Imaging Timing: Beginning of diastole, the first frame of aortic valve closure	Ao: From the blood-tissue interface at the midpoint of the right aortic sinus to the commissure between the noncoronary and left coronary aortic cusps LA: Extension of the aortic line to the blood-tissue interface of the left atrial wall, immediately lateral to the pulmonary vein
LAD Max ^{33,34}	View: RPLAX4ch Imaging modality: 2D Imaging Timing: Beginning of diastole, the last frame before the mitral valve opening	Mid-interatrial septum to the leading edge of the pericardium, bisecting the left atrium parallel to the mitral valve annulus
LVWT Max ³⁴	View: RPLAX4ch, RPLAX5ch, and RPSAX at the level of the papillary muscles Imaging modality: 2D Imaging Timing: End-diastole, the last frame before the aortic valve opens (RPLAX5ch), the first frame after the mitral valve closes (RPLAX4ch), or when the left ventricular internal diameter was the largest (RPSAX)	Leading edge technique avoiding the papillary muscles or false tendon attachments Only the average of 3 measurements from the area that measures the maximal thickness is used
LVIDd ³⁵	View: RPSAX at the level of the papillary muscles Imaging modality: 2D Imaging Timing: End-diastole	Leading edge technique bisecting the left ventricle in end-diastole
LVIDs ³⁵	View: RPSAX at the level of the papillary muscles Imaging modality: 2D Imaging Timing: End-systole	Leading edge technique bisecting the left ventricle in end-systole
FS% ³⁵	-	Calculated using (LVIDd-LVIDs)/LVIDd
LVOT Vmax ¹⁶	View: LA5ch with the cursor transecting the left ventricular outflow tract. Imaging modality: Spectral Doppler (either pulse wave or continuous wave) imaging Timing: Systole, when the modal velocity was the greatest	Maximal modal velocity in systole was recorded. The measurement was not recorded when there was inadequate cursor alignment, absence of a dynamic profile, or contamination of the mitral regurgitation
DLVOTO ^{15,16}	-	Defined as LVOT Vmax ≥ 2.5 m/s
Regional wall hypokinesis ^{12,36}	View: Overview of the RPLAX4ch, RPLAX5ch, RPSAX, LA4ch, LA5ch Imaging modality: 2D Imaging Timing: Across the entire cardiac cycle	Subjective recognition of a segment with asynchronous motion to the adjacent segment or minimal excursion on 2D imaging

Note: All echocardiographic measurements in this study were taken over 3 different cardiac cycles and averaged.

Abbreviations: DLVOTO, dynamic left ventricular outflow tract obstruction; FS%, left ventricular fractional shortening; LA:Ao, left atrium to aortic ratio; LA4ch, left apical 4-chamber view, LA5ch, and 5-chamber view; LAD Max, maximal left atrial diameter; LVIDd, left ventricular internal diameter in diastole, LVIDs, and systole; LVOT Vmax, maximal left ventricular outflow tract velocity; LVWT Max, maximal end-diastolic left ventricular wall thickness; RPLAX4ch, right parasternal long axis 4-chamber view, RPLAX5ch, and 5-chamber view; RPSAX, right parasternal short axis view.

2 different methods, both after the instructions of the reference laboratory (IDEXX Veterinary Laboratories, Wetherby, Workshire, United Kingdom): For the samples taken at the QMHA, the samples were centrifuged at 800g for 3 minutes and plasma was transferred to a separate plain tube and refrigerated at 4°C. The samples then were transported within 24 hours to the reference laboratory where the assays were performed. For samples collected at the rehoming centers, samples were centrifuged at 4000 rpm for 15 minutes. Plasma was separated and immediately frozen at -20°C. Batches of frozen samples were transferred on dry ice to the reference laboratory and underwent batch analysis.

Cardiac troponin-I was analyzed using 2 different analyzers: For all samples from the rehoming centers and for some from the QMHA, a blood sample was collected in a plain tube and processed in an identical manner as samples for the NT-proBNP assay. These samples then were analyzed at the reference laboratory. Some cats from the QMHA had a point-of-care (POC) high sensitivity cTnI measurement performed with a handheld analyzer (VetScan i-STAT Analyzer, Abaxis, Abbott Point of Care Inc, Union City, California). Either whole blood in plain tubes or heparinized whole blood samples were used immediately or within 10 minutes after collection, respectively.

The lower and upper limits of detection for NT-proBNP were 24 and 1500 pmol/L, respectively. Results reported as <24 or >1500 pmol/L were entered as 23 or 1501 pmol/L, respectively, for statistical analysis. The high sensitivity cTnI assay from the reference laboratory had a lower limit of detection of 0.01 ng/mL and no upper limit. The POC analyzer had the same lower limit of detection but an upper limit of 50 ng/mL. Results reported as <0.01 ng/mL were entered as 0.009 ng/mL. The POC results >50 ng/mL were entered as 51 ng/mL.

4 | STATISTICAL ANALYSIS

Commercial software programs were used for statistical analysis (IBM SPSS Statistics Version 24) and generating graphs (GraphPad Prism 7 Version 7.0d). Normality of the continuous variables was tested visually and by a Shapiro-Wilk test. Normally distributed data are presented as mean (\pm SD) and non-normally distributed data as median (interquartile range, IQR). Continuous variables were compared using either an independent *t* test or Mann-Whitney test. Categorical demographic variables were compared by a Pearson chi-squared test, with posthoc analysis done by analyzing the standardized residuals. Significance was set at $P < .05$.

The association between the dependent variables (NT-proBNP and cTnI) and others was tested first by univariable analysis, by either Pearson's or Spearman's test for continuous variables, and either an independent *t* test or Mann-Whitney test for categorical variables. Significantly associated variables with $P < .25$ then were selected for multivariable analysis.

Multivariable analysis was performed by constructing a separate general linear model for NT-proBNP and cTnI. Both models were created in a backward stepwise manner, starting with all significant

variables from the univariable analysis ($P < .25$) and then removing non-significant variables 1 at a time until only significant variables remained ($P < .05$). Variables were log-transformed if the distribution was severely skewed, and the residuals of each model were visually inspected by quantile plots, ensuring the assumption of normal distribution of linear models. Collinearity statistics were checked and multicollinearity was eliminated.

5 | RESULTS

A total of 210 cats met the initial inclusion criteria. Of these, 70 cats subsequently were excluded because of anemia (1), limited echocardiographic study (8), lymphoma (1), incomplete medical records (23), hyperthyroidism (3), diabetes mellitus (2), serum creatinine concentration ≥ 2.9 mg/dL (8), serum creatinine concentration ≥ 1.6 mg/dL with unavailable blood pressure measurement (4), and any additional cats after matching the 2 groups (20). This process resulted in a final number of 140 enrolled cats: 40 cats from rehoming centers and 100 cats from the QMHA. In total, 119 measurements were available for NT-proBNP, and 123 measurements for cTnI. These cats were divided into HCM^{SAM+} (70 cats) and HCM^{SAM-} groups (70 cats; Table 2).

No significant differences in breed, sex, medications, or the proportion of cats with CHF, ATE, or arrhythmias were found between the groups. Cats in the HCM^{SAM+} group were significantly younger ($P < .001$) with lower body weight ($P < .001$) than cats in the HCM^{SAM-} group. A significant difference in presenting complaints was found between the 2 groups, with most HCM^{SAM+} cats being presented for a heart murmur ($P < .001$) and most HCM^{SAM-} cats being presented for HCM screening ($P < .001$). Additional demographic information is outlined in Table 2.

Based on the study design, no differences were found in LAD Max, LA : Ao, LVIDd, LVIDs, and FS%. The LVWT Max was higher in the HCM^{SAM+} cats ($P < .001$). Forty-nine of 65 HCM^{SAM+} cats (68.4%) had DLVOTO at the time of examination, with a mean LVOT Vmax of 3.3 ± 1.2 m/s. Additional echocardiographic data are presented in Table 3.

Plasma NT-proBNP concentrations were significantly higher in the HCM^{SAM+} group (729 pmol/L [IQR, 275-1467] versus 65 pmol/L [IQR, 25-271] in the HCM^{SAM-} group; $P < .001$; Figure 1). Circulating concentrations of cTnI also were significantly higher in the HCM^{SAM+} group (0.27 ng/mL [IQR, 0.10-0.81] versus 0.07 ng/mL [IQR, 0.01-0.43] in the HCM^{SAM-} group; $P = .002$). This increase was observed in both the non-DLVOTO and DLVOTO populations of the SAM group ($P = .3$ for NT-proBNP and $P = .6$ for cTnI comparing non-DLVOTO and DLVOTO subgroups).

Results of the univariable analysis are summarized in TableS1. Using these results, the following variables were tested in a general linear model to explain the variation of NT-proBNP: source population (QMHA versus rehoming centers), presenting complaint, age, body weight, body condition score, breed, SABP, murmur grade, gallop sound, sedation, medication, urea, creatinine, CHF, ATE, arrhythmias, SAM, DLVOTO, LVOT Vmax, LVIDs, FS%, LVWT Max, LAD Max, regional wall hypokinesis, and cTnI. Similarly, the following variables

TABLE 2 Demographic characteristics for the 140 cats with hypertrophic cardiomyopathy

	HCM ^{SAM+} (n = 70)	HCM ^{SAM-} (n = 70)	P value
Population			
CatScan	9 (12.9%)	31 (44.3%)	<.001
QMHA	61 (87.1%)	39 (55.7%)	
Age (years)	4.1 [2.4-7.0]	6.9 [4.0-10.6]	<.001
Weight (kg)	4.4 (±1.0)	5.2 (±1.2)	<.001
BCS (/9)	5.0 [4.0-5.0]	5.0 [5.0-6.4]	.332
Presenting complaint (%)			
Arrhythmias	0	1 (1.4%)	
Collapsing episodes	3 (4.3%)	2 (2.9%)	
Gait abnormality	4 (5.7%)	5 (7.1%)	
Murmur	32 (45.7%)	8 (11.4%)	
Respiratory signs	7 (10%)	9 (12.9%)	
HCM screening	13 (18.6%)	39 (55.7%)	
HCM reassessment	11 (15.7%)	6 (8.6%)	
Murmur (%)	66/68 (94.3%)	42/69 (60.9%)	<.001
Murmur grade	3 [3-4]	2 [0-3]	
Gallop sound (%)	3/26 (11.5%)	8/57 (11.4%)	1.000
Arrhythmias ^a	5 (7.1%)	8 (11.4%)	.562 ^a
Congestive heart failure ^a	8 (11.4%)	13 (18.6%)	.344 ^a
Aortic thromboembolism ^a	3 (4.3%)	3 (4.3%)	1.000 ^a
Sex			
Female (%)	25 (35.7%)	21 (30.0%)	.590
Neutered : Entire	22:3	18:3	
Male (%)	45 (64.3%)	49 (70.0%)	
Neutered : Entire	42:3	42:7	
Number of pedigree cats (%)			
Bengal	2 (2.9%)	2 (2.9%)	
British Short Hair	3 (4.3%)	2 (2.9%)	
Domestic Long Hair	5 (7.1%)	6 (8.6%)	
Domestic Medium Hair	2 (2.9%)	1 (1.4%)	
Domestic Short Hair	48 (68.6%)	43 (61.4%)	
European Short Hair	3 (4.3%)	0	
Maine Coon	0	1 (1.4%)	
Norwegian Forest Cat	1 (1.4%)	8 (11.4%)	
Persian	2 (2.9%)	1 (1.4%)	
Russian Blue/White	1 (1.4%)	2 (2.9%)	
Scottish Fold	1 (1.4%)	1 (1.4%)	
Siamese	0	2 (2.9%)	
Sphynx	2 (2.9%)	1 (1.4%)	
Doppler blood pressure	124.1 (±18.8) (n = 41)	129.7 (±21.6) (n = 39)	.224
Renal markers			
Urea (mg/dL)	27.2 [21.8-39.2] (n = 14)	28.3 [25.8-35.6] (n = 35)	.603
Creatinine (mg/dL)	1.52 (±0.28) (n = 16)	1.58 (±0.35) (n = 35)	.618
Medications (number of cats)			
Aspirin	0	1 (1.4%)	.512
Atenolol	1 (1.4%)	0	

(Continues)

TABLE 2 (Continued)

	HCM ^{SAM+} (n = 70)	HCM ^{SAM-} (n = 70)	P value
Benazepril	2 (2.9%)	2 (2.9%)	
Clopidogrel	3 (4.3%)	1 (1.4%)	
Diltiazem	1 (1.4%)	0	
Furosemide	4 (5.7%)	4 (5.7%)	
Pimobendan	0	1 (1.4%)	

Note: The cats were divided in 2 groups: Hypertrophic cardiomyopathy with systolic anterior motion of the mitral valve (HCM^{SAM+}) and hypertrophic cardiomyopathy without systolic anterior motion of the mitral valve (HCM^{SAM-}).

^aActively matched variables.

TABLE 3 Echocardiographic variables and cardiac biomarker results for the 140 cats with hypertrophic cardiomyopathy

	HCM ^{SAM+} (n = 70)	HCM ^{SAM-} (n = 70)	P value
SAM	70 (100%)	0	
Sedation	1 (1.4%)	2 (2.9%)	1.000
LVOT Vmax (m/s)	3.3 (±1.2) (n = 65)	1.1 (±0.5) (n = 29)	<.001
DLVOTO	49/65 (68.4%)	0/50 (0%)	<.001
LAD Max (mm) ^a	16.4 [14.5-17.9]	15.4 [13.7-17.4]	.163 ^a
LA : Ao ^a	1.3 [1.2-1.5]	1.3 [1.2-1.5]	.762 ^a
LVIDd (mm) ^a	13.9 [12.5-15.4]	14.2 [12.5-16.1]	.216 ^a
LVIDs (mm) ^a	5.0 [3.9-5.9]	5.6 [3.8-7.2]	.115 ^a
FS ^a	62.4 [57.6-71.4]	61.9 [50.2-69.9]	.253 ^a
Regional wall hypokinesis	5 (7.1%)	6 (8.6%)	1.000
LVWT Max (mm)	7.2 [6.6-8.2]	6.5 [6.1-6.8]	<.001
NT-proBNP (pmol/L)	729 [275-1467] (n = 61)	65 [25-271] (n = 58)	<.001
cTnl (ng/mL)	0.27 [0.10-0.81] (n = 59)	0.07 [0.01-0.43] (n = 64)	.002
POC cTnl	7/59 (11.9%)	7/64 (10.9%)	1.000

Abbreviations: DLVOTO, dynamic left ventricular outflow tract obstruction; FS%, fractional shortening %; LA : Ao, left atrium-to-aortic ratio; LAD Max, maximal left atrial diameter; LVIDd, LVIDs, left ventricular internal diameter in diastole and systole; LVOT Vmax, maximal left ventricular outflow tract velocity; LVWT Max, maximal left ventricular wall thickness; SAM, systolic anterior motion of the mitral valve.

^aActively matched variables.




FIGURE 1 Box and whisker plots graphs comparing plasma concentrations of A, NT-proBNP and B, cTnl in cats with hypertrophic cardiomyopathy with and without systolic anterior motion of the mitral valve (groups HCM^{SAM+} and HCM^{SAM-}, respectively). cTnl, cardiac troponin-I; HCM^{SAM+}/HCM^{SAM-}, cats with hypertrophic cardiomyopathy and with/without systolic anterior motion of the mitral valve; NT-proBNP, N-terminal pro B-type natriuretic peptide

TABLE 4 Two multivariable models with NT-proBNP and cTnI as dependent variables after log-transformation

1. log(NT-proBNP)			
Adjusted R ² = .524	B coefficient (95% CI)	Standardized B coefficient	P value
CHF	0.496 (0.209-0.783)	0.229	.001
SAM	0.583 (0.398-0.767)	0.448	<.001
LAD Max	0.050 (0.025-0.076)	0.268	<.001
LVWT Max	0.137 (0.038-0.236)	0.208	.007
2. log(cTnI)			
Adjusted R ² = .493	B coefficient (95% CI)	Standardized B coefficient	P value
CHF	1.060 (0.694-1.425)	0.407	<.001
SAM	0.436 (0.188-0.685)	0.237	.001
LAD Max	0.070 (0.038-0.102)	0.309	<.001
LVWT Max	0.169 (0.050-0.288)	0.196	.006

Abbreviations: CHF, congestive heart failure; cTnI, cardiac troponin-I; LAD Max, maximal left atrial diameter; LVWT Max, maximal left ventricular wall thickness; NT-proBNP, N-terminal pro B-type natriuretic peptide; SAM, systolic anterior motion of the mitral valve.

were tested for cTnI: source population, presenting complaint, body weight, body condition score, SABP, gallop sound, sedation, medication, urea, CHF, ATE, arrhythmias, SAM, DLVOTO, LVOT Vmax, LVIDs, FS%, LVWT Max, LAD Max, regional wall hypokinesis, POC cTnI, and NT-proBNP. Because of their skewed distribution, NT-proBNP and cTnI were log-transformed for further analysis.

Results of multivariable analysis are presented in Table 4. For both general linear models, LAD Max, CHF, SAM, and LVWT Max were the final independent variables with an adjusted R² of .52 for NT-proBNP and .49 for cTnI. The beta coefficient for SAM was higher than for LVWT Max in both models. Other variables including age, source population (QMHA versus rehoming centers), POC cTnI, creatinine, urea, SABP, and presence of regional wall hypokinesis were not significant in either model. Visual inspection of the residuals was consistent with good model fit.

6 | DISCUSSION

We compared cardiac biomarker concentrations in cats with HCM and SAM to those without SAM that were otherwise similar for sex, breed, left atrial and left ventricular size, clinical signs, blood pressure, and renal markers. Concentrations of circulating NT-proBNP and cTnI were significantly higher in the cats with SAM, and multivariable analysis confirmed that presence of SAM was associated with higher NT-proBNP and cTnI concentrations, independent of other factors.

Left atrial size is widely considered 1 of the most important prognostic factors in cats with cardiomyopathy,^{31,36,37} and although NT-proBNP also has been shown to have prognostic value, at least 1 study has shown left atrial size to be a more important indicator.¹² In situations where echocardiography is unavailable, NT-proBNP concentrations sometimes are measured as a means of screening for more advanced disease. We were not able to determine whether the higher NT-proBNP concentrations in the cats with SAM corresponded with a

worse prognosis, but we were able to show that cats with SAM can have high NT-proBNP even with normal left atrial size. Approximately one-third of cats with HCM phenotype have SAM on presentation,^{18,19} and our findings suggest that the presence or absence of SAM should be taken into consideration when interpreting NT-proBNP concentrations.

Other independent explanatory variables in our final multivariable model of NT-proBNP were LAD Max, CHF, and LVWT Max. These findings are expected based on previous studies that showed increased NT-proBNP concentrations with ventricular hypertrophy, left atrial enlargement, and CHF.^{1-9,25} Renal disease and systemic hypertension are other known conditions associated with increased NT-proBNP concentrations in cats.³⁰ These variables were not significant in our final multivariable model, likely because of our study design in which cats with systemic hypertension and azotemia were excluded.

Cardiac troponin-I also was higher in cats with SAM. In people, SAM is thought to confer greater risk of myocardial ischemia as a result of increased myocardial oxygen demand and decreased coronary perfusion, potentially associated with worsening hypertrophy, microvascular dysfunction,^{26,38,39} and arteriosclerosis of intramural small coronary arteries.^{38,39} Similar mechanisms are hypothesized in cats with HCM and SAM. The higher cTnI concentrations in cats with HCM have been associated with shorter survival times,^{12,13} but studies to date have failed to show any association between the presence of SAM and increased cardiac morbidity or mortality in cats.^{16,19,36,40,41} These observations contrast with those of human medical literature in which SAM is considered a negative prognostic factor in HCM.²⁶ In our study and in other studies, cats with HCM and SAM were younger than cats without SAM. The recent REVEAL study showed similar survival times between cats with obstructive and nonobstructive HCM, but cats with obstructive HCM also were younger than cats with nonobstructive HCM.¹⁹ Other significant independent explanatory variables in our study were LAD Max, CHF, and LVWT Max, findings that are similar to those of previous studies.^{10,11}

Systolic anterior motion of the mitral valve usually is associated with DLVOTO, which can result in increased left ventricular wall stress and ischemia.^{26,42} However, although the presence of SAM was a statistically significant explanatory variable for both NT-proBNP and cTnI in our study, neither LVOT Vmax nor the presence of DLVOTO was significant in our general linear models. We defined DLVOTO as LVOT velocities >2.5 m/s, although to confirm the site of dynamic obstruction, color Doppler imaging is necessary to show variance and aliasing in the LVOT and an eccentric jet of mitral regurgitation. Dynamic LVOT obstruction is known to be labile, and is affected by loading conditions and sympathetic tone.⁴³ Provocative maneuvers often are used to provoke a gradient across the LVOT in people with HCM,⁴⁴ and auditory stimulation sometimes is used in cats with HCM for this purpose. It is difficult to standardize echocardiographic conditions in cats, and the echocardiographic examination itself could be considered a provocative maneuver, with variable effects in different cats. The magnitude of LVOT gradient that should be considered clinically relevant in cats with HCM has not been established, nor whether or not this conclusion should be based on a resting or provoked gradient. It is not clear why SAM has a closer association with NT-proBNP and cTnI concentrations than with presence of DLVOTO or LVOT Vmax. It might reflect greater ease of recording SAM compared with DLVOTO or LVOT Vmax, or that SAM is more consistently present in cats with intermittent LVOT obstruction.

Our study had a number of limitations, many of which were a consequence of its retrospective nature. First, the NT-proBNP samples were processed by 2 different methods according to the instructions provided by the reference laboratory. However, no significant effect was seen when using the general linear model with the variable "population source." Similarly, cTnI concentrations were measured using 2 different analyzers, following the instructions provided by either the reference laboratory or the manufacturer of the POC cTnI analyzer. Although this POC cTnI assay has not been validated in cats, cTnI is well conserved across species and statistical analysis with or without these cats showed no effect of including this unvalidated assay in our results, and no significant effect of the POC cTnI was found in the final models. We elected to include the cats with cTnI analyzed on the POC assay because doing so resulted in better balancing of the 2 groups. The HCM^{SAM} + cats were younger and had higher LVWT Max. Systemic hypertension previously has been shown to result in left ventricular hypertrophy in cats.⁴⁵ Although strict exclusion criteria were followed, systemic blood pressure recordings were not available in all cats, and some cats with systemic hypertension inadvertently may have been included. However, SAM still had a significant influence on both NT-proBNP and cTnI concentrations independent of wall thickness, and the influence of age was not significant in the final multivariable model. Finally, no attempt was made to evaluate outcome. Doing so was not an aim of our study, and therefore we cannot determine whether increased NT-proBNP and cTnI

concentrations in cats with SAM are associated with a worse prognosis regardless of left atrial size.

7 | CONCLUSIONS

Cats with HCM and SAM had higher NT-proBNP and cTnI concentrations than did cats without SAM. The presence of SAM should be taken into consideration when interpreting NT-proBNP and cTnI results in cats with HCM phenotype.

ACKNOWLEDGMENT

The findings of this study were presented at the 2019 ACVIM Forum, Phoenix, AZ.

CONFLICT OF INTEREST DECLARATION

This study was not supported by any grant or other source of direct funding. Drs JR Payne and V Luis Fuentes previously received financial support from IDEXX Ltd for research on NT-proBNP and cTnI.

OFF-LABEL ANTIMICROBIAL DECLARATION

Authors declare no off-label use of antimicrobials.

INSTITUTIONAL ANIMAL CARE AND USE COMMITTEE (IACUC) OR OTHER APPROVAL DECLARATION

Authors declare no IACUC or other approval was needed.

HUMAN ETHICS APPROVAL DECLARATION

Authors declare human ethics approval was not needed for this study.

ORCID

Joonbum Seo  <https://orcid.org/0000-0002-8215-1237>

Jose Novo Matos  <https://orcid.org/0000-0001-8128-8111>

Virginia Luis Fuentes  <https://orcid.org/0000-0001-8076-3806>

REFERENCES

- Connolly DJ, Magalhaes RJ, Syme HM, et al. Circulating natriuretic peptides in cats with heart disease. *J Vet Intern Med.* 2008;22:96-105.
- Hsu A, Kittleson MD, Paling A. Investigation into the use of plasma NT-proBNP concentration to screen for feline hypertrophic cardiomyopathy. *J Vet Cardiol.* 2009;11(Suppl 1):S63-S70.
- Fox PR, Rush JE, Reynolds CA, et al. Multicenter evaluation of plasma N-terminal pro-brain natriuretic peptide (NT-pro BNP) as a biochemical screening test for asymptomatic (occult) cardiomyopathy in cats. *J Vet Intern Med.* 2011;25:1010-1016.
- Biondo AW, Ehrhart EJ, Sisson DD, Bulmer BJ, de Moraes HSA, Solter PF. Immunohistochemistry of atrial and brain natriuretic peptides in control cats and cats with hypertrophic cardiomyopathy. *Vet Pathol.* 2003;40:501-506.
- Wess G, Daisenberger P, Mahling M, Hirschberger J, Hartmann K. Utility of measuring plasma N-terminal-pro-brain natriuretic peptide in detecting hypertrophic cardiomyopathy and differentiating grades of severity in cats. *Vet Clin Pathol.* 2011;40:237-244.

6. Connolly DJ, Soares Magalhaes RJ, Fuentes VL, et al. Assessment of the diagnostic accuracy of circulating natriuretic peptide concentrations to distinguish between cats with cardiac and non-cardiac causes of respiratory distress. *J Vet Cardiol.* 2009;11(Suppl 1):S41-S50.
7. Hezzell MJ, Rush JE, Humm K, et al. Differentiation of cardiac from noncardiac pleural effusions in cats using second-generation quantitative and point-of-care NT-proBNP measurements. *J Vet Intern Med.* 2016;30:536-542.
8. Humm K, Hezzell M, Sargent J, Connolly DJ, Boswood A. Differentiating between feline pleural effusions of cardiac and non-cardiac origin using pleural fluid NT-proBNP concentrations. *J Small Anim Pract.* 2013;54:656-661.
9. Singletary GE, Rush JE, Fox PR, Stepien RL, Oyama MA. Effect of NT-pro-BNP assay on accuracy and confidence of general practitioners in diagnosing heart failure or respiratory disease in cats with respiratory signs. *J Vet Intern Med.* 2012;26:542-546.
10. Hori Y, Iguchi M, Heishima Y, et al. Diagnostic utility of cardiac troponin I in cats with hypertrophic cardiomyopathy. *J Vet Intern Med.* 2018;32:922-929.
11. Wells SM, Shofer FS, Walters PC, Stamoulis ME, Cole SG, Sleeper MM. Evaluation of blood cardiac troponin I concentrations obtained with a cage-side analyzer to differentiate cats with cardiac and noncardiac causes of dyspnea. *J Am Vet Med Assoc.* 2014;244:425-430.
12. Borgeat K, Sherwood K, Payne JR, Luis Fuentes V, Connolly DJ. Plasma cardiac troponin I concentration and cardiac death in cats with hypertrophic cardiomyopathy. *J Vet Intern Med.* 2014;28:1731-1737.
13. Langhorn R, Tarnow I, Willesen JL, Kjelgaard-Hansen M, Skovgaard IM, Koch J. Cardiac troponin I and T as prognostic markers in cats with hypertrophic cardiomyopathy. *J Vet Intern Med.* 2014;28:1485-1491.
14. Peterson EN, Moise NS, Brown CA, Erb HN, Slater MR. Heterogeneity of hypertrophy in feline hypertrophic heart disease. *J Vet Intern Med.* 1993;7:183-189.
15. Schober K, Todd A. Echocardiographic assessment of left ventricular geometry and the mitral valve apparatus in cats with hypertrophic cardiomyopathy. *J Vet Cardiol.* 2010;12:1-16.
16. Fox PR, Liu SK, Maron BJ. Echocardiographic assessment of spontaneously occurring feline hypertrophic cardiomyopathy. An animal model of human disease. *Circulation.* 1995;92:2645-2651.
17. MacLea HB, Boon JA, Bright JM. Doppler echocardiographic evaluation of midventricular obstruction in cats with hypertrophic cardiomyopathy. *J Vet Intern Med.* 2013;27:1416-1420.
18. Payne JR, Brodbelt DC, Luis FV. Cardiomyopathy prevalence in 780 apparently healthy cats in rehoming centres (the CatScan study). *J Vet Cardiol.* 2015;17(Suppl 1):S244-S257.
19. Fox PR, Keene BW, Lamb K, et al. International collaborative study to assess cardiovascular risk and evaluate long-term health in cats with preclinical hypertrophic cardiomyopathy and apparently healthy cats: the REVEAL study. *J Vet Intern Med.* 2018;32:930-943.
20. Cavalcante JL, Barboza JS, Lever HM. Diversity of mitral valve abnormalities in obstructive hypertrophic cardiomyopathy. *Prog Cardiovasc Dis.* 2012;54:517-522.
21. Ramos LW, Murad N, Goto E, et al. Ischemia/reperfusion is an independent trigger for increasing myocardial content of mRNA B-type natriuretic peptide. *Heart Vessels.* 2009;24:454-459.
22. Jung SW, Kittleson MD. The effect of atenolol on NT-proBNP and troponin in asymptomatic cats with severe left ventricular hypertrophy because of hypertrophic cardiomyopathy: a pilot study. *J Vet Intern Med.* 2011;25:1044-1049.
23. Goetze JP, Gore A, Moller CH, et al. Acute myocardial hypoxia increases BNP gene expression. *FASEB J.* 2004;18:1928-1930.
24. Iwanaga Y, Nishi I, Furuichi S, et al. B-type natriuretic peptide strongly reflects diastolic wall stress in patients with chronic heart failure: comparison between systolic and diastolic heart failure. *J Am Coll Cardiol.* 2006;47:742-748.
25. Lukowicz TV, Fischer M, Hense HW, et al. BNP as a marker of diastolic dysfunction in the general population: importance of left ventricular hypertrophy. *Eur J Heart Fail.* 2005;7:525-531.
26. Maron MS, Olivotto I, Betocchi S, et al. Effect of left ventricular outflow tract obstruction on clinical outcome in hypertrophic cardiomyopathy. *N Engl J Med.* 2003;348:295-303.
27. Ferasin L, Ferasin H. Plasma NT-pro-BNP and serum troponin-I concentrations in cats with systolic anterior motion of the mitral valve (SAM) not accompanied by left ventricular hypertrophy. *J Vet Intern Med.* 2012;26:1513.
28. Maron MS, Olivotto I, Maron BJ, et al. The case for myocardial ischemia in hypertrophic cardiomyopathy. *J Am Coll Cardiol.* 2009;54:866-875.
29. Connolly DJ, Cannata J, Boswood A, Archer J, Groves EA, Neiger R. Cardiac troponin I in cats with hypertrophic cardiomyopathy. *J Feline Med Surg.* 2003;5:209-216.
30. Lalor SM, Connolly DJ, Elliott J, Syme HM. Plasma concentrations of natriuretic peptides in normal cats and normotensive and hypertensive cats with chronic kidney disease. *J Vet Cardiol.* 2009;11(Suppl 1):S71-S79.
31. Luis Fuentes V, Abbott J, Chetboul V, et al. ACVIM consensus statement guidelines for the classification, diagnosis, and management of cardiomyopathies in cats. *J Vet Intern Med.* 2020. <https://doi.org/10.1111/jvim.15745>.
32. Hansson K, Haggstrom J, Kvarn C, Lord P. Left atrial to aortic root indices using two-dimensional and M-mode echocardiography in cavalier King Charles spaniels with and without left atrial enlargement. *Vet Radiol Ultrasound.* 2002;43:568-575.
33. Schober KE, Wetli E, Drost WT. Radiographic and echocardiographic assessment of left atrial size in 100 cats with acute left-sided congestive heart failure. *Vet Radiol Ultrasound.* 2014;55:359-367.
34. Marz I, Wilkie LJ, Harrington N, et al. Familial cardiomyopathy in Norwegian Forest cats. *J Feline Med Surg.* 2015;17:681-691.
35. Campbell FE, Kittleson MD. The effect of hydration status on the echocardiographic measurements of normal cats. *J Vet Intern Med.* 2007;21:1008-1015.
36. Payne JR, Borgeat K, Connolly DJ, et al. Prognostic indicators in cats with hypertrophic cardiomyopathy. *J Vet Intern Med.* 2013;27:1427-1436.
37. Payne JR, Borgeat K, Brodbelt DC, et al. Risk factors associated with sudden death vs. congestive heart failure or arterial thromboembolism in cats with hypertrophic cardiomyopathy. *J Vet Cardiol.* 2015;17(Suppl 1):S318-S328.
38. Gori F, Basso C, Thiene G. Myocardial infarction in a patient with hypertrophic cardiomyopathy. *N Engl J Med.* 2000;342:593-594.
39. Misawa K, Nitta Y, Matsubara T, et al. Difference in coronary blood flow dynamics between patients with hypertension and those with hypertrophic cardiomyopathy. *Hypertens Res.* 2002;25:711-716.
40. Rush JE, Freeman LM, Fenolosa NK, Brown DJ. Population and survival characteristics of cats with hypertrophic cardiomyopathy: 260 cases (1990-1999). *J Am Vet Med Assoc.* 2002;220:202-207.
41. Payne J, Luis Fuentes V, Boswood A, Connolly D, Koffas H, Brodbelt D. Population characteristics and survival in 127 referred cats with hypertrophic cardiomyopathy (1997 to 2005). *J Small Anim Pract.* 2010;51:540-547.
42. Abbott JA. Feline hypertrophic cardiomyopathy: an update. *Vet Clin North Am Small Anim Pract.* 2010;40:685-700.
43. Lamont LA, Bulmer BJ, Sisson DD, Grimm KA, Tranquilli WJ. Doppler echocardiographic effects of medetomidine on dynamic left ventricular outflow tract obstruction in cats. *J Am Vet Med Assoc.* 2002;221:1276-1281.
44. Ayoub C, Geske JB, Larsen CM, Scott CG, Klarich KW, Pellikka PA. Comparison of Valsalva maneuver, amyl nitrite, and exercise echocardiography to demonstrate latent left ventricular outflow obstruction in hypertrophic cardiomyopathy. *Am J Cardiol.* 2017;120:2265-2271.

45. Chetboul V, Lefebvre HP, Pinhas C, Clerc B, Boussof M, Pouchelon JL. Spontaneous feline hypertension: clinical and echocardiographic abnormalities and survival rate. *J Vet Intern Med.* 2003;17:89-95.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of this article.

How to cite this article: Seo J, Payne JR, Novo Matos J, Fong WW, Connolly DJ, Luis Fuentes V. Biomarker changes with systolic anterior motion of the mitral valve in cats with hypertrophic cardiomyopathy. *J Vet Intern Med.* 2020;1-10. <https://doi.org/10.1111/jvim.15807>