

This is the author's accepted manuscript. The version of record is available online via the *Journal of Antibiotics*: <https://doi.org/10.1038/s41429-019-0149-0>.

The full details of the published version of the article are as follows:

TITLE: Doxycycline inhibits pre-rRNA processing and mature rRNA formation in *E. coli*

AUTHORS: CU Chukwudi, Liam Good

JOURNAL TITLE: Journal of Antibiotics

PUBLICATION DATE: 8 February 2019 (online)

PUBLISHER: Springer Nature

DOI: 10.1038/s41429-019-0149-0

# 8 **Doxycycline inhibits Pre-rRNA Processing and Mature rRNA**

## 9 **Formation in *E. coli***

10 Chinwe .U. Chukwudi <sup>1,2\*</sup>, Liam Good <sup>1</sup>

11 <sup>1</sup> Department of Pathology and Infectious Diseases, Royal Veterinary College, University of London,  
12 UK

13 <sup>2</sup> Department of Veterinary Pathology and Microbiology, University of Nigeria, Nsukka, Nigeria

14 \* To whom correspondence should be addressed. Email: [chinwe.chukwudi@unn.edu.ng](mailto:chinwe.chukwudi@unn.edu.ng)

15

### 16 **Abstract**

17 In bacteria, RNase III cleaves the initial long primary ribosomal RNA transcripts/precursors  
18 (pre-rRNAs), thereby releasing the pre-16S and pre-23S rRNAs for maturation. This cleavage  
19 is specified by the double-stranded secondary structures flanking the mature rRNAs, and not  
20 necessarily by the nucleotide sequences. Inhibition of this cleavage would lead to a build-up  
21 of pre-rRNA molecules. Doxycycline has earlier been shown to bind synthetic double-  
22 stranded RNAs and inhibit their cleavage by RNase III. Since bacterial rRNA processing is  
23 primarily dependent on RNase III cleavage (which is inhibited by doxycycline), doxycycline  
24 could therefore inhibit the normal processing of bacterial rRNA. In this study, the effect of  
25 doxycycline on bacterial rRNA processing was investigated by analyzing the amounts of  
26 various rRNAs in growing *E. coli* cells treated with doxycycline. The results showed a  
27 doxycycline dose-dependent decrease in mature 16S and 23S rRNAs, concurrent with an  
28 accumulation of the initial rRNA transcripts and long precursors. Morphologically, treated  
29 cells were elongated at low drug concentrations, while nucleoid degeneration indicative of  
30 cell death occurred at higher drug concentrations. These observations suggest that

31 doxycycline inhibits the cleavage and processing of bacterial rRNA transcripts/precursors,  
32 leading to impaired formation of mature rRNAs, and the consequent inhibition of protein  
33 synthesis for which the tetracycline group of antibiotics are renowned. Since rRNA structure  
34 and processing pathway is conserved among bacterial species, this mechanism may account  
35 for the broad spectrum of antibiotic activity and selective microbial protein synthesis  
36 inhibition of doxycycline and the tetracyclines.

37 Keywords: doxycycline/pre-rRNA/ribosomal RNA processing/RNase III  
38 cleavage/tetracycline antibiotics.

### 39 **Introduction**

40 Doxycycline is a member of the broad spectrum group of antibiotics known as the  
41 tetracyclines. The tetracyclines are known to inhibit bacterial protein synthesis by binding to  
42 the 16S ribosomal RNA (rRNA) and inhibiting the binding of aminoacyl-tRNA to the  
43 mRNA-ribosome complex [1-4]. However, their activity against other microbes which do not  
44 possess the 16S rRNA such as viruses, protozoa, and helminths has raised further questions  
45 as to the exact mechanism of action. In addition, despite conservation of ribosome structure  
46 and function between bacteria and host cells, the tetracyclines are sufficiently selective that  
47 the protein synthesis machinery of the host organism remains relatively unaffected. Despite  
48 their long history of usage as therapeutic agents, the mechanism(s) by which the tetracyclines  
49 achieve their wide range of effects and selectively inhibit microbial protein synthesis is not  
50 yet fully understood.

51 Even though binding interactions with both the 16S and 23S rRNAs had earlier been  
52 indicated for the tetracyclines [5], an *in vitro* study to correlate ribosomal subunit activity  
53 with drug binding suggested that inhibition of tRNA binding to the A-site is solely due to  
54 tetracycline crosslinked to the strong binding site on the 30S ribosomal subunit [6]. Hence,

55 subsequent investigations on the mechanism of action of the tetracyclines and their  
56 interaction with ribosomal RNA concentrated on the 16S rRNA of the 30S ribosomal subunit  
57 [7]. Nevertheless, a recent study has shown that the tetracyclines (doxycycline and  
58 minocycline) bind to various synthetic double-stranded RNAs of random base sequence and  
59 inhibit their cleavage by RNase III *in vitro* [8]. This could imply that the double stranded  
60 secondary structures that frequently occur in cellular RNAs may be more crucial for the  
61 binding of the tetracyclines to RNA than the specific base pairs; and that the mechanism of  
62 action of the tetracyclines may be linked to the effect of the drugs on the processing of such  
63 cellular RNAs. If this is correlated *in vivo*, it could offer insights into the mechanism that  
64 underlie the activity of the tetracyclines against a wide range of pathogens, as well as in other  
65 non-infectious therapeutic indications of the drugs.

66 Ribosomal RNAs constitute about 95% of total cellular RNA in *E. coli* [9]. They form the  
67 active sites of the ribosomes for decoding the message of the mRNA, as well as perform  
68 enzymatic functions in the translation process [10]. The rRNAs of prokaryotes are co-  
69 transcribed from an operon (Fig 1), and *E. coli* has 7 copies of rRNA operons in the  
70 chromosome [11]. RNase III then cleaves the nascent rRNA transcripts at the double-  
71 stranded stem regions that flank the mature 16S and 23S ribosomal RNA sequences to release  
72 the pre-16S and pre-23S rRNAs for further maturation [11-13]. In wild type cells, RNase III  
73 cleavage is very rapid, and occurs concurrently with transcription. Hence, only a very small  
74 amount of the long primary transcript (1-2% of total rRNA) is reported to be detectable in *E.*  
75 *coli* [14]. However, in RNase III-deficient strains, the 30S pre-rRNA accumulates [15]. In  
76 this study, the effects of doxycycline on the processing of bacterial ribosomal RNAs were  
77 investigated and correlated with their antibiotic activity in growing *E. coli* cells, with a view  
78 to elucidate the molecular mechanism of their antimicrobial activity.

79

## 80 **Materials and methods**

81 Probes and primers used in this study are listed in Table 1, and obtained from Sigma®  
82 Aldrich. Doxycycline was also purchased from Sigma® Aldrich. Nylon membranes were  
83 purchased from Roche (Roche # 11209299001). Hybridization probe labelling and detection  
84 was done using AlkPhos Direct™ labelling and detection System with ECF™ from GE  
85 healthcare life sciences (RPN3692).

86 All procedures (sample collection, RNA extraction, gel electrophoresis, transfer to nylon  
87 membranes and northern blot hybridization) were performed at least in replicates of three  
88 independent experiments.

89

## 90 **Total RNA extraction**

91 An overnight culture of *E. coli* strain K-12 grown in LB broth was diluted 20 fold with fresh  
92 medium and incubated at 37°C in a Stuart orbital incubator S1500 with shaking for 1hr to  
93 ensure growth is activated. The culture was divided into aliquots to which were added 0-200  
94 µM doxycycline (0-96 µg/ml) and incubated at 37°C with shaking (180 rpm). Optical density  
95 (OD) of cultures was measured at 550nm using Biotek Powerwave XS universal spectrometer.  
96 2 ml of culture samples were taken from each treatment group at the indicated time points (0-  
97 240 min). Nucleic acid decay was stopped in collected samples by immediate transfer of the  
98 samples to a cold microcentrifuge tube (on ice) containing 200 µl ethanol and 20 µl phenol  
99 [16]. The bacterial cells were harvested by centrifuging at 8000 rpm for 1 min, and lysed by  
100 re-suspending the pellet in 400 µl of Sigma B cell lysis reagent (Sigma® Aldrich). Total  
101 RNA was harvested by the phenol-chloroform extraction method with ethanol precipitation

102 [17]. 400 µl phenol (Sigma) was added to the samples and the tube vortexed vigorously for at  
103 least 2 min total. 400 µl RNase-free water and 400 µl chloroform (Sigma) was then added  
104 and vortexed vigorously for another 2 min. The tube was then centrifuged at 12000 rpm for  
105 15 min and 600 µl of the aqueous phase transferred to a fresh 2 mL tube. 60 µl of 3 M  
106 sodium acetate and 1.4 ml ethanol was added, mixed and held on ice for 10 min, and  
107 centrifuged at 10000 rpm in a microcentrifuge for 15 min. The supernatant was removed, and  
108 the pellet rinsed with 70% ethanol and dried. The resultant nucleic acid pellets were  
109 resuspended in 1x TE buffer (10 mM Tris, pH 8.0, 1 mM EDTA) and stored at -20°C until  
110 further use. The nucleic acid concentration of the extracts was quantified using NanoDrop  
111 ND-1000 spectrophotometer, and scored as an average of at least three readings.

#### 112 **Northern blot analysis of mature rRNA from *E. coli* total RNA extract**

113 Ribosomal RNA was separated from the total RNA extracts by agarose gel electrophoresis. 5  
114 µl total RNA extracts from each sample was loaded in 1% agarose gels containing 1X MOPS  
115 buffer. For denaturing gel electrophoresis, the RNA samples were incubated with 3X volume  
116 of formaldehyde loading dye (Ambion) at 65°C for 10 min before loading. Electrophoresis  
117 was carried out at 90 V for 40 min in 1X MOPS buffer and gels were stained with 1X EtBr or  
118 SYBR® Gold (Invitrogen™). Images were taken with SynGene G:Box camera using  
119 GeneSnap software. The rRNA was subsequently transferred unto positively charged nylon  
120 membranes by gravity and capillary action. The gel was soaked in 20X saline-sodium citrate  
121 (SSC) buffer for 30 min, placed right side up on the nylon membrane without trapping air  
122 bubbles in-between and covered on both sides with Whatman filter paper that had been cut to  
123 size and soaked in 20X SSC buffer. The gel was wrapped around the edges with parafilm to  
124 prevent drying and ensure that transfer proceeds only through the gel. About 2 cm of paper  
125 towel was also cut to size, soaked in 10X SSC buffer and placed on top of the filter paper. A

126 small flat weight was then placed on top of the stack which was allowed to stand overnight.  
127 After overnight transfer, the membrane was rinsed with 2X SSC buffer and air-dried. RNA  
128 was cross-linked to the membrane by UV illumination for 1-1.5 min using SynGene G:Box  
129 transilluminator. Membranes were then stained with 2% methylene blue and de-stained in 2%  
130 SSC buffer to check for mature rRNA bands (visible to the naked eye). Images were taken  
131 using SynGene G:Box image camera. Membranes were subsequently stored dry at 4°C until  
132 used for hybridization.

### 133 **Probe design**

134 CCPR1 was designed to target a sequence 21 base pairs from the 3' end of *E. coli* K-12 16S  
135 rRNA, and purchased from Sigma. CCPR2 was designed to target the sequence between 10  
136 bp downstream of K-12 16S rRNA and 10 bp upstream of 23S rRNA, including the  
137 intergenic sequences (Fig 1B). BLAST search (Basic Local Alignment Search Tool, NCBI)  
138 indicated no similarity to any other *E. coli* gene segment apart from the ribosomal RNA.  
139 CCPR2 was synthesized by PCR amplification using the primers shown on Table 1. The  
140 amplicon size was verified by matching the band of the PCR product on agarose gel  
141 electrophoresis with the band of DNA ladder of the expected size.

### 142 **Nucleic acid hybridization for detection of pre-rRNAs**

143 The probe was labelled with alkaline phosphatase using AlkPhos Direct™ labelling and  
144 detection System with ECF™ (GE healthcare life sciences) following the manufacturer's  
145 instructions. For this experiment, a ten-fold dilution of overnight culture of *E. coli* K12 cells  
146 was grown to exponential phase for about 1.5 hrs in LB broth (to ensure a high cell  
147 harvest/rRNA yield) before adding doxycycline at the specified concentrations. The cultures  
148 were then incubated for only 20 min to minimize the differential inhibitory effects of the  
149 antibiotic concentrations on culture growth (OD) and concentration of the RNA extracted.

150 Hence, samples were collected at 20 min incubation time. The total RNA was extracted and  
151 separated by gel electrophoresis, then transferred onto nylon membranes. Membrane blots of  
152 the total RNA extracts were equilibrated in hybridization buffer for 15 min before overnight  
153 hybridization with the labelled probe in a hybridization oven at 42°C. Membranes were  
154 washed at 42°C with the primary wash buffer, and at room temperature with the secondary  
155 wash buffer. Detection reagent was applied to the membranes, which were then wrapped with  
156 cling film and incubated overnight in the dark to enhance development of the fluorescence  
157 signal. Blots were scanned and images were taken using SynGene G:Box camera.

### 158 **Fluorescence microscopy**

159 *E. coli* strain K-12 was grown in LB broth containing 0-200 µM doxycycline at 37°C for 20  
160 min. Cells were harvested by centrifuging and washed twice with 2x volume of 1x PBS. They  
161 were stained using DAPI (to examine the nucleoid morphology) by adding 1x volume of 1  
162 µM DAPI (Sigma) and incubating in the dark for about 5 min. Samples were then mounted  
163 on glass slides with cover slips. Cell morphology was examined with Leica DM4000B  
164 fluorescence microscope using DAPI filter. Images were captured with DC500 camera using  
165 Leica IM500 software programme.

### 166 **Data Analysis**

167 All RNA band intensities were quantified using the image analysis software, GeneTools from  
168 SynGene (Cambridge, UK). Statistical analysis was done using GraphPad Prisms 7.02.  
169 Means were compared using repeated measure ANOVA or paired t-test (as appropriate),  
170 while dose-response effects were analysed using non-linear regression fitted for direct (non-  
171 normalized) response. Statistical significance was considered at 95% confidence interval ( $P$   
172  $\leq 0.05$  significance level).

173 **Results**

174

175 **Doxycycline reduces the amounts of mature ribosomal RNA *in vivo***

176 In view of earlier reports that doxycycline inhibits RNase III degradation/cleavage of double-  
177 stranded RNA *in vitro* [8], the effect of doxycycline on RNase III-dependent dsRNA  
178 cleavage/processing pathways *in vivo* was investigated. The most important and generalized  
179 RNase III-dependent processing pathway in bacteria cells with respect to protein synthesis is  
180 the processing of ribosomal RNA. RNase III cleavage is the rate limiting step for the  
181 formation of mature rRNAs which is necessary for protein synthesis in growing bacteria cells.  
182 To assess the effect of doxycycline on this processing pathway *in vivo*, total RNA was  
183 harvested from doxycycline-treated and untreated *E. coli* K-12 cells and analysed by native  
184 agarose gel electrophoresis. The intensities of the mature ribosomal RNA bands in the cells  
185 growing in the presence of 100 $\mu$ M doxycycline over a given time was compared with those  
186 of cells growing in the absence of the drug. The results show a significant ( $P= 0.0046$  for 23S  
187 and 0.0091 for 16S rRNA) and progressive reduction ( $r = -0.7365$  for 23S and  $-0.8126$  for  
188 16S rRNA) in the band intensities of mature ribosomal RNAs over time in the cells growing  
189 with doxycycline, in contrast to those growing without the drug, which showed a progressive  
190 increase in rRNA band intensities that peaked at about 210 min (Fig 2). In the sample  
191 containing doxycycline, there was a sharp drop in the band intensities of the 16S and 23S  
192 ribosomal RNAs at 20 min, which subsequently increased slightly between 40-90 min before  
193 ultimately fading away. When gels of samples containing other concentrations of doxycycline  
194 (2-200 $\mu$ M) over time were analysed, it was observed that this sharp drop in the band  
195 intensities of the 16S and 23S rRNAs at 20 min only occurred at higher doxycycline  
196 concentrations (100-200 $\mu$ M); but at lower doxycycline concentrations (0-50 $\mu$ M), the

197 decrease is gradual and steady. These results suggest that doxycycline inhibits the formation  
198 of mature rRNAs, although the inhibition of bacterial growth in the presence of the drug may  
199 contribute to this effect.

200

201 To determine whether the observed depletion of 16S and 23S rRNAs in doxycycline-treated  
202 samples was caused by inhibited rRNA formation due to processing rather than inhibited  
203 synthesis due to growth inhibition or death of bacteria cells, the RNA samples were also  
204 analysed by northern blot hybridization to detect the pre-16S rRNA (using CCPR1 probe). If  
205 the reduction in mature rRNA band intensities seen in Fig 2A was simply due to reduced cell  
206 numbers by the growth inhibitory activity of the drug, one would expect a commensurate and  
207 concurrent reduction in the amounts of pre-rRNA detected by northern blotting. However,  
208 northern blot hybridization assay of the samples showed the presence of long pre-rRNAs in  
209 both treated and untreated samples, with no significant difference ( $P= 0.7157$ ) between the  
210 treated and untreated groups (Fig 2B). There was no reduction in pre-rRNA band intensity in  
211 doxycycline-treated cells. Instead, there was smearing of the pre-rRNA bands in doxycycline-  
212 treated cells from earlier incubation time points (20-60 min), indicating the presence of  
213 variable sizes of pre-16S rRNA in all doxycycline-treated samples. This smearing only  
214 occurred at longer incubation times (120-240 min) in untreated samples. These results  
215 suggest that rRNA was still being transcribed in doxycycline-treated cells at all time points  
216 studied, but the rRNA transcripts and pre-rRNA were not adequately processed to form the  
217 mature 16S and 23S rRNAs.

218 Interestingly, in the course of these experiments, some samples were analysed by denaturing  
219 gel electrophoresis that was run much longer (2-4hr) to allow a better separation of the RNA  
220 bands. In these experimental conditions, we observed a difference in the profile of the long

221 pre-rRNA bands between the samples that contain doxycycline and those without the drug.  
222 Whereas only one band was seen in the samples without the drug from about 60 min, the  
223 samples containing the drug showed an additional second band from about 210 min  
224 incubation time which represents different species/sizes of the long pre-rRNA. Since the pre-  
225 rRNA are cleavage products of the initial transcripts, the observation of different pre-rRNA  
226 sizes between doxycycline-treated and untreated cells is suggestive of impaired/abnormal  
227 cleavage or processing of the rRNA transcripts in the presence of doxycycline.

228

229 **Doxycycline induces a dose-dependent inhibition of mature ribosomal RNA in growing**  
230 **bacteria cells**

231 In order to further investigate the involvement of doxycycline in the observed reduction in the  
232 amounts of mature ribosomal RNA in *E. coli* K-12 cells, total RNA from cells grown in the  
233 presence of various concentrations of doxycycline was assessed. Samples collected at both 20  
234 and 120 min of incubation time showed a dose-dependent decrease in the amounts of mature  
235 16S and 23S rRNA with increasing amounts of doxycycline (Fig 3). Statistical analysis  
236 showed that the concentration of doxycycline that gave a response half-way between baseline  
237 and maximal ( $IC_{50}$ ) at 20 min incubation time was  $8.327\mu\text{M}$  ( $\pm$ -SE 2.465,  $R^2= 0.9554$ ), and  
238 within the range of 4.295-17.95  $\mu\text{M}$  at 95% confidence interval (CI). This increased at 120  
239 min incubation time to 76.51  $\mu\text{M}$  ( $\pm$ -SE 49.6,  $R^2= 0.8947$ ), and within 24.05-392.6  $\mu\text{M}$  at  
240 95% CI.  $IC_{95}$  was observed at 100  $\mu\text{M}$  doxycycline concentration, which had the lowest  
241 rRNA band intensity observed at both 20 and 120 min incubation times. The lower  
242 concentrations of doxycycline (2-20  $\mu\text{M}$ ) showed a slight increase in the intensity of mature  
243 rRNA bands at the longer incubation time (Fig 3A, 120 min). Further analysis of samples at  
244 increasing incubation time of different doses showed that this increase in mature rRNA band

245 intensity at lower concentrations of doxycycline was sustained over the duration of the  
246 experiments (240 min), and was highest at the lowest concentration of doxycycline used (2  
247  $\mu\text{M}$ ). This suggests that sub-inhibitory concentrations of doxycycline may induce rRNA  
248 formation/synthesis with time. In addition, there was also a general increase in the total RNA  
249 concentration of doxycycline treated samples at 120 min compared to the untreated samples  
250 (which was highest at 5  $\mu\text{M}$  drug concentration; Fig 3F), despite decreased culture  
251 growth/OD at that incubation time (Fig 3E). However, in spite of the fact that the higher drug  
252 concentrations (50-200  $\mu\text{M}$ ) had higher total RNA concentration than the untreated cells at  
253 120 min incubation time (Fig 3E), they still showed decreased 16S and 23S rRNA band  
254 intensities in the gel (Fig 3A). This suggests that much of the RNAs at these drug  
255 concentrations are not mature rRNA. The growth curves also showed that all drug  
256 concentrations produced similar growth inhibitory effects at 20 min incubation time, but had  
257 variable effects at longer incubation times. At 120 min incubation time, the maximum growth  
258 inhibition was achieved with 10-20  $\mu\text{M}$  (Fig 3E). These results therefore indicate that the  
259 effect of doxycycline on the formation of rRNAs in growing bacteria cells is affected by both  
260 drug dosage and incubation/treatment time, and that very low doses of doxycycline may  
261 induce rRNA transcription/formation over time.

262 When the samples were analysed by northern blot hybridization to detect pre-16S rRNA  
263 using CCPR1 probe (Fig 3B), the intensity of the long rRNA precursors (initial transcript and  
264 pre-rRNA) increased with increasing drug concentration at 20 min incubation time, with  
265 smearing at the lower drug concentrations (2-20  $\mu\text{M}$ ). This indicates that more pre-16S rRNA  
266 is being retained in the long precursors with increasing drug concentration. At 120 min, all  
267 the rRNA bands (except the rRNA transcript band) were smeared. Besides smearing, the pre-  
268 rRNA bands became faint whereas the mature rRNA bands became more prominent at 120

269 min, especially at lower drug concentrations (2-20  $\mu$ M) when compared to the 20 min  
270 samples. This indicates that the pre-rRNA was being cleaved (albeit inadequately) into  
271 smaller particles about the size of the mature 16S and 23S rRNAs with time, especially at low  
272 drug concentrations. This is consistent with the observed increase in band intensities of the  
273 16S and 23S rRNAs at these low drug concentrations and longer incubation time in the gel  
274 image (Fig 3A). Altogether, these results suggest that the inhibition of mature rRNA  
275 formation by doxycycline could be due to inadequate cleavage/processing of the long rRNA  
276 transcripts and pre-rRNAs.

277

### 278 **Doxycycline induces accumulation of pre-rRNAs in growing bacteria cells**

279 The observation that doxycycline inhibits RNase III cleavage of total RNA extracts *in vitro*  
280 and the formation of mature rRNAs *in vivo* could imply that the drug inhibits the cleavage  
281 and processing of the primary rRNA transcripts and pre-rRNA. If this is true, then  
282 doxycycline would induce accumulation of the unprocessed pre-rRNAs. This would  
283 substantiate the decrease in 16S and 23S rRNA bands as resulting from the effect of the drug  
284 on ribosomal RNA processing, rather than just a reflection of the rate of culture growth.  
285 Hence, the effect of doxycycline on the cleavage/processing of the primary ribosomal RNA  
286 transcript was further investigated *in vivo* by northern blot hybridization assay to assess the  
287 amounts of long primary rRNA transcripts and pre-rRNAs in growing bacteria cells treated  
288 with various concentrations of doxycycline. To minimize the growth inhibitory effect of the  
289 antibiotic and ensure good RNA yield in this experiment, bacterial cultures were initially  
290 grown to exponential phase before treatment, and thereafter, samples were harvested at 20  
291 min incubation time to minimize differential growth in the antibiotic media (culture OD and  
292 total RNA extract concentrations were also assessed for confirmation). A probe (CCPR2) that

293 is complementary to the spacer region between the mature 16S and 23S rRNAs (including  
294 about 10 nucleotides upstream and downstream as shown in Fig 1) was used to detect  
295 uncleaved rRNA transcripts and long pre-rRNAs in the total RNA extracts. If rRNA  
296 processing occurs normally, this region is cleaved off by RNase III, and further processing  
297 yields mature 16S and 23S rRNAs. If RNase III cleavage is inhibited, this region is retained  
298 and would accumulate in the initial transcripts and long pre-rRNA.

299 Northern blot hybridization assay of the total RNA extracts from cells grown in the presence  
300 of various concentrations of doxycycline for 20 min (using the probe CCPR2) showed a dose-  
301 dependent accumulation of the initial transcripts and long pre-rRNA species, concomitant  
302 with a dose-dependent decrease in the 16S and 23S rRNA species (Fig 4). There were two  
303 distinct bands representing two uncleaved long rRNA precursors detected by the probe: a  
304 long initial rRNA transcript (positioned just below the wells), and a long pre-rRNA  
305 (estimated to be about 9KB in size). Linear regression analysis of culture OD showed no  
306 significant change in the culture OD ( $P=0.7745$ ). For total RNA concentration, 50% of the  
307 observed reduction was induced at about 192 $\mu$ M doxycycline concentration ( $IC_{50}$  at  
308  $R^2=0.8534$ ). Despite the culture OD being stable at all drug concentrations, and decreasing  
309 total RNA concentrations from 50-200 $\mu$ M drug concentrations (Fig 4C), both the initial  
310 rRNA transcript and the pre-rRNA band intensities increased with increasing doxycycline  
311 concentration (Fig 4D). It is interesting to note that the highest pre-rRNA band intensity  
312 occurred at doxycycline concentrations at which total RNA concentrations decreased. These  
313 observations strongly indicate that the effects of doxycycline on rRNA band intensities may  
314 be due to inhibition of rRNA transcript processing, and not essentially a reflection of culture  
315 growth inhibition by the antibiotic.

316 When the RNA band intensities were quantified by densitometry, statistical analysis of the  
317 results showed that 27.21 (+/- 19)  $\mu\text{M}$  doxycycline concentration (or 8.832-104.2  $\mu\text{M}$  at 95%  
318 CI,  $R^2=0.7949$ ) induced 50% of the observed accumulation of pre-rRNAs ( $\text{EC}_{50}$ ). On the  
319 other hand, 4.149 (+/- 1.535)  $\mu\text{M}$  doxycycline concentration induced 50% of the observed  
320 reduction in 16S and 23S rRNAs (2.028-8.553  $\mu\text{M}$  at 95% CI,  $R^2=0.9596$ ). It should be noted  
321 that the 16S and 23S rRNAs detected in this experiment are not yet fully mature (as they still  
322 contain some base sequences that are excised at maturation, which is detectable by the probe),  
323 and may differ slightly from the fully matured ones described in the previous sections. The  
324 concurrent increase in precursor rRNA species and decrease in 16S and 23S rRNAs indicates  
325 that much of the rRNAs are increasingly present as long precursor rRNAs with increasing  
326 doxycycline concentration. This reaches a peak at the higher drug concentrations (50-200 $\mu\text{M}$ )  
327 when only about 10% of the rRNA detected by the probe is in the 16S and 23S rRNA bands  
328 (Fig 4D inset). Taken together, these results indicate that doxycycline induces accumulation  
329 of uncleaved/long rRNA precursors, while inhibiting the formation of 16S and 23S rRNAs.  
330 This implies that doxycycline inhibits the cleavage of the rRNA transcripts and pre-rRNA  
331 into the smaller 16S and 23S fragments.

332

### 333 **Doxycycline induces bacterial cell elongation**

334 In order to correlate the molecular observations of the effects of doxycycline on ribosomal  
335 RNAs with the effect of the drug on the whole bacteria cell *in vivo*, the nucleoid morphology  
336 of cells treated with increasing concentrations of doxycycline for 20 min was examined. The  
337 results show that doxycycline induces elongation of bacteria cells at low doses ( $\leq 50\mu\text{M}$ ),  
338 which is indicative of cell division inhibition. At higher drug concentrations, nucleoid  
339 degeneration was observed, which is indicative of early stages of cell death (Fig 5). This is

340 consistent with the observed decrease in total RNA concentration at 50-200 $\mu$ M doxycycline  
341 concentration, suggesting that bacterial cell death occurs at high drug concentrations.

342

### 343 **Discussion**

344 The currently held 16S rRNA binding mechanism of action of the tetracyclines so far have  
345 not been sufficiently correlated with *in vivo* effects of the drug and their wide range of  
346 antimicrobial (not just antibacterial) activities [18]. The recently reported double-stranded  
347 RNA binding may therefore be a mechanism worth investigating to help elucidate the  
348 molecular basis of their wide range of activities [8]. If the tetracyclines bind to dsRNA and  
349 inhibit their cleavage/degradation by RNase III as previously reported [8], it could induce the  
350 accumulation of rRNA transcripts/precursors in growing bacteria cells.

351 The results presented here show a dose-dependent reduction of 16S and 23S rRNAs,  
352 concurrent with the accumulation of long rRNA precursors by doxycycline. Although any  
353 antibiotic that causes a reduction in bacterial growth would result in fewer cells growing in a  
354 culture medium, the observations in this study cannot be merely attributed to the growth  
355 inhibitory effects of an antibiotic. Several factors point towards a specific effect of  
356 doxycycline on mature rRNA formation rather than a reflection of the amount of cells in the  
357 culture. For instance, these effects were mostly observed at 20 min incubation time, when the  
358 effect of the drug on culture growth (OD) and total RNA concentration was minimal.

359 Particularly, the greatest increase in pre-rRNA band intensity (Fig 4) occurred at drug  
360 concentrations at which there was a decrease in total RNA concentration (50-200 $\mu$ M). Even  
361 the cell morphology changes, which are consistent with the molecular observations and  
362 earlier reports for tetracycline [19], were also observed at 20 min incubation time. Moreover,

363 the increase in mature rRNA band intensities of low drug concentrations at longer incubation  
364 periods (120mins) when the growth inhibitory effects of the drug should have been more  
365 pronounced (Fig 3 A, D) indicate an effect on rRNA processing rather than culture growth.  
366 This is in agreement with earlier reports that Chlortetracycline induces initial stimulation of  
367 RNA synthesis especially at low concentrations, and subsequent accumulation of RNA while  
368 inhibiting protein synthesis [20, 21]. These reports suggested that the accumulated RNA  
369 species differ from both 23S and 16S rRNAs in their sedimentation properties (attributed to  
370 “incomplete precursors”), but could synthesize ribosomes during recovery from the antibiotic  
371 effects. In this study, the concurrent decrease in mature rRNAs and increase in precursor  
372 rRNAs as detected by northern blot hybridization (Fig 4D) indicate effects on rRNA  
373 processing by doxycycline. Furthermore, the observations of smeared pre-rRNA bands at  
374 longer incubation periods which decrease in intensity as the mature rRNAs increase in  
375 intensity (Fig 3B) also indicate effects on rRNA cleavage/processing. In view of the ability of  
376 doxycycline to inhibit RNase III degradation/cleavage of dsRNA [8], these results indicate  
377 that doxycycline inhibits the cleavage of long rRNA transcripts/precursors by RNase III;  
378 leading to the accumulation of the pre-rRNAs [15]. This initial inhibition of cleavage of the  
379 long rRNA precursors by doxycycline is subsequently relieved with time (Fig 3), as has also  
380 been demonstrated *in vitro* with synthetic dsRNA [8]. A combination of this subsequent  
381 recovery from the inhibitory effects of doxycycline with time and possible alternate  
382 processing pathway which is less efficient than the RNase III cleavage pathway [15, 22],  
383 would lead to improved processing of the rRNA precursors at longer incubation periods. This  
384 could explain the observation of increased mature rRNA band intensities at longer incubation  
385 time with lower drug concentrations (Fig 3A). The dose-dependent increase in the long rRNA  
386 precursors (Fig 4B) seems to suggest that doxycycline also stimulates rRNA transcription.  
387 This may occur via a positive feedback mechanism, as the transcribed rRNA is not being

388 processed to yield functional mature rRNA. Such feedback mechanisms involved in  
389 transcriptional regulation have been described in bacteria [23-25], and have recently been  
390 associated with the regulation of rRNA transcription [26]. On the other hand, it is unlikely  
391 that the inhibition of mature rRNA formation was due to inhibition of transcription. If that  
392 was so, one would expect a decrease in the initial rRNA transcript amounts. However, the  
393 reverse was the case in this study (Fig 3B, 4B&D), indicating the possibility of a positive  
394 feedback mechanism instead. The general picture appears to be like this: As doxycycline is  
395 added, the mature rRNA decreases and the cells react to the shortage of mature rRNAs by  
396 increasing rRNA transcription. At higher doxycycline concentrations, more  
397 uncleaved/unprocessed pre-rRNA accumulate, and the cells activate/enhance alternative  
398 cleavage/processing pathways (such as by other nucleases) in an attempt to clear the  
399 accumulating pre-rRNAs.

400 It has been reported that although RNase III cleavage is necessary for the maturation of  
401 23S rRNA, it is not essential for its function [27]. On the other hand, maturation of 16S  
402 rRNA could proceed in the absence of RNase III cleavage, as has been demonstrated in  
403 RNase III-deficient strain, even though such strains are known to grow slowly [28]. This is  
404 believed to be due to an alternative processing pathway in the absence of RNase III by other  
405 nucleases acting independently of RNase III [29, 30]. However, unlike the immature 23S  
406 rRNA which is functional in protein synthesis, the immature 16S rRNA is not functional in  
407 protein synthesis [28]. In this study, doxycycline was found to inhibit the amounts of both the  
408 16S and 23S rRNAs. It is therefore possible that the non-functionality of the immature 16S  
409 rRNA, in contrast to the functionally active immature 23S rRNA, led to the previous belief  
410 that the tetracyclines exert their antibacterial action solely by binding to the 16S rRNA [6].

411 It is interesting to note that the inhibitory effects of doxycycline on rRNA processing were  
412 observed at the effective antibacterial concentrations of the drug. MIC of doxycycline for *E.*  
413 *coli* K-12 and the range of plasma concentrations following clinical therapeutic usage is  $\approx$ 4-  
414 8 $\mu$ g/ml ( $\approx$ 10-20 $\mu$ M). However, drug concentrations in organs may reach 10-25 times that of  
415 serum [31]. Also, time-kill studies have shown that doxycycline exhibits time-dependent  
416 antibacterial effect on *E. coli* at low concentrations (2-4 times the MIC), but optimal dose-  
417 dependent killing is achieved at higher drug concentrations of about 8-16 times the MIC [32].  
418 This complex interplay of dose and time was also observed in this study on the effect of  
419 doxycycline on mature rRNA formation (Fig 3), and could have clinical implications for the  
420 effective use of doxycycline and other tetracycline antibiotics. Also, mutations in the 16S  
421 rRNA sequence that have been shown to confer resistance to tetracycline often occur at the  
422 double-stranded stem regions, and disrupt base pairing and formation of the secondary  
423 structures necessary for RNase III recognition and cleavage [33].

424 The broad spectrum of antibacterial activity of the tetracyclines can be attributed to the  
425 highly conserved nature of rRNA processing via RNase III cleavage pathway among  
426 prokaryotes. In eukaryotes however, the processing of the ribosomal RNA involves a much  
427 more complex pathway that is not dependent on RNase III [34]. In addition, eukaryotic rRNA  
428 processing, occurs in a protected environment (nucleolus) where ionic conditions (especially  
429  $Mg^{2+}$ /divalent metal ion concentrations) are not ideal for doxycycline binding [8]. These  
430 differences in the processing pathway of prokaryotic and eukaryotic ribosomal RNAs could  
431 account for the selective inhibition of microbial protein synthesis, with minimal effects on  
432 eukaryotic protein synthesis [18]. The recovery from the inhibitory effects of the drug on the  
433 formation of mature ribosomal RNA with time supports the bacteriostatic mode of action of  
434 the tetracyclines.

435 Although the results presented here for doxycycline slightly digress from the 16S rRNA  
436 binding mechanism of action currently held for the tetracycline antibiotics, many of the  
437 underlying principles have been indicated long ago for various tetracyclines [5, 13, 19-21, 35-  
438 37]. However, those leads seem to have been largely ignored in favour of certain postulations  
439 from *in vitro* studies [6, 18]. Nevertheless, this work would serve as a basis for further studies  
440 with other tetracycline antibiotics in this perspective. When correlated with their effects on  
441 non-bacterial and eukaryotic rRNA processing and non-infectious disease conditions, the  
442 molecular mechanism of action of the tetracyclines would be more definitively elucidated.

#### 443 **Acknowledgments**

444 This work was supported by the Commonwealth Scholarship Commission in the UK.

#### 445 **References**

- 446 1 Goldman RA, Hasan T, Hall CC, Strycharz WA, Cooperman BS. Photoincorporation of  
447 Tetracycline into Escherichia-Coli Ribosomes - Identification of the Major Proteins  
448 Photolabeled by Native Tetracycline and Tetracycline Photoproducts and Implications for the  
449 Inhibitory-Action of Tetracycline on Protein-Synthesis. *Biochemistry* 1983; 22: 359-368.
- 450  
451 2 Chopra I, Roberts M. Tetracycline antibiotics: Mode of action, applications, molecular  
452 biology, and epidemiology of bacterial resistance. *Microbiology and Molecular Biology  
453 Reviews* 2001; 65: 232-260.
- 454  
455 3 Schnappinger D, Hillen W. Tetracyclines: Antibiotic action, uptake, and resistance  
456 mechanisms. *Archives of Microbiology* 1996; 165: 359-369.
- 457  
458 4 Chopra I, Hawkey PM, Hinton M. Tetracyclines, Molecular and Clinical Aspects. *Journal of  
459 Antimicrobial Chemotherapy* 1992; 29: 245-277.
- 460  
461 5 Day LE. Tetracycline inhibition of cell-free protein syntesis I. Binding of tetracycline to  
462 components of the system. *Journal of Bacteriology* 1966; 91: 1917-1923.
- 463  
464 6 Oehler R, Polacek N, Steiner G, Barta A. Interaction of tetracycline with RNA:  
465 photoincorporation into ribosomal RNA of Escherichia coli. *Nucleic Acids Res* 1997; 25: 1219-  
466 1224.

467  
468 7 Noah JW, Dolan MA, Babin P, Wollenzien P. Effects of Tetracycline and Spectinomycin on the  
469 Tertiary Structure of Ribosomal RNA in the Escherichia coli 30 S Ribosomal Subunit. *J Biol*  
470 *Chem* 1999; 274: 16576-16581.

471  
472 8 Chukwudi CU, Good L. Interaction of the tetracyclines with double-stranded RNAs of random  
473 base sequence: new perspectives on the target and mechanism of action. *J Antibiot (Tokyo)*  
474 2016; 69: 622-630.

475  
476 9 Yi H, Cho Y-J, Won S, Lee J-E, Jin Yu H, Kim S *et al.* Duplex-specific nuclease efficiently  
477 removes rRNA for prokaryotic RNA-seq. *Nucleic Acids Research* 2011; 39: e140-e140.

478  
479 10 Simons rw, Grunberg-Manago m. *RNA structure and function*. Cold Spring Harbor Laboratory  
480 Press, 1998.

481  
482 11 Srivastava AK, Schlessinger D. Mechanism and regulation of bacterial ribosomal RNA  
483 processing. *Annu Rev Microbiol* 1990; 44: 105-129.

484  
485 12 Dunn JJ, Studier FW. T7 early RNAs and Escherichia coli ribosomal RNAs are cut from large  
486 precursor RNAs in vivo by ribonuclease 3. *Proc Natl Acad Sci U S A* 1973; 70: 3296-3300.

487  
488 13 Nikolaev N, Schlessinger D, Wellauer PK. 30 S pre-ribosomal RNA of Escherichia coli and  
489 products of cleavage by ribonuclease III: length and molecular weight. *J Mol Biol* 1974; 86:  
490 741-747.

491  
492 14 King TC, Schlessinger D. S1 nuclease mapping analysis of ribosomal RNA processing in wild  
493 type and processing deficient Escherichia coli. *Journal of Biological Chemistry* 1983; 258:  
494 12034-12042.

495  
496 15 Gegenheimer P, Apirion D. Processing of procaryotic ribonucleic acid. *Microbiol Rev* 1981; 45:  
497 502-541.

498  
499 16 Nakashima N, Tamura T, Good L. Paired termini stabilize antisense RNAs and enhance  
500 conditional gene silencing in Escherichia coli. *Nucleic Acids Res* 2006; 34: e138.

501  
502 17 Mironov KS, Los DA. RNA Isolation from *Synechocystis*. *Bio-protocol* 2015; 5: e1428.

503  
504 18 Chukwudi CU. Ribosomal RNA binding sites and the molecular mechanism of action of the  
505 tetracyclines. *Antimicrobial Agents and Chemotherapy* 2016; 60: 4433-4441.

506

507 19 Rye RM, Wiseman D. Cell size changes during the growth of Escherichia coli partially  
508 inhibited by some antibacterial agents. *J Pharm Pharmacol* 1968; 20 Suppl: 8S-13S.

509  
510 20 Holmes IA, Wild DG. The synthesis of ribonucleic acid during inhibition of Escherichia coli by  
511 chlortetracycline. *Biochem J* 1965; 97: 277-283.

512  
513 21 Holmes IA, Wild DG. Inhibition of the growth of Escherichia coli by chlortetracycline.  
514 *Biochem J* 1967; 104: 679-685.

515  
516 22 Apirion D, Gegenheimer P. Processing of bacterial RNA. *FEBS Lett* 1981; 125: 1-9.

517  
518 23 Guespin-Michel J, Kaufman M. Positive Feedback Circuits and Adaptive Regulations in  
519 Bacteria. *Acta Biotheoretica* 2001; 49: 207-218.

520  
521 24 Ray JCJ, Igoshin OA. Adaptable Functionality of Transcriptional Feedback in Bacterial Two-  
522 Component Systems. *PLoS Comput Biol* 2010; 6: e1000676.

523  
524 25 Todar K. Regulation and Control of Metabolism in Bacteria. *Online Textbook of Bacteriology*  
525 [www.textbookofbacteriology.net](http://www.textbookofbacteriology.net), 2008.

526  
527 26 Nikolay R, Schmidt S, Schlömer R, Deuerling E, Nierhaus KH. Ribosome Assembly as  
528 Antimicrobial Target. *Antibiotics (Basel)* 2016; 5.

529  
530 27 King TC, Sirdeshmukh R, Schlessinger D. RNase III cleavage is obligate for maturation but not  
531 for function of Escherichia coli pre-23S rRNA. *Proceedings of the National Academy of*  
532 *Sciences of the United States of America* 1984; 81: 185-188.

533  
534 28 Srivastava AK, Schlessinger D. Processing pathway of Escherichia coli 16S precursor rRNA.  
535 *Nucleic Acids Res* 1989; 17: 1649-1663.

536  
537 29 Gegenheimer P, Watson N, Apirion D. Multiple pathways for primary processing of  
538 ribosomal RNA in Escherichia coli. *Journal of Biological Chemistry* 1977; 252: 3064-3073.

539  
540 30 Babitzke P, Granger L, Olszewski J, Kushner SR. Analysis of mRNA decay and rRNA processing  
541 in Escherichia coli multiple mutants carrying a deletion in RNase III. *J Bacteriol* 1993; 175:  
542 229-239.

543  
544 31 Agwuh KN, MacGowan A. Pharmacokinetics and pharmacodynamics of the tetracyclines  
545 including glycylicyclines. *J Antimicrob Chemother* 2006; 58: 256-265.

546

547 32 Cunha BA, Domenico P, Cunha CB. Pharmacodynamics of doxycycline. *Clin Microbiol Infect*  
548 2000; 6: 270-273.

549  
550 33 De Stasio EA, Moazed D, Noller HF, Dahlberg AE. Mutations in 16S ribosomal RNA disrupt  
551 antibiotic--RNA interactions. *EMBO J* 1989; 8: 1213-1216.

552  
553 34 Eichler DC, Craig N. Processing of eukaryotic ribosomal RNA. *Prog Nucleic Acid Res Mol Biol*  
554 1994; 49: 197-239.

555  
556 35 Wei Y, Bechhofer DH. Tetracycline induces stabilization of mRNA in *Bacillus subtilis*. *Journal*  
557 *of Bacteriology* 2002; 184: 889-894.

558  
559 36 Amin AR, Attur MG, Thakker GD, Patel PD, Vyas PR, Patel RN *et al*. A novel mechanism of  
560 action of tetracyclines: effects on nitric oxide synthases. *Proc Natl Acad Sci U S A* 1996; 93:  
561 14014-14019.

562  
563 37 Atherly AG. Specific inhibition of ribosomal RNA synthesis in *Escherichia coli* by tetracycline.  
564 *Cell* 1974; 3: 145-151.

565  
566 38 Cangelosi GA, Brabant WH. Depletion of pre-16S rRNA in starved *Escherichia coli* cells. *J*  
567 *Bacteriol* 1997; 179: 4457-4463.


568  
569  
570

571 **TABLES**572 **Table 1: Hybridization probes used for pre-rRNA northern blotting**

<b>Probe/primer</b>	<b>Target description*</b>	<b>Length</b>	<b>Sequence</b>	<b>Reference</b>
<b>ECR2</b>	Mature 16S rRNA	28	5'-gtccccctcttggcttgcgacgttat-3'	[38]
<b>ECPR2</b>	3' pre-16S rRNA tail (rrnA, -D, -G, -H)	30	5'-gtgtgagcactgcaaagtagccttctttaa-3'	[38]
<b>CCPR1</b>	Pre-16S rRNA (3' rrnA, -H; 5' -D, -G)	50	5'-cctgtagaggtttactgctcatttca tcagacaatctgtgtgagcact-3'	This work
<b>CCPR2</b>	Pre-rRNA (3' end of 16S to 5' end of 23S)	457	*	This work
<b>CCPR2 forward primer</b>	3' end of 16S rRNA	22	5'-cacctccttacctaaagaagc-3'	This work
<b>CCPR2 reverse primer</b>	5' end of 23S rRNA	19	5'-tcgcttaacctcacaacc-3'	This work

573 \*See Fig 1B for illustration of region of complementarity with target pre-rRNA.

574




576

577 **Fig 1: Schematic representation of the primary transcript of ribosomal RNA of *E. coli*.**


578 (A) Mature rRNA sequences are indicated as bold line loops, and dsRNA within the  
 579 precursor sequences represented by the stems (not drawn to scale). Arrows indicate proposed  
 580 regions of double-stranded primary transcript RNase III cleavage sites, where cleavage  
 581 releases the pre-16S and pre-23S rRNAs for further maturation to produce mature rRNAs.  
 582 The \* and \*\* symbols indicate the number of tRNA molecules within the operon at the  
 583 indicated sites. \* = 1 - 2, \*\* = 0 - 4. In addition, the *rrnD* in *E. coli* has two genes encoding  
 584 5S rRNA. (B) Target position of the hybridization probes in relation to the mature ribosomal  
 585 RNAs in the long primary rRNA transcript of *E. coli*.

586



587

588 **Fig 2: Effect of doxycycline on mature rRNA amounts and rRNA sizes in growing**  
 589 **bacteria cells over time.** Northern blot membrane stained with methylene blue (A) of total  
 590 RNA extract from *E. coli* cells growing in the absence and presence of 100µM doxycycline at  
 591 various time points during growth showing the 23S and 16S rRNAs, and the hybridized  
 592 membrane blot (B) showing rRNA primary transcript and pre-rRNA (that indicate continued  
 593 transcription of rRNA) and smearing of the pre-rRNA in the presence of doxycycline.  
 594 Graphical analysis of the rRNA band intensities (C,D) show significantly decreasing amounts  
 595 of 16S and 23S rRNAs with time ( $P= 0.0046$ ,  $r = -0.7365$  for 23S, and  $P= 0.0091$ ,  $r = -$   
 596  $0.8126$  for 16S rRNA) in cells that were grown in medium containing doxycycline, when  
 597 compared to the increasing amounts of the rRNAs in cells growing without the drug. RFU=  
 598 Relative fluorescence unit.




600

601 **Fig 3: Effect of increasing concentrations of doxycycline on mature rRNA formation in**  
 602 **growing bacterial cells.** EtBr-stained denaturing agarose gel image (A) of total RNA  
 603 extracted from *E. coli* cells grown in increasing concentrations of doxycycline (0-200μM) at  
 604 20 min and 120 min incubation periods, showing decreasing amounts of 23S and 16S

605 ribosomal RNAs with increasing concentration of doxycycline as illustrated in the graphs (C,  
 606 D), hybridized membrane blot of the gel (B) showing smearing of the RNA bands at 120 min,  
 607 growth curve (E) and total RNA concentration of the samples (F).  $IC_{50} = 8.327\mu M$  (+/-SE  
 608 2.465,  $R^2 = 0.9554$ ) at 20 min incubation time and  $76.51\mu M$  (+/-SE 49.6,  $R^2 = 0.8947$ ) at  
 609 120min. RFU= Relative fluorescence unit, T=rRNA transcript.

610




611

612 **Fig 4: Effect of increasing concentrations of doxycycline on the processing of *E. coli***  
 613 **rRNA *in vivo*.** (A) EtBr-stained denaturing agarose gel of total RNA extract from *E. coli* K-  
 614 12 cells grown in increasing concentrations of doxycycline. Samples were collected 20 min  
 615 after treating with doxycycline, and total RNA extracted from the samples were fractionated  
 616 in 1% denaturing agarose gel. S=starting culture sample at 0 min, 0=untreated culture at

617 20mins, M=NEB log 2-log DNA ladder (0.1-10.0 kb). (B) Northern blot nylon membrane  
618 hybridized with pre-rRNA probe CCPR2 showing the initial rRNA transcript (T) and long  
619 pre-rRNA. (C) Graphical presentation of the optical density of cultures and the concentration  
620 of total RNA extracted from them, showing no significant change in culture OD ( $P= 0.7745$ ),  
621 and a slight decrease in RNA conc, from 50  $\mu\text{M}$  doxycycline conc. S=starting culture sample  
622 at 0 min, 0=untreated culture at 20mins. (D) Graph of densitometric analysis of the various  
623 rRNA bands in B. The blot and graph show a dose-dependent increase in the long pre-rRNAs  
624 and concurrent decrease in 16S and 23S rRNAs with increasing doxycycline concentrations.  
625 RFU= Relative fluorescence unit. Inset shows percentage contribution of each rRNA species,  
626 as detected with the probe.

627



628

629 **Fig 5: Effect of doxycycline on nucleoid morphology of *E. coli*.** Fluorescent microscopy  
630 images of *E. coli* K-12 cells treated with 0-200 $\mu$ M doxycycline and incubated for 20 min  
631 before sample collection and processing for microscopy. Cells appear elongated at 20 and 50  
632  $\mu$ M doxycycline, with nucleoid degeneration at 100-200 $\mu$ M. (x630).

633

634