RVC OPEN ACCESS REPOSITORY – COPYRIGHT NOTICE

This is the author's version of the work. It is posted here by permission of the AAAS for personal use, not for redistribution. The definitive version was published in *Science*, Vol. 362, Issue 6411, pp. 165-166 (12 October, 2018), DOI: <u>https://doi.org/10.1126/science.aav4096</u>.



Letter for Science

<u>Title</u>: PPR virus threatens wildlife conservation

Authors:

Xavier Fernandez Aguilar¹,*, Amanda E. Fine², Mathieu Pruvot², Felix Njeumi³, Christian Walzer^{2,4}, Richard Kock¹, Enkhtuvshin Shiilegdamba²

Author institutions:

1 Department of Pathobiology and Population Sciences, Royal Veterinary College, London NW1 0TU, UK.

2 Wildlife Conservation Society, Bronx, NY 10460, USA.

3 Food and Agriculture Organization of the United Nations, FAO, 00153 Rome, Italy.

4 Research Institute of Wildlife Ecology, University of Veterinary Medicine, A-1160 Vienna, Austria.

* Corresponding author. Email: <u>xfdezaguilar@gmail.com</u>

Peste des petits ruminants (PPR), a viral disease that affects domestic small ruminants with high morbidities and mortalities across more than 70 countries, engenders a global cost of US\$2.1 billion and compromises livelihoods of some 900 million poor and low-income people (1, 2). PPR has generally been regarded globally as only a livestock problem (1), but it is also a wildlife conservation challenge.

PPR has spread from its historic range of Africa, West Asia, and the Middle East to the vast and remote steppes and mountains of Eastern Asia, threatening even more wildlife. The mass mortality event affecting more than two-thirds of the critically endangered Mongolian saiga (Saiga tatarica mongolica) population in 2017 is one example of PPR's reach (3). PPR has also been diagnosed in mountain ungulate mortalities in the Middle East (4) and detected along the Himalayas (5) and the Tian Shan and Altai ranges (6).

PPR clearly threatens saiga populations, but its impact on other steppe and mountain ungulates of Asia is unknown. The inaccessibility of their habitats, together with the lack of wildlife health surveillance programs across large areas, contributes to underestimating and underreporting of PPR mortality events. The circulation of PPR in Asia may have grave consequences in wild populations that already struggle with overhunting, poaching, livestock competition, and stochastic climatic events. For example, in 2018, multiple PPR-related mortalities of the Siberian ibex (Capra sibirica) were recorded in Mongolia (7), and PPR-related mortalities of wild ungulates were found in Iran (8). PPR is also of special concern for the conservation of the susceptible markhor (Capra falconeri), argali (Ovis ammon), and goitered gazelle (Gazella subgutturosa), considered vulnerable by the IUCN (9), and for the survival of the snow leopard (Panthera uncia), which relies on prey abundance. A robust assessment will require better baseline information on population size and trends. More efforts to conduct PPR surveillance and integrate wildlife protection into control strategies are urgently required.

References

- 1. "Global strategy for the control and eradication of PPR" (Tech. Rep. FAO-OIE, 2015).
- 2. B. Jones et al., PLOS ONE 11, e0149982 (2016).
- 3. R. Kock, "Investigation of peste des petits ruminants (PPR) among wild animals and its potential impact on the current PPR situation in livestock" (Crisis Management Centre for Animal Health, Tech. Rep. FAO-OIE, 2017).
- 4. M. Marashi et al., Emerg. Infect. Dis. 23, 704 (2017).
- 5. J. Bao et al., China Res. Vet. Sci. 90, 238 (2011).
- 6. J. Li et al., Transbound. Emerg. Dis. 64, e43 (2017).
- 7. Science for Nature and People Partnership (SNAPP) team: Steppe Health (<u>https://snappartnership.net/teams/steppe-health/</u>).
- 8. ProMED-mail, "Peste des petits ruminants Iran: (AL) wildlife" (Archive number 20180928.6058637, 28 September 2018); www.promedmail.org
- 9. The IUCN Red List of Threatened Species, Gazella subgutturosa (www.iucnredlist.org/details/8976/0).