

This author's accepted manuscript may be used for non-commercial purposes in accordance with [Wiley Terms and Conditions for Self-Archiving](#).

The full details of the published version of the article are as follows:

TITLE: C7 vertebra homeotic transformation in domestic dogs – are Pug dogs breaking mammalian evolutionary constraints?

AUTHORS: Brocal, J; De Decker, S; José-López, R; Manzanilla, E G; Penderis, J; Stalin, C; Bertram, S; Schoenebeck, J J; Rusbridge, C; Fitzpatrick, N; Gutierrez-Quintana, R

JOURNAL: Journal of Anatomy

PUBLISHER: Wiley

PUBLICATION DATE: 14 May 2018 (online)

DOI: <https://doi.org/10.1111/joa.12822>

1 C7 VERTEBRA HOMEOTIC TRANSFORMATION IN DOMESTIC DOGS- ARE PUG
2 DOGS BREAKING MAMMALIAN EVOLUTIONARY CONSTRAINTS?

3 J. Brocal¹, S. De Decker², R. José-López¹, E.G. Manzanilla³, J. Penderis⁴, C. Stalin¹, S.
4 Bertram², J.J. Schoenebeck⁵, C. Rusbridge^{6,7}, N. Fitzpatrick⁶, R. Gutierrez-Quintana^{1*}

5
6 ¹School of Veterinary Medicine, College of Medical, Veterinary and Life Sciences,
7 University of Glasgow, Bearsden Road, Glasgow G61 1QH, UK.

8 ² Department of Veterinary Clinical Science and Services, The Royal Veterinary College,
9 University of London, Hawkshead Lane, North Mymms, Hertfordshire AL97TA, UK.

10 ³Teagasc Animal and Grassland Research and Innovation Centre, Moorepark, Fermoy, Co.
11 Cork, P61 C996, Ireland.

12 ⁴Vet-Extra Neurology, Broadleys Veterinary Hospital, Craig Leith Road, Stirling, FK7 7LE,
13 UK.

14 ⁵Royal (Dick) School of Veterinary Studies and Roslin Institute, University of Edinburgh,
15 Easter Bush, Midlothian EH25 9RG, UK.

16 ⁶Fitzpatrick Referrals, Eashing, Surrey GU7 2QQ, UK.

17 ⁷School of Veterinary Medicine, Faculty of Health & Medical Sciences, University of Surrey,
18 Guildford, Surrey, United Kingdom.

19
20

21 *Corresponding author: College of Medical, Veterinary and Life Sciences, University of
22 Glasgow, Bearsden Road, Glasgow G61 1QH, UK phone: +44(0)141 330 5848, fax: +44
23 (0)141 330 3663, email: Rodrigo.GutierrezQuintana@glasgow.ac.uk

24

25 Key words: vertebral malformation, dog, pug, homeotic, cervical ribs, cervical transitional
26 vertebra

27

28

29 **Abstract**

30 The number of cervical vertebrae in mammals is almost constant at seven, regardless of their
31 neck length, implying that there is selection against variation in this number. Homeobox (*Hox*)
32 genes are involved in this evolutionary mammalian conservation and homeotic
33 transformation of cervical into thoracic vertebrae (cervical ribs) is a common phenotypic
34 abnormality when *Hox* gene expression is altered. This relatively benign phenotypic change
35 can be associated with fatal traits in humans. Mutations in genes upstream of *Hox*, inbreeding
36 and stressors during organogenesis can also cause cervical ribs. The aim of this study was to
37 describe the prevalence of cervical ribs in a large group of domestic dogs of different breeds
38 and explore a possible relation with other congenital vertebral malformations (CVM) in the
39 breed with the highest prevalence of cervical ribs. By phenotyping we hoped to give clues as
40 to the underlying genetic causes. Twenty computed tomography studies from at least two
41 breeds belonging to each of the nine groups recognised by the Federation Cynologique
42 Internationale, including all the brachycephalic “screw-tailed” breeds that are known to be
43 overrepresented for CVM’s were reviewed. The Pug dog was more affected by cervical ribs
44 than any other breed (46%) ($p < 0.001$) and was selected for further analysis. No association
45 was found between the presence of cervical ribs and vertebral body formation defect, bifid
46 spinous process, caudal articular process hypoplasia/aplasia and an abnormal sacrum, which
47 may infer they have a different etiopathogenesis. However, Pug dogs with cervical ribs were
48 more likely to have a transitional thoraco-lumbar vertebra ($p = 0.041$) and a pre-sacral
49 vertebral count of 26 ($p < 0.001$). Higher C7/T1 dorsal spinous processes ratios were
50 associated with the presence of cervical ribs ($P < 0.001$) supporting this is a true homeotic
51 transformation. Relaxation of the stabilizing selection has likely occurred and the Pug dog
52 appears to be a good naturally occurring model to further investigate the aetiology of cervical
53 ribs, other congenital vertebral anomalies and numerical alterations.

54 **Introduction**

55 The number of cervical vertebrae in mammals is almost constant at seven, regardless of their
56 neck length (Schulz, 1961; Leboucq, 1895; Starck, 1978; Badlangana et al. 2009).

57 Interestingly, the vertebral count in other regions does vary among mammals, implying there
58 is a selection against variation in the number of cervical vertebrae (Schulz, 1961; Leboucq,
59 1895).

60 Manatees (*Trichechus*) and sloths (*Bradypus* and *Choloepus*) are exceptions to this
61 evolutionary rule (Buchholtz & Stepien, 2009; Varela-Lasheras et al. 2011; Buchholtz et al.
62 2007). This relaxation of stabilizing selection leading to persistence of features, against
63 which there is normally a strong selection, can also be found in domesticated mammals
64 through artificial selection. Due to the relaxation of stabilizing selection through human care,
65 dogs with polydactyly can breed and reproduce despite the strong evolutionary constraint on
66 polydactyly in amniotes (Galis et al. 2001; Galis et al. 2010).

67 Homeobox (*Hox*) genes are involved with this evolutionary mammalian conservation and have
68 proven to be critical regulators in the establishment of axial skeleton morphology during
69 embryogenesis in all vertebrae classes (Krumlauf, 1994; Wellik 2007). It is generally
70 assumed that shifts of the boundary of vertebral regions are associated with shifts of *Hox*
71 expression patterns and therefore, homeotic transformations of cervical into thoracic
72 vertebrae (*C7* transitional vertebra commonly known in humans as cervical ribs) is a common
73 phenotypic abnormality. Abnormal expression of *Hox* genes does not just lead to vertebral
74 column anomalies, but also neural problems, increased susceptibility to early childhood
75 cancer, other congenital anomalies and stillbirths (Galis, 1999; Hostikka et al. 2009; Furtado
76 et al. 2011; ten Broek et al 2012; Schumacher, 1992; Gladstone et al. 1932). Upstream

77 organizers of *Hox* genes are equally crucial in the head-to-tail patterning of the vertebral
78 column

79 An extreme selection in humans against homeotic transformations of cervical vertebrae
80 occurs (ten Broek et al 2012) and the total incidence of cervical ribs in people is low (0.04-
81 6.2%) (Henderson, 1913; Spadlinski et al. 2016). Two previous studies in dogs have
82 described congenital vertebral malformations of the whole vertebral column in 145 and 228
83 cases submitted to necropsy and reported a cervical ribs prevalence that varied from 2.1% to
84 11.4% (Morgan, 1968; Breit and Kunzel 1998). These studies date from 1968 and 1998 and
85 the dog population has changed in size, genetic diversity, breed prevalence and phenotypic
86 characteristics over time (The Kennel Club 2016, Vila et al. 1999)

87 Normal dogs have seven cervical, 13 thoracic, seven lumbar, three sacral and a variable
88 number of caudal vertebrae. The first and second vertebrae present easily distinct
89 recognisable anatomical features. The rest of cervical vertebrae differ slightly from each
90 other, except for the expanded sagittal plate-like transverse processes of C6. Thoracic
91 vertebrae have shorter bodies compared to cervical and lumbar vertebrae and articulate with a
92 rib. The spinous process of the first thoracic vertebra is the longest of all vertebrae, is its most
93 conspicuous feature and its length gradually decreases with successive vertebrae (Evans and
94 de Lahunta, 2013) (**Figure 1**).

95 Congenital vertebral malformations (CVM) are relatively common in brachycephalic “screw-
96 tailed” breed dogs, such as the French bulldogs, the English bulldogs, Boston terriers and Pug
97 dogs-(Gutierrez-Quintana et al. 2014; Steffen et al. 2004; Wrzosek et al. 2014). Unlike
98 CVM’s in other less frequently seen regions, thoracic CVM’s have been described and
99 compared in some brachycephalic dogs. Overall, Pug dogs had significantly fewer vertebral
100 body formation defects and more thoracolumbar transitional vertebrae and bifid spinous

101 process compared to the other two breeds. Despite having fewer vertebral body formation
102 defects, when present, these had greater clinical relevance. Furthermore, all French and
103 English bulldogs in the study had a normal thoracic vertebral count, while 25% of the Pug
104 dogs had 12 instead of 13 thoracic vertebrae (Ryan et al. 2017). Therefore, the Pug dog
105 appeared to be a good naturally occurring canine model to phenotype and further investigate
106 the genetic background of CVM's.

107 The aim of this study was to describe the frequency of cervical ribs in a large group of
108 domestic dogs of different breeds. We were particularly interested in breeds reported to be
109 commonly affected by CVM's and we wanted to explore a possible relation between them in
110 a naturally occurring model, providing clues as to their genetic causes. Given the high
111 number of thoraco-lumbar transitional vertebrae previously identified in Pug dogs (Ryan et
112 al. 2017), we hypothesised that a high number of cervical ribs compared to other dog breeds
113 and species would be identified. We also hypothesised that having cervical ribs would be
114 associated with a higher incidence of other CVM's.

115 **Material and methods**

116 Ethical approval from the University of Glasgow, School of Veterinary Medicine was obtained
117 for this study.

118 To quantify the frequency of cervical ribs in different breeds of domestic dogs, the medical
119 records of the Royal Veterinary College, University of Glasgow Small Animal Hospital, Vet
120 Extra Neurology and Fitzpatrick Referrals were retrospectively reviewed to identify
121 computed tomographic studies (CTs) that included the C7 vertebra regardless of their clinical
122 indication. A total of 19 breeds with at least two breeds belonging to each of the nine groups
123 recognised by the Federation Cynologique Internationale (FCI) (FCI 2017) were examined.
124 These included Basset hounds, Beagle, Border collie, Chow-chow, Cocker spaniel, German

125 shepherd dog, German pointer, Greyhound, Hungarian Vizsla, Husky, Labrador retriever,
126 Dachshund, Staffordshire bull terrier, Whippet, West Highland white terrier and all
127 brachycephalic “screw-tailed” breeds (French bulldog, the English bulldog, Boston terrier
128 and Pug dog) known to be overrepresented for CVM’s were selected for analysis (Gutierrez-
129 Quintana et al. 2014; Ryan et al. 2017). A maximum of 20 randomly selected CTs were
130 reviewed for each breed. Dogs underwent imaging investigations under sedation or general
131 anaesthesia for a variety of clinical indications (related or unrelated to spinal disease). The
132 breed having the highest frequency of cervical ribs was then selected for further analysis. All
133 the available CTs that included the entire vertebral column (including or excluding coccygeal
134 vertebrae) or C7 and the thoracic region were then reviewed and the signalment (age, breed,
135 sex and colour) was recorded when available. We also retrieved and reviewed orthogonal
136 radiographic views of dogs with thoracic CVM’s causing neurological deficits from the breed
137 that was selected for further analysis.

138 Computed tomographic studies and radiographs were reviewed by a board-certified
139 neurologist (RG) and a third-year neurology resident (JB). Images were displayed using an
140 open-source Workstation DICOM viewer (Osirix Imaging Software, v 3.9.2, Pixmeo, Geneva,
141 Switzerland). Congenital vertebral malformations were identified and classified into vertebral
142 body formation defects (commonly known as hemivertebra) (Gutierrez-Quintana et al. 2014),
143 transitional vertebrae and dorsal process formation defects (e.g. bifid spinous process or
144 fusion of the dorsal spinal process). Ambiguous vertebrae located at the border between two
145 segments of the vertebral column and possessing characteristics of both segments were
146 considered to be transitional vertebra (Simoens, 1983). The total number of pre-sacral
147 vertebrae was recorded and the sacrum was classified as being normal or abnormal. The
148 caudal articular processes from T9-L2 of whole body CTs were classified as normal or
149 hypoplastic/aplastic. All dogs were considered to have seven cervical vertebrae, even if two

150 large ribs were identified at C7. When identified on CT, cervical ribs were classified as being
151 either short (less than 1.5cm) or long (more than 1.5cm), and with or without articulation.
152 One of the key morphological differences between cervical and thoracic vertebrae is the
153 comparatively long dorsal spinous process in the latter. To test whether other aspects of the
154 C7 vertebrae took on thoracic vertebrae characteristics, the heights of the dorsal spinous
155 processes of C7 and T1 vertebrae were measured and the C7/T1 dorsal spinous process
156 height ratio was calculated. When identified on radiographs, the presence or absence of
157 cervical ribs was recorded with no further sub classification. Thoraco-lumbar vertebrae were
158 considered thoracic if they were transitional (including transverse process asymmetry) and
159 lumbar if they had bilateral and normal size symmetric transverse processes. Dogs in which
160 L7 was considered to have fused with the sacrum (absent intervertebral disc) were considered
161 to have six lumbar vertebrae.

162 **Statistical Analysis**

163 All analyses were done using SAS 9.4 (SAS Inst. Inc., Cary, NC). Data was analysed using
164 non-parametric methods (Wilcoxon-Mann-Whitney, Kruskal-Wallis and Chi-square/Fisher
165 tests, and Spearman correlation) except for the analysis of C7/T1 dorsal spinous process ratio
166 depending on the presence of transitional vertebrae, cervical ribs and joints. The later was
167 analysed using t-test. Alpha level for determination of significance was 0.05 and multiple
168 comparison were analysed using Bonferroni correction. Given that CT provides better bone
169 definition compared to radiography, data from orthogonal radiographic views of dogs with
170 thoracic CVM's causing neurological deficits was analysed separately and only to study the
171 relation between CVM's.

172 **Results**

173 *Frequency of cervical ribs in domestic pure breed dogs*

174 Twenty CTs were available for review from each of the breeds mentioned above except for
175 the Husky (n=15), the Basset hound (n=15), the German pointer (n=13), the Miniature
176 Schnauzer (n=6) and the Chow-chow (n=4). The percentage of dogs with cervical ribs from
177 each of the reviewed breeds is illustrated in **Figure 2**. Cervical ribs were more common in the
178 Pug dog compared to any other breed and was therefore selected for further analysis (45/97-
179 46%) (P<0.001). The overall prevalence of cervical ribs was 14% (61/430). Excluding the
180 Pug dog the percentage decreased to 4.8% (16/333).

181 *Further analysis in the breed with highest frequency of cervical ribs: the Pug dog*

182 A total of 51 CTs of the whole vertebral column and 46 cervico-thoracic CTs from Pug dogs
183 with no clinical signs caused by CVM's (scanned for respiratory signs (n=81), metastatic
184 check (n=6), orthopaedic problems (n=6) and gastrointestinal signs (4)) and 33 thoracic
185 radiographs from Pug dogs with neurological deficits caused by CVM's (radiographed for
186 T3-L3 myelopathy) Pug dogs were available and reviewed (n=130 Pug dogs).

187 A total of 57/130 Pug dogs had cervical ribs, 50/130 had vertebral body formation defects
188 (**Figure 3**), 48/130 had bifid spinous process (without myelomeningocele) (**Figure 4**), 52/130
189 had a thoraco-lumbar transitional vertebra (**Figure 5**), 12/51 had a lumbo-sacral transitional
190 vertebra (**Figure 6**) and 47/51 had caudal articular process hypoplasia/aplasia (**Figure 7**).
191 T13 was absent in 31/130 and L7 was absent in 17/51. Out of 51 whole vertebral column
192 CTs, twenty-eight (55%) had 26 pre-sacral vertebrae, 23 (45%) had 27 pre-sacral vertebrae
193 and eleven had no transitional vertebrae (**Table 1**).

194 Mean and median age were 38 and 31 months respectively. Sixty-one Pug dogs were female
195 and 69 were male. Females (30/45-65%) were affected more frequently by cervical ribs than

196 males (15/45-29%) (P=0.012). Coat colour was only available for Pug dogs with a whole-
197 body CT. No association between coat colour and the presence of any transitional vertebrae,
198 vertebral body formation defects or a bifid spinous process was identified. In most cases,
199 cervical ribs were bilateral (32/45-71%). Among cases with unilateral changes (13/45-29%),
200 the left (10/13-77%) side was more frequently affected compared to the right (3/13-23%)
201 (P=0.04; **Table 2**). Cervical ribs were classified as short with no joint in 35% (27/77) of the
202 cases, long with no joint in 2.5% (2/77), short with a joint in 34% (26/77) and long with a
203 joint in 28.5% (22/77) of the cases (**Figure 8**).

204 No association was found between the presence of vertebral body formation defects,
205 transitional vertebra, bifid spinous process, caudal articular process hypoplasia/aplasia and an
206 abnormal sacrum in the CTs of Pug dogs with no clinical signs associated to CVM's or in the
207 radiographs of Pug dogs with CVM's causing neurological deficits. Pug dogs in which
208 cervical ribs were identified on CTs were more likely to have a thoraco-lumbar transitional
209 vertebra (P= 0.041) than those without cervical ribs, though this was not replicated in our
210 analysis of radiographs. Pug dogs with cervical ribs (P=0.025) or a lumbo-sacral transitional
211 vertebra (P<0.001) were also more likely to have a pre-sacral vertebral count of 26 instead of
212 27. The pre-sacral vertebral count was not associated with the presence of hemivertebrae,
213 bifid spinous process or an abnormal sacrum. Longer C7 dorsal spinal processes and higher
214 C7/T1 ratios were associated with the presence of cervical ribs (P<0.001) and were correlated
215 (r=0.44 and r = 0.55 respectively) with the added length of left and right cervical ribs (**Figure**
216 **9**). Higher C7/T1 ratios were also associated with having fewer vertebrae affected by caudal
217 articular process hypoplasia/aplasia (r = -0.36; P = 0.010).

218 **Discussion**

219 The previously reported prevalence of cervical ribs in domestic dogs varied from 2.1% to
220 11.9% (Morgan, 1968; Breit and Kunzel 1998). The present study investigated a larger
221 number of dogs equally distributed between breeds and found an overall similarly low
222 prevalence, but identified the Pug dog as being significantly more affected than any other
223 breed. The German Shepherd (15%), the Hungarian Vizsla (15%) and the Boxer (10%) also
224 exhibited unusually high percentages of cervical ribs.

225

226 It has been proposed that cervical ribs may be largely composed of ossification sites that
227 disappear after birth (Chernoff and Rogers, 2004). One of the key morphological differences
228 between cervical and thoracic vertebrae is the comparatively long dorsal spinous process in
229 the latter. Longer C7 dorsal spinous processes and high C7/T1 dorsal spinous process height
230 ratios were associated with the presence of cervical ribs in this study, suggesting this is a true
231 homeotic transformation of a cervical vertebra into a thoracic one and not an unmasking of
232 vestigial cervical ribs left behind in evolution. The Spearman correlation between C7/T1 ratio
233 and the added length of left and right cervical ribs was moderate, suggesting that longer
234 cervical ribs are not always associated with longer C7 dorsal spinous process. In combination
235 with the length of the cervical ribs, this ratio may be used as an objective measure of the
236 degree of C7 homeotic transformation.

237 In humans, bilateral cervical ribs are the most common; and when unilateral, they are
238 traditionally considered to be more frequently left-sided. Although inconsistent results have
239 been reported, females seem to be more frequently affected (Furtado et al. 2011; Galis et al
240 2006; Bots et al. 2011, Schumacher et al. 1992). The results of this study were in accordance
241 with these findings. Deficient signalling in mice leads to delayed somite formation on the
242 right side, causing a diminished coordination between the development of the left and right

243 somites, and leading to a differential expression of the *Hox* genes (Vermot et al. 2005;
244 Griesinger et al. 2005).

245 Mammals under harsh conditions and stressors like hyperthermia or exposure to boric acid
246 during organogenesis (Chernoff and Rogers, 2004; Steigenga et al. 2006; Wery et al. 2003, Li
247 and Shiota, 1999) and mutations that influence *Hox* gene expression can also cause cervical
248 ribs. *T*-gene (also known as brachyury) knockdown in vivo experiments cause T1
249 transformation of cervical vertebrae in 30% of mice (Yamaguchi, 1999; Pennimpede et al.
250 2012) and a spontaneous mutation in this same gene, has been associated with a cervical
251 vertebral homeotic transformation in the *Bos Taurus* with vertebral and spinal dysplasia
252 (VSD) (Kromic et al. 2015). Data suggests that a distinct amino acid position (p.66) in the *T*-
253 *gene* is relevant for a coordinated Wnt–brachyury–HOX signaling cascade, which is
254 important for cervical vertebral and spinal cord development (Galis, 1999; Yamaguchi, 1999;
255 Pennimpede et al. 2012). To the authors' knowledge, no other spontaneous mutation in
256 mammalian genes causes a homeotic transformation of cervical vertebrae similar to VSD.

257 Some familial cases of cervical ribs with a probable autosomal-dominant mode of inheritance
258 have been reported in human literature (Weston, 1956; Schapera 1987). A high incidence of
259 cervical ribs has been observed in mammals with a high rate of inbreeding (e.g 33% in
260 mammoths before extinction (Reumer et al. 2014), 11% in minipigs (KD, 1998) or 7.5% in
261 humans (Palma and Carine, 1990)). Pedigrees were not available for analysis in any of the
262 dogs included in this study; and therefore, the mode of inheritance or the degree of
263 inbreeding could not be studied. Familial cases could potentially help determine the genetic
264 basis of cervical ribs.

265 Children with neuroblastoma, brain tumours, leukaemia, soft tissue sarcoma, Wilms tumour,
266 and Ewing sarcoma also have a high incidence of cervical ribs. Pug dogs are not known to be

267 overrepresented for any of the above (Gough and Thomas, 2010). Cervical ribs have also
268 been associated with human stillbirths (Furtado et al. 2011; ten Broek et al 2012). Among the
269 100 most popular breeds in a Norwegian dog population, the Pug dog ranked eighth highest
270 for stillbirths and fourth highest for total perinatal mortality risk at the individual puppy.
271 Mean litter size was like other similar size dog breeds (Parker et al. 2017; Tonnessen et al.
272 2012). Our study was based on the United Kingdom Pug dog population and the prevalence
273 of cervical ribs in the Norway Pug dog population is currently unknown, making
274 comparisons difficult. Significant associations between congenital anomalies and cervical
275 ribs in stillborn human foetuses have been found (ten Broek et al 2012). One study did not
276 identify this association; however, an association with aneuploidy was identified, implying
277 that congenital anomalies would have been detected if they had survived to be born (Furtado
278 et al. 2011). We only evaluated the available images for the presence of other CVM's and not
279 for other major anomalies (renal agenesis, ventricular septum defects, anal atresia, etc.).

280

281 Most human adults with cervical ribs are asymptomatic, but some develop thoracic outlet
282 syndrome (TOS) (Chang et al. 2013). Thoracic outlet syndrome represents a group of distinct
283 disorders that result from the compression of one or more neurovascular elements as they
284 traverse the thoracic outlet. Unlike humans in which the brachial plexus and the subclavian
285 artery pass through the interscalene triangle, which is formed by first rib and the ventral and
286 middle components of the scalene muscles (Ferrante and Ferrante, 2017); in domestic
287 mammals, the ventral scalene muscle is not separated from the middle component and the
288 interscalene triangle is not formed (Evans and de Lahunta, 2013). The scalene muscles can
289 play an important role in the pathophysiology of this syndrome (Ferrante and Ferrante,
290 2017a), of which some forms typically occur in athletes after vigorous activity (Ferrante and
291 Ferrante, 2017b). None of the Pug dogs in our study presented with clinical signs suggestive

292 of TOS. To the authors' knowledge, TOS has not been reported in dogs, and the anatomical
293 difference between species and the fact that that Pug dogs are not too athletic may be some of
294 the reasons why; however, given the high prevalence of cervical ribs within the breed,
295 clinicians should bear in mind this differential diagnosis in Pug dogs with cervical ribs and
296 compatible clinical signs.

297 Even though they do not usually cause clinical signs in humans, cervical ribs and transitional
298 lumbo-sacral vertebra are associated with dermatomal variations and alterations in function
299 (Makhoul and Machleder, 1992; Roos, 1996; Seyfert 1997; Kim et al. 2008). Lumbo-sacral
300 transitional vertebrae can predispose dogs to cauda equina syndrome (Fluckiger et al. 2006),
301 asymmetrical hip joint development and more severe hip dysplasia (Komsta et al. 2015;
302 Fluckiger et al. 2017). The prevalence of lumbo-sacral transitional vertebrae in some dog
303 breeds has been reported as high as 22%; however, the overall prevalence is low (3.5%)
304 (Damur-Djuric et al. 2006). Despite high prevalence of lumbo-sacral transitional vertebrae
305 (12/51-24%), none of the Pugs dogs in our study presented with clinical signs suggestive of
306 hip dysplasia and/or cauda equina syndrome. Pug dogs are predisposed to hip dysplasia, with
307 an approximate prevalence of 70% (OFA 2016). To the author's knowledge, the Pug dog is
308 not predisposed to cauda equina syndrome (Meij and Bergknut 2010). If spinal stenosis
309 occurs in the absence of clinical signs, and how clinically significant hip dysplasia is in Pug
310 dogs, remains to be explored.

311 The prominent and sharp ventral process of C1, the large transverse processes of C6, the first
312 and last rib, the first lumbar transverse process and the last lumbar vertebra are all useful
313 anatomical landmarks for localisation during the surgical approach to vertebral column in
314 dogs (Sharp and Wheeler 2005). Two out of 51 (4%), 8/51 (16%) and 23/51 (45%) of Pug
315 dogs with a whole body CTs were considered to have a C5, a C6 and a C7 abnormal

316 transverse process morphology, respectively, and 52/130 (40%) were considered to have a
317 thoraco-lumbar transitional vertebra. Given the frequency of altered anatomical surgical
318 landmarks, special consideration should be taken when using them to localise during surgery
319 in pug dogs.

320 No association was found between the presence of vertebral body formation defects,
321 transitional vertebrae, bifid spinous process or an abnormal sacrum, and this may reflect a
322 different etiopathogenesis. However, an association between cervical ribs and thoracolumbar
323 transitional vertebrae was identified using CT. A possible reason why this was not significant
324 when analysing radiographs is that CT provides better bone definition compared to
325 radiography, and the presence of cervical ribs could had been underestimated. In sloths,
326 cervical disruptions occur coincident with abnormalities at the thoraco-lumbar, lumbo-sacral,
327 and sacro-caudal boundaries. In people, significant associations between cervical ribs and
328 sacralisation (Erken et al. 2002) and thoraco-lumbar transitional vertebrae have been found
329 (Bots et al. 2011). Thoraco-lumbar transitional vertebrae have also been associated with a
330 higher incidence of concomitant lumbo-sacral transitional vertebrae (Carrino et al. 2011) and
331 vice versa (Nakajima et al. 2014).

332 A total of 49 bifid spinous processes were identified in 47/130 (36%) Pug dogs. This
333 prevalence is in accordance with a previous study in which bifid spinous process was
334 identified exclusively at T1 in Pug dogs (Ryan et al. 2017). One Pug dog in our study had
335 bifid spinous process at C5 and C6 and another Pug dog had bifid spinous process at T5. It is
336 currently unclear why bifid spinous process occurs more frequently at T1 and what the
337 clinical relevance of this finding may be. A higher than normal incidence of cervical ribs has
338 been previously identified in fetuses and infants with open spina bifida (myelomeningocele)
339 (5/112-4.5%) and spina bifida (4/10-40%), suggesting a possible relation between both

340 malformations (Galis et al. 2006; Naik et al. 1978). Our study identified no association
341 between bifid spinous process and cervical ribs.

342 Twenty-eight out of 51 (55%) Pug dogs had 26 pre-sacral vertebrae instead of the fixed 27
343 pre-sacral vertebrae common to all the canine species (de Lahunta et al. 2015). Having 26
344 pre-sacral vertebrae was associated with the presence of cervical ribs or a lumbo-sacral
345 transitional vertebra. In humans, the presence of lumbo-sacral transitional vertebra is
346 associated with an anomalous total number of vertebrae, but the presence of thoraco-lumbar
347 transitional vertebra is not (Carrino et al. 2011). The mechanisms of regulation of the number
348 of segments in the vertebral column and its associations with homeotic transformations are
349 still not well understood (Gomez and Pourquie 2009).

350 Hypoplasia or aplasia of the T9-L2 caudal articular processes are thought to sometimes lead
351 to the formation of fibrous tissue and cause a constrictive myelopathy along these spinal cord
352 segments in Pug dogs (Fisher et al. 2013, Breit, 2002). The association between cervical ribs
353 and thoraco-lumbar transitional vertebra identified in our study supports the previously
354 demonstrated fact that cervical ribs appear to be accompanied by homeotic changes of several
355 adjacent cervical and thoracic vertebrae, and sometimes all thoracic vertebrae (ten Broek et al
356 2012; Oostra et al 2005). As a result, the normal vertebral anatomy, including the caudal
357 articular processes of T9-T13, could be altered. However, we found no association between
358 the presence of caudal articular process hypoplasia/aplasia, the presence of vertebral body
359 formation defects, transitional vertebra at any location or bifid spinous process. Furthermore,
360 higher C7/T1 dorsal spinous process ratios were associated with having fewer vertebrae
361 affected by caudal articular process hypoplasia/aplasia, but the correlation coefficient was
362 low. The reasoning behind this finding remains unknown.

363 Mutations in *Hox* genes are almost absent in humans, other than a few mutations in caudal
364 *Hox* genes (e.g. *HoxA11*, *Hox A13*, *HoxD10*, *HoxD13*) (Kosaki et al. 2002, Dobbs et al.
365 2006) and supposedly because most mutations are lethal at a very young stage. As supported
366 by this study, cervical ribs are rarely the only homeotic change along the vertebral column
367 and are often associated with shifts of the thoraco-lumbar and lumbo-sacral boundary
368 (Varela-Lasheras et al. 2011). This combination of homeotic transformations can be well
369 explained by mutations of genes upstream of *Hox* (Alkema et al 1995, Ikeya and Takada
370 2001, van den Akker et al 2002). Alternatively, it can be caused by mutations in several *Hox*
371 genes at the same time, which is unlikely, given that such mutations are already not found in
372 a single *Hox* gene. Therefore, it is more likely that mutations in genes upstream of *Hox* genes
373 are involved in the induction of cervical ribs in dogs and other mammals.

374

375 To the authors' knowledge, no previous CT based study in veterinary literature including a
376 large number of dogs has described CVM's of the whole vertebral column (excluding
377 coccygeal vertebrae) in any specific breed. Due to the high percentage of altered pre-sacral
378 vertebral counts and the high number and variety of CVM's identified in the Pug dog, it
379 appears to be a good naturally occurring model to further investigate the aetiology of CVM's
380 and numerical alterations.

381

382 **Conclusion**

383 Despite the high prevalence of C7 homeotic transformation, affected Pug dogs survive to
384 adulthood, and reproduce; whereas most conceived individuals with a deviation number of
385 cervical vertebrae die prenatally, or as infant (ten Broek et al 2012). We hypothesise that
386 stabilizing selection relaxed due to human care and the pleiotropic constraints broke to allow
387 the evolution of novelties in body plans. Sloths, manatees, *Bos taurus* with VDS and Pug

388 dogs are unique examples of mammals with a naturally occurring exceptional number of
389 cervical vertebrae. Further research is required to answer questions regarding the aetiology
390 and frequent natural occurrence of the different CVM's and numerical anomalies of the axial
391 skeleton in Pug dogs.

392 **Author contributions**

393 J.B.: concept/design, acquisition of data, data analysis/interpretation, drafting of the
394 manuscript, assembled figures, critical revision of the manuscript and approval of the article.
395 S.D.D.: acquisition of data, critical revision of the manuscript and approval of the article.
396 R.J.L: acquisition of data, critical revision of the manuscript and approval of the article. J.P:
397 acquisition of data, critical revision of the manuscript and approval of the article. E.M: data
398 analysis/interpretation, critical revision of the manuscript and approval of the article. C.S:
399 critical revision of the manuscript and approval of the article. S.B.: acquisition of data,
400 critical revision of the manuscript and approval of the article. J.J.S.: critical revision of the
401 manuscript and approval of the article. C.R.: acquisition of data, critical revision of the
402 manuscript and approval of the article. N.F.: critical revision of the manuscript and approval
403 of the article. R.G: concept/design, data analysis/interpretation, drafting of the manuscript,
404 critical revision of the manuscript and approval of the article.

405

406 **Conflict of interest statement**

407 None of the authors has any financial or personal relationships that could inappropriately
408 influence or bias the content of the paper.

409

410 An abstract with some of the results of this study was submitted to the European College of

412 **References:**

- 413 Alkema MJ, van der Lugt NM, Bobeldijk RC, Berns A, van Lohuizen M (1995)
414 Transformation of axial skeleton due to overexpression of bmi-1 in transgenic mice. *Nature*
415 374, 724-727.
- 416 Badlangana NL, Adams JW, Manger PR (2009) The giraffe (*Giraffa camelopardalis*) cervical
417 vertebral column- a heuristic example in understanding evolutionary processes? *ZOOLOGICAL JOURNAL OF THE LINNEAN SOCIETY* 155, 736-757.
418
419
- 420 Bots J, Wijnaendts LC, Delen S, Van Dongen S, Heikinheimo K, Galis F (2011) Analysis of
421 cervical ribs in a series of human fetuses. *J Anat* 219, 403-409.
422
- 423 Breit S, Kunzel W (1998) Osteologische Besonderheiten an Wirbelsäulen von Rassehunden:
424 eine röntgenologische und morphologische Studie. *Wiener Tierärztliche Monatsschrift* 85,
425 340-350.
426
- 427 Breit S (2002) Osteological and Morphometric Observations on Intervertebral Joints in the
428 Canine Pre-diaphragmatic Thoracic Spine (Th1–Th9). *Vet J* 164, 216-223.
429
- 430 Buchholtz EA, Stepien CC (2009) Anatomical transformation in mammals: developmental
431 origin of aberrant cervical anatomy in tree sloths. *Evol Dev* 11, 69-79.
432
- 433 Buchholtz EA, Booth AC, and Webbink KE (2007) Vertebral anatomy in the Florida
434 manatee, *Trichechus manatus latirostris*: a developmental and evolutionary analysis. *Anat Rec (Hoboken)* 290, 624-637.
435
- 436 Carrino JA, Campbell PD, Lin DC et al. (2011) Effect of Spinal Segment Variants on
437 Numbering Vertebral Levels at Lumbar MR Imaging. *Radiology* 259, 196-202.
438
439
- 440 Chang KZ, Likes K, Davis K, Demos J, Freischlag JZ (2013) The significance of cervical
441 ribs in thoracic outlet syndrome. *J Vasc Surg* 57, 771-5.
442
- 443 Chernoff N, Rogers JM (2004) Supernumerary ribs in developmental toxicity bioassays and
444 in human populations: incidence and biological significance. *J Toxicol Environ Health B Crit Rev* 7, 437-439.
445
446
- 447 Damur-Djuric N, Steffen F, Hassig M, Morgan JP, Fluckiger MA (2006) Lumbosacral
448 Transitional Vertebrae in Dogs: Classification, Prevalence, and Association with Sacroiliac
449 Morphology. *Vet Radiol Ultrasound* 47, 32-38.
450
- 451 de Lahunta, A., Glass, E., Kent, M., (2015). *Veterinary neuroanatomy and clinical neurology*
452 4th ed. Missouri: Elsevier, pp. 86.
453
- 454 Dobbs MB, Gurnett CA, Pierce B (2006) HOXD10 M319K mutation in a family with
455 isolated congenital vertical talus. *J Orthop Res* 24, 448–453.
456
- 457 Erken E, Ozer HT, Gulek B, Durgun B (2002) The association between cervical rib and
458 sacralization. *Spine* 27, 1659-1664.
459

460 Evans, de Lahunta A (2013). The skeleton. In: HE. Evans, A. de Lahunta, ed., Miller's
461 Anatomy of the Dog, 4th ed. Missouri: Elsevier, pp. 81-156.

462

463 Ferrante MA, Ferrante ND (2017a) The thoracic outlet syndromes: Part 1. Overview of the
464 thoracic outlet syndromes and review of true neurogenic thoracic outlet syndrome. *Muscle*
465 *Nerve* 55, 782-793.

466 Ferrante MA, Ferrante ND (2017b) The thoracic outlet syndromes: part 2. The arterial,
467 venous, neurovascular, and disputed thoracic outlet syndromes. *Muscle Nerve* 54, 663-673.

468

469 fci.be, (2017). FCI breeds nomenclature. [online] Available at:
470 <http://www.fci.be/en/Nomenclature/>

471

472 Fisher SC, Shores A, Simpson ST (2013) Constrictive myelopathy secondary to hypoplasia
473 or aplasia of the thoracolumbar caudal articular processes in Pugs - 11 cases. *J Am Vet Med*
474 *Assoc* 242, 223-229.

475 Fluckiger MA, Damur-Djuric N, Hassig M, Morgan JP, Steffen F (2006) A Lumbosacral
476 Transitional Vertebra in the Dog Predisposes to Cauda Equina Syndrome. *Vet Radiol*
477 *Ultrasound* 47, 39-44.

478

479 Fluckiger MA, Steffen F, Hassig M, Morgan JP (2017) Asymmetrical lumbosacral
480 transitional vertebrae in dogs may promote asymmetrical hip joint development. *Vet Comp*
481 *Orthop Traumatol* 30, 137-142.

482

483 Furtado LV, Thaker HM, Erickson LK, Shirts BH, Opitz JM (2011) Cervical ribs are more
484 prevalent in stillborn fetuses than in live-born infants and are strongly associated with fetal
485 aneuploidy. *Pediatr Dev Pathol* 14, 431-437.

486

487 Galis F (1999) Why do almost all mammals have seven cervical vertebrae? Developmental
488 constraints, Hox genes, and cancer. *J Exp Zool* 285, 19-26.

489

490 Galis F, van Alphen JJM, Metz JAJ (2001) Why five fingers- Evolutionary constraints on
491 digit numbers. *Trends Ecol Evol* 16, 637-646.

492

493 Galis F, Van Dooren TJM, Feuth JD, Metz JAJ, Witkam A, Ruinard S et al. (2006) Extreme
494 selection in humans against homeotic transformations of cervical vertebrae. *Evolution* 60,
495 2643-2654.

496 Galis F, Arntzen JW, Lande R (2010) Dollo's law and the irreversibility of digit loss in
497 *Bachia*. *Evolution* 64, 2466-2485.

498

499 Gladstone, R. J., and C. P. G. Wakeley. (1932) Cervical ribs and rudimentary first thoracic
500 ribs considered from the clinical and etiological standpoints. *J. Anat.* 66,334-337.

501

502 Gough A, Thomas A (2010) Breed predisposition to disease in dogs and cats 2nd ed. Oxford:
503 Willey-Blackwell, pp. 579-585.

504

505 Gomez C and Pourquie O (2009) Developmental control of segment numbers in vertebrates.
506 *J Exp Zool B Mol Dev Evol* 312, 533-44.

507 Griesinger CB, Richards CD, Ashmore JF (2005). Retinoic acid coordinates somitogenesis
508 and left-right patterning in vertebrate embryos. *Nature* 435, 212-215.
509

510 Gutierrez-Quintana R, Guevar J, Stalin C, Faller K, Yeaman C, Penderis J (2014) A
511 proposed radiographic classification scheme for congenital thoracic vertebral malformations
512 in brachycephalic "screw-tailed" dog breeds. *Vet Radiol Ultrasound* 55, 585-591.
513

514 Henderson MS (1913) Cervical rib: report of thirty-one cases. *Am J Orthop Surg* 11, 408-
515 430.
516

517 Hostikka SL, Gong J, Carpenter EM (2009) Axial and appendicular skeletal transformations,
518 ligament alterations, and motor neuron loss in Hoxc10 mutants. *Int J. Biol Sci* 5, 397-410.

519 Ikeya M, Takada S (2001) Wnt-3a is required for somite specification along the
520 anterior-posterior axis of the mouse embryo and for regulation of cdx-1 expression. *Mech Dev*
521 103, 27-33.

522 KD J (1998) Minipig in reproduction toxicology. *Scandinavian Journal of laboratory*
523 *Animal Science* 25, 63-75.
524

525 Komsta R, Łojczyk-Szczepaniak A, Dębiak P (2015) Lumbosacral Transitional Vertebrae,
526 Canine Hip Dysplasia, and Sacroiliac Joint Degenerative Changes on Ventrodorsal
527 Radiographs of the Pelvis in Police Working German Shepherd Dogs. *Top Companion Anim*
528 *Med* 30, 10-15.
529

530 Kosaki K, Kosaki R, Suzuki T et al. (2002) Complete mutation analysis panel of the 39
531 human HOX genes. *Tetralogy* 65, 50-62.
532

533 Kim YH, Lee PB, Lee CJ, Lee SC, Kim YC, Huh J (2008) Dermatome variation of
534 lumbosacral nerve roots in patients with transitional lumbosacral vertebrae. *Anesth Analg*
535 106, 1279-1283.
536

537 Kromik A, Ulrich R, Kusenda M et al. (2015) The mammalian cervical vertebrae blueprint
538 depends on the T (brachyury) gene. *Genetics* 199, 873-83.
539

540 Krumlauf R (1994) Hox genes in vertebrate development. *Cell* 78, 191-201.

541 Leboucq HLF (1895) Recherches sur les variations anatomiques de la première côte chez
542 l'homme, Hayez. *Archiv Biol* 15, 9-178.

543 Li ZL, Shiota K (1999) Stage-specific homeotic vertebral transformations in mouse fetuses
544 induced by maternal hyperthermia during somitogenesis. *Dev Dyn* 216, 336-348.
545

546 Makhoul RG, Machleder HI (1992) Developmental anomalies at the thoracic outlet: An
547 analysis of 200 consecutive cases. *J Vasc Surg* 16, 534-545.
548

549 Meij BP, Bergknot N (2010) Degenerative lumbosacral stenosis in dogs. *Vet Clin North Am*
550 *Small Anim Pract* 40, 983-1009.
551
552 Morgan JP (1968) Congenital Anomalies of the Vertebral Column of the Dog- A Study of the
553 Incidence and Significance Based on a Radiographic and Morphologic Study. *Vet Radiol*
554 *Ultrasound* 21-29.
555
556 Naik DR, Lendon RG, Barson AJ (1978) A radiological study of vertebral and rib
557 malformations in children with myelomeningocele. *Clin Radiol* 29, 427-430.
558
559 Nakajima A, Usui A, Hosokai Y, Kawasumi Y, Abiko K, Funayama M, Saito H (2014) The
560 prevalence of morphological changes in the thoracolumbar spine on whole-spine computed
561 tomographic images. *Insights Imaging* 5, 77-83.
562
563 Ofa.org, (2016). Hip dysplasia statistics. [online] Available at:
564 http://www.ofa.org/stats_hip.html
565
566 Oostra RJ, Hennekam RC, De Rooij L, Moorman AF (2005) Malformations of the axial
567 skeleton in Museum Vrolijk I: homeotic transformations and numerical anomalies. *Am J Med*
568 *Genet A* 134, 268-281.
569
570 Palma A, Carini F (1990) Variazioni dell'apofisi trasversa della settima vertebra cervicale:
571 studio anatomo-radiologico su una popolazione "segregate". *Arch Ital Anat Embriol* 95, 11-
572 16
573
574 Pennimpede T, Proske J, Konig A, VidigaL JA, Morkel M, Bramsen JB et al. (2012) In vivo
575 knockdown of Brachyury results in skeletal defects and urorectal malformations resembling
576 caudal regression syndrome. *Dev Biol* 372, 55-67.
577
578 Reumer JW, Ten Broek CM, Galis F (2014) Extraordinary incidence of cervical ribs indicates
579 vulnerable condition in Late Pleistocene mammoths. *PeerJ* Mar 25;2:e318.
580
581 Roos DB (1996) Historical perspectives and anatomic considerations. Thoracic outlet
582 syndrome. *Semin Thoracic Cardiovasc Surg* 8, 183-189.
583
584 Ryan R, Gutierrez-Quintana R, Ter Haar G, De Decker S (2017) Prevalence of thoracic
585 vertebral malformations in French bulldogs, Pugs and English bulldogs with and without
586 associated neurological deficits. *Vet J* 221, 25-29.
587
588 Seyfert S (1997) Dermatome variations in patients with transitional vertebrae. *J Neurol*
589 *Neurosurg Psychiatry* 63, 801-803.
590
591 Schapera J (1987) Autosomal dominant inheritance of cervical ribs. *Clin Genet* 31, 386-388.
592
593 Schulz AH (1961) Vertebral column and thorax. In: Hofer HSA, Starck D, ed., *Primatologia,*
594 *handbook of primatology.* 1st ed. Basel/New York: S. Karger, pp. 51-56.
595
596 Schumacher R, Mai A, Gutjahr P (1992) Association of rib anomalies and malignancy in
597 childhood. *Eur J Pediatr* 151, 432-434.
598

599 Sharp NJH, Wheeler SJ (2005) Cervical disc disease. In: NJH. Sharp, SJ. Wheeler, ed., Small
600 Animal Spinal Disorders Diagnosis and Surgery, 2nd ed. Elsevier, pp. 93-120.
601

602 Spadlinski L, Cecot T, Majos A et al. (2016) The Epidemiological, Morphological, and
603 Clinical Aspects of the Cervical Ribs in Humans. *Biomed Res Int* 2016, 8034613.

604 Starck D (1978) Vergleichende anatomie der wirbeltiere. Berlin: Springer Verlag.

605 Steigenga MJ, Helmerhorst FM, De Koning J, Tijssen Ans MI, Ruinard SAT, Galis F
606 (2006) Evolutionary conserved structures as indicators of medical risks- increased incidence
607 of cervical ribs after ovarian hyperstimulation in mice. *Animal Biology* 56, 63-68.

608 Steffen F, Berger M, Morgan JP (2004) Asymmetrical, Transitional, Lumbosacral Vertebral
609 Segments in Six Dogs- A Characteristic Spinal Syndrome. *J Am Anim Hosp Assoc* 40, 338-
610 344.
611

612 ten Broek, CMA, Bakker AJ, Varela-Lasheras I, Bugiani M, Van Dongen S, Galis F (2012)
613 Evo-Devo of the Human Vertebral Column: On Homeotic Transformations, Pathologies and
614 Prenatal Selection. *Evol Biol* 39, 456-471.

615 thekennelclub.org.uk, (2016). Top twenty breeds in registration order for the years 2015 and
616 2016 [online] Available at:
617 https://www.thekennelclub.org.uk/media/1098176/top_20_breeds_2015_-_2016.pdf

618 Tonnessen R, Borge KS, Nodtvedt A, Indrebo A (2012) Canine perinatal mortality: a cohort
619 study of 224 breeds. *Theriogenology* 77, 1788-801.
620

621 van den Akker E, Forlani S, Chawengsaksophak K, et al. (2002): Cdx1 and Cdx2 have
622 overlapping functions in anterioposterior patterning and posterior axis elongation.
623 *Development* 129, 2181-2193.

624 Varela-Lasheras I, Bakker AJ, van der Mije SD, Metz JAJ, van Alphen J and Galis F (2011)
625 Breaking evolutionary and pleiotropic constraints in mammals- On sloths, manatees and
626 homeotic mutations. *EvoDevo* 2, 11.

627 Vermot J, Llamas JG, Fraulob V, Niederreither K, Chambon P, Dolle P (2005)
628 Retinoic acid controls the bilateral symmetry of somite formation in the mouse embryo.
629 *Science* 308, 563-566.

630 Vila C, Maldonado JE, Wayne RK (1999) Phylogenetic relationships, evolution, and genetic
631 diversity of the domestic dog. *J Hered* 90, 71-77.
632

633 Wellik DM (2007) Hox patterning of the vertebrate axial skeleton. *Dev Dyn* 236, 2454-63.
634

635 Wery N, Narotsky MG, Pacico N, Kavlock RJ, Picard JJ, Gofflot F (2003) Defects in cervical
636 vertebrae in boric acid-exposed rat embryos are associated with anterior shifts of hox gene
637 expression domains. *Birth Defects Res A Clin Mol Teratol* 67, 59-67.
638

639 Weston WJ (1956) Genetically Determined Cervical Ribs—A Family Study. *Br J Radiol* 29,
640 455-456.

641

642 Wrzosek M, Plonek M, Zeira O, Biezynski J, Kinda W, Guzinski M (2014) Congenital
643 bipartite atlas with hypodactyly in a dog: clinical, radiographic and CT findings. *J Small*
644 *Anim Pract* 55, 375-378.

645 Yamaguchi TP, Takada S, Yoshikawa Y, Wu N, McMahon AP (1999) T (Brachyury) is a
646 direct target of Wnt3a during paraxial mesoderm specification. *Genes Dev* 15, 3185-3190.

647

648

649

650

651

652

653

654

655

656

657

658

659

660

661

662

663

664

665

666 **Table 1.** Number (N) and percentages (%) of Pug dogs with different congenital vertebral
667 malformations and number of pre-sacral vertebrae identified on whole body computed
668 tomography studies (WBCT), cervico-thoracic computed studies (C-TCT) and radiographs.

	WBCT % (affected/total)	C-TCT % (affected/total)	Radiographs % (affected/total)	Total N and %
Homeotic free	22 (11/51)	33 (15/46)	45 (15/33)	32 (41/130)
C7 cervical ribs	45 (23/51)	48 (22/46)	36 (12/33)	44 (57/130)
Transitional TL	53 (27/51)	33 (15/46)	30 (10/33)	40 (52/130)
Transitional LS	24 (12/51)	-	-	24 (12/51)
Bifid spinous process	27 (14/51)	46 (21/46)	39 (13/33)	37 (48/130)
Hemivertebra	16 (8/51)	21 (9/46)	100 (33/33)	38 (50/130)
Articular process hypoplasia/aplasia	92 (47/51)	-	-	92 (47/51)
Absent T13 vertebra	22 (11/51)	24 (11/46)	27 (9/33)	24 (31/130)
Absent L7 vertebra	33 (17/51)	-	-	33 (17/51)
26 pre-sacral vertebrae	55 (28/51)	-	-	55 (28/51)
27 pre-sacral vertebrae	45 (23/51)	-	-	45 (23/51)


669
670
671
672
673
674
675
676

677 **Table 2.** Distribution of C7 vertebra cervical ribs in Pug dogs according to gender and colour
 678 coat.

	Bilateral	Unilateral right	Unilateral left	% (affected/total number)
C7 cervical ribs	32	3	10	46 (45/97)
Gender:				
• Female	22	2	6	65 (30/46)
• Male	10	1	4	29 (15/51)
Colour coat:				
• Fawn	10	0	3	50 (13/26)
• Black	8	1	1	40 (10/25)

679
 680
 681
 682
 683
 684
 685
 686
 687
 688
 689
 690
 691
 692
 693

694 **Figure 1.** Lateral (A), ventral (B) and dorsal (C) 3D-computed tomography reconstructed
695 images of the axial skeleton (excluding some caudal vertebrae), ribs, sternum and pelvis of a
696 Pug dog with no congenital vertebral malformations.



697

698


699

700 **Figure 2.** Bar chart illustrating the number of computed tomography studies reviewed and
701 the number and percentage of dogs with C7 homeotic transformation from the breeds
702 included in the study.

703

704


705 **Figure 3.** Sagittal reconstructed computed tomography image of the thoracic vertebral
706 column of a Pug dog with a vertebral body formation defect (arrow) and ventral spondylosis.



707

708

709 **Figure 4.** Transverse computed tomography image (A) and 3D-computed tomography
710 reconstruction image (caudo-cranial oblique view-B) of the T1 vertebra of a Pug dog with a
711 bifid spinous process.




712

713

714

715 **Figure 5.** Transverse computed tomography image of T13 (A) and 3D-computed tomography
716 reconstruction image (dorsal view-B) of T11, T12 and T13 vertebrae of a Pug dog. T13
717 vertebra is a transitional thoraco-lumbar vertebra and has two long thick transverse processes
718 (no joints).





719

720

721


722 **Figure 6.** Transverse computed tomography image of L7 (A) and 3D-computed tomography
723 reconstruction image (ventral view-B) of L6, L7 and the sacrum of a Pug dog. L7 is a lumbo-
724 sacral transitional vertebra and has asymmetric transverse processes.



725

726

727 **Figure 7.** Two 3D-computed tomography reconstructed images (dorsal views) of T11, T12,
728 T13, L1 and L2 vertebrae of a Pug dog. Image A illustrates normal caudal articular processes.
729 Image B has bilateral hypoplastic/aplastic caudal articular processes at T11-T12, T12-T13
730 (small arrow), unilateral hypoplastic/aplastic caudal articular processes at T13-L1 (medium
731 sized arrow), and normal caudal articular processes at L1-L2 (large arrow).




732


733

734

735 **Figure 8.** Transverse computed tomography images (left) and 3D-computed tomography
736 reconstructions (right) of the C7 vertebra of multiple Pug dogs: (A) Normal vertebra (B)
737 Vertebra with bilateral short ribs and no joint (C) Vertebra with a left long rib and no joint,
738 and a right long rib with a joint not visible in the transverse image (D) Vertebra with bilateral
739 short ribs and joints (E) Vertebrae with bilateral long ribs and joints.



741 **Figure 9.** Spearman correlation for the added length (cm) of left and right cervical ribs and
742 the dorsal spinous process C7/T1 ratio, $r = 0.55$.



743