

This author's accepted manuscript may be used for non-commercial purposes in accordance with [Wiley Terms and Conditions for Self-Archiving](#).

The full details of the published version of the article are as follows:

TITLE: Inhibition of arterial medial calcification and bone mineralization by extracellular nucleotides: The same functional effect mediated by different cellular mechanisms

AUTHORS: **Patel, J J**; Zhu, D; Opdebeeck, B; D'Haese, P; Millán, J L; Bourne, L E; **Wheeler-Jones, C P D**; Arnett, T R; MacRae, V E; **Orriss, I R**

JOURNAL: Journal of Cellular Physiology

PUBLISHER: Wiley

PUBLICATION DATE: 4 October 2017 (online)

DOI: <http://dx.doi.org/10.1002/jcp.26166>

# **Inhibition of arterial medial calcification and bone mineralisation by extracellular nucleotides: the same functional effect mediated by differing cellular mechanisms**

**Jessal J Patel<sup>1</sup>, Dongxing Zhu<sup>2</sup>, Britt Opdebeeck<sup>3</sup>, Patrick D'Haese<sup>3</sup>, José Luis Millán<sup>4</sup>, Lucie Bourne<sup>1</sup>, Caroline PD Wheeler-Jones<sup>1</sup>, Timothy R Arnett<sup>5</sup>, Vicky E MacRae<sup>2</sup>, Isabel R Orriss<sup>1</sup>**

<sup>1</sup> Department of Comparative Biomedical Sciences, Royal Veterinary College, London, UK

<sup>2</sup> The Roslin Institute and Royal (Dick) School of Veterinary Studies, University of Edinburgh, Edinburgh, UK

<sup>3</sup> Laboratory of Pathophysiology, Department of Biomedical Sciences, University of Antwerp, Belgium

<sup>4</sup> Sanford Burnham Prebys Medical Discovery Institute, La Jolla, USA

<sup>5</sup> Department of Cell and Developmental Biology, University College London, London, UK

Address correspondence to: Isabel R Orriss  
Department of Comparative Biomedical Science  
Royal Veterinary College  
London, NW1 0TU  
Tel: 020 7468 1238 (ext 5468)  
Email: [iorriss@rvc.ac.uk](mailto:iorriss@rvc.ac.uk)

**Short title:** Arterial medial calcification and ATP

**Key words:** arterial medial calcification, bone formation, ATP, pyrophosphate, NPP1, osteoblasts, vascular smooth muscle cells

**Conflict of Interest:** The authors have no conflict of interest

## ABSTRACT

Arterial medial calcification (AMC) is thought to share some outward similarities to skeletal mineralisation and has been associated with the transdifferentiation of vascular smooth muscle cells (VSMCs) to an osteoblast-like phenotype. ATP and UTP have previously been shown to inhibit bone mineralisation. This investigation compared the effects of extracellular nucleotides on calcification in VSMCs with those seen in osteoblasts. ATP, UTP and the ubiquitous mineralisation inhibitor, pyrophosphate (PP<sub>i</sub>), dose dependently inhibited VSMC calcification by  $\leq 85\%$ . Culture of VSMCs in calcifying conditions was associated with an increase in apoptosis; treatment with ATP, UTP and PP<sub>i</sub> reduced apoptosis to levels seen in non-calcifying cells. Extracellular nucleotides had no effect on osteoblast viability. Basal alkaline phosphatase (TNAP) activity was over 100-fold higher in osteoblasts than VSMCs. ATP and UTP reduced osteoblast TNAP activity ( $\leq 50\%$ ) but stimulated VSMC TNAP activity ( $\leq 88\%$ ). The effects of extracellular nucleotides on VSMC calcification, cell viability and TNAP activity were unchanged by deletion or inhibition of the P2Y<sub>2</sub> receptor. Conversely, the actions of ATP/UTP on bone mineralisation and TNAP activity were attenuated in osteoblasts lacking the P2Y<sub>2</sub> receptor. Ecto-nucleotide pyrophosphatase/phosphodiesterase 1 (NPP1) hydrolyses ATP and UTP to produce PP<sub>i</sub>. In both VSMCs and osteoblasts, deletion of NPP1 blunted the inhibitory effects of extracellular nucleotides suggesting involvement of P2 receptor independent pathways. Our results show that although the overall functional effect of extracellular nucleotides on AMC and bone mineralisation is similar there are clear differences in the cellular mechanisms mediating these actions.

## INTRODUCTION

Vascular calcification is a common consequence of ageing, atherosclerosis, diabetes and chronic kidney disease. It is the pathological deposition of calcium phosphate mineral, usually as hydroxyapatite, in the medial and/or intimal layer of the arteries and heart valves. Arterial medial calcification (AMC) refers to the calcification that occurs within the tunica media of blood vessels and is characterised by increased vessel stiffness and reduced blood flow (Young et al., 1993). Traditionally, AMC was thought to be a passive process caused by high serum levels of phosphate and calcium. However, it is now accepted that it is a complex cell-mediated process that shares some similarities with the process of physiological bone formation. Specifically, AMC is thought to involve the loss of calcification inhibitors (e.g. pyrophosphate (PP<sub>i</sub>), osteopontin, fetuin A), gain of calcification inducers (e.g. alkaline phosphatase (TNAP)) and increased apoptosis (Proudfoot et al., 2000; Zhu et al., 2012). Whilst many cell types can contribute towards the development of AMC, vascular smooth muscle cells (VSMCs) are thought to be the major cell type involved (Narisawa et al., 2007; Zhu et al., 2011). Within a calcifying environment (high phosphate), VSMCs can undergo a phenotypic transdifferentiation to take on characteristics usually associated with bone-forming osteoblasts (Shroff and Shanahan, 2007; Zhu et al., 2011).

ATP has long been recognized for its role in intracellular energy metabolism; however, it is also an important extracellular signalling molecule. ATP and related compounds (UTP, ADP, UDP) act via purinergic P2 receptors to regulate cell proliferation, differentiation, survival and function in many tissues (Burnstock, 2007). The P2 receptor family is made up of seven P2X ion channels (P2X1-7) and eight P2Y G-protein coupled receptors (P2Y<sub>1,2,4,6,11-14</sub>) (Abbracchio and Burnstock, 1994; Burnstock and Kennedy, 1985).

Purinergic signalling plays a number of roles in the cardiovascular system (see review (Burnstock and Ralevic, 2014)). VSMCs express multiple P2 receptor subtypes (Wang et al., 2002) and in recent years several investigations have examined the role of purinergic signalling in the different forms of vascular calcification (Fish et al., 2013). However, they provide conflicting evidence as to whether extracellular nucleotides are harmful or protective. P2Y<sub>2</sub> receptor mediated signalling has been shown to promote the survival of aortic valve interstitial cells and protect against aortic valve calcification (Cote et al., 2012) and arterial intimal calcification (Qian et al., 2017). Additionally, a biochemical study showed that ATP inhibited calcium phosphate deposition in rat VSMCs (Villa-Bellosta and Sorribas, 2013). In contrast, activation of P2YRs by Up<sub>4</sub>A (a non-selective P2R agonist) has been reported to enhance VSMC calcification (Schuchardt et al., 2012).

Once released extracellular nucleotides are rapidly broken down by ecto-nucleotidases to limit their actions to cells within close proximity of the release site. PP<sub>i</sub>, a ubiquitous and potent inhibitor of calcification (Fleisch and Bisaz, 1962), is generated when nucleotide triphosphates are hydrolysed by ecto-nucleotide pyrophosphatase/phosphodiesterases (NPPs). VSMCs express many ecto-nucleotidases, including NPP1, and can release ATP in a controlled manner (Prosdocimo et al., 2009; Prosdocimo et al., 2010; Villa-Bellosta et al., 2011). Accumulating evidence now suggests that the hydrolysis of locally released ATP by NPP1 is the major source of extracellular PP<sub>i</sub> (Orriss et al., 2016; Prosdocimo et al., 2009). This PP<sub>i</sub> can then act locally to regulate the level of calcification. Consistent with this, NPP1 knockout mice (*Enpp1*<sup>-/-</sup>) display extensive ectopic calcification in a variety of soft tissues including the aorta, kidney, cartilage, ear pinna and whisker vibrissae (Hajjawi et al., 2014; Johnson et al., 2003; Mackenzie et al., 2012). Furthermore, mutations in the gene encoding NPP1 lead to the recessive condition Generalised Arterial Calcification of Infancy (GACI) which is characterised by extensive vascular calcification (Rutsch et al., 2001).

The regulation of bone formation by extracellular nucleotides has been widely studied often with conflicting results (see reviews (Burnstock et al., 2013; Gartland et al., 2012; Noronha-Matos and Correia-de-Sa, 2016; Orriss, 2015)). Activation of several P2 receptors (e.g. P2Y<sub>2</sub>, P2X1, P2X7) has been shown to both inhibit and promote bone mineralisation (Hoebertz et al., 2002; Noronha-Matos et

al., 2014; Orriss et al., 2012a; Orriss et al., 2007; Panupinthu et al., 2007; Xing et al., 2014). We have previously reported that ATP and UTP selectively inhibit the mineralisation of the organic matrix and TNAP expression and activity (Orriss et al., 2007). These actions are due to both P2 receptor mediated signalling and also direct hydrolysis by NPP1 to produce PP<sub>i</sub> (Orriss et al., 2012a; Orriss et al., 2013; Orriss et al., 2007). Recently, we also showed that activation of the P2Y<sub>2</sub> receptor exerts some of its effects on bone mineralisation indirectly by promoting the release of ATP from osteoblasts (Orriss et al., 2017).

The aim of this study was to investigate the effects of extracellular nucleotides on AMC and, given the apparent similarities to bone mineralisation, to compare any functional effects to those seen in osteoblasts. Established *in vitro* mouse models of AMC and bone formation were used to determine the cellular mechanisms mediating these actions. Since the inhibitory effects of ATP and UTP on bone mineralisation involve both P2 receptor dependent and independent signalling we examined both pathways in parallel.

## METHODS

### Reagents

All tissue culture reagents were purchased from Life Technologies (Paisley, UK); unless mentioned, all chemicals were purchased from Sigma Aldrich (Poole, UK). The selective P2Y<sub>2</sub> receptor antagonist, ARC118935XX, was obtained from Tocris Bioscience (Bristol, UK).

### Animals

Mice lacking the P2Y<sub>2</sub> receptor gene (*P2Y<sub>2</sub>R*<sup>-/-</sup>) were obtained from Jackson Laboratories (Bar Harbor, Maine, USA). The generation and characterisation of *P2Y<sub>2</sub>R*<sup>-/-</sup> mice, which are on a C57BL/6J background, has been previously described (Homolya et al., 1999). Animals were bred from homozygote (*P2Y<sub>2</sub>R*<sup>-/-</sup>) and parental strain wildtype (*P2Y<sub>2</sub>R*<sup>+/+</sup>) breeding pairs. The generation and characterisation of mice lacking NPP1 (*Enpp1*<sup>-/-</sup>), which are on a 129Sv/TerJ genetic background, has previously been described (Sali et al., 1999). Animals were bred from heterozygote (*Enpp1*<sup>+/-</sup>) breeding pairs. All mice were housed under standard conditions with free access to food and water. All procedures complied with the UK animals (Scientific Procedures) Act 1986 and were reviewed and approved by the Royal Veterinary College Research Ethics Committee.

### Vascular smooth muscle cell (VSMC) calcification assay

Primary VSMCs were isolated from aortas of *P2Y<sub>2</sub>R*<sup>-/-</sup> or *Enpp1*<sup>-/-</sup> mice and their corresponding wildtypes (*P2Y<sub>2</sub>R*<sup>+/+</sup> or *Enpp1*<sup>+/+</sup>). After removal of the adventitia, the aorta was opened to expose the endothelial layer under a dissection microscope. Tissues from 6-8 animals were pooled and incubated with trypsin (0.25% w/v) for 10 minutes to remove any remaining adventitia and endothelium. Tissues were incubated overnight in alpha Minimum Essential Medium, supplemented with 10% foetal calf serum (FCS), 100U/ml penicillin, 100µg/ml streptomycin and 0.25µg/ml amphotericin (complete mixture abbreviated to  $\alpha$ MEM) before being digested with 425U/ml collagenase type II (Worthington Biomedical Corporation, Lakewood, USA) for 5 hours. Isolated VSMCs were expanded in T25 tissue culture flasks in a humidified atmosphere of 5% CO<sub>2</sub>-95% air at 37°C until confluent. Following seeding into 24-well plates at a density of 2.5 x 10<sup>4</sup> cells/well, VSMCs were cultured in control ( $\alpha$ MEM only) or calcifying medium ( $\alpha$ MEM + 2mM phosphate) for up to 14 days, with half medium changes every 3 days. Cells were treated with 1-100µM extracellular nucleotides (ATP, ADP, UTP, UDP) or pyrophosphate (PP<sub>i</sub>) for the duration of the culture. The selective P2Y<sub>2</sub> receptor antagonist, ARC118935XX, was used to confirm results obtained with *P2Y<sub>2</sub>R*<sup>-/-</sup> cells.

### Aortic ring calcification assay

Aortas were isolated from *P2Y<sub>2</sub>R*<sup>+/+</sup> or *P2Y<sub>2</sub>R*<sup>-/-</sup> mice and the adventitia layer removed. The vessels were cut into 2-3mm rings and cultured overnight in serum free  $\alpha$ MEM. After 24 hours, the rings were transferred to calcification medium ( $\alpha$ MEM plus 2.5mM phosphate and 2.7mM calcium chloride). Aortic rings were cultured for a further 9 days with half medium changes every 3 days.

### Determination of VSMC and aortic ring calcification

Calcifying VSMCs or aortic rings were washed twice with phosphate buffered saline (PBS) and incubated with 0.6M HCl at room temperature for 24 hours. Calcium content was measured colorimetrically by stable interaction with o-cresolphthalein using a commercially available kit (Sigma-Adrich, Poole, UK) and corrected for total protein concentration using the Bradford assay. Calcium deposition was visualised by alizarin red staining of VSMC cell layers as previously described (Taylor et al., 2014).

### Osteoblast bone formation assay

Osteoblasts were isolated from the calvariae of 3-5 day old mice by trypsin/collagenase digestion as previously described (Orriss et al., 2014; Orriss et al., 2012b; Taylor et al., 2014). Cells were obtained from *P2Y<sub>2</sub>R*<sup>-/-</sup>, *P2Y<sub>2</sub>R*<sup>+/+</sup>, *Enpp1*<sup>-/-</sup> or *Enpp1*<sup>+/+</sup> animals. Following isolation, cells were resuspended in

$\alpha$ MEM and cultured for 2-4 days in a humidified atmosphere of 5% CO<sub>2</sub>-95% air at 37°C in 75 cm<sup>2</sup> flasks until fully confluent. Cells were sub-cultured into 6-well trays in  $\alpha$ MEM supplemented with 2mM  $\beta$ -glycerophosphate and 50 $\mu$ g/ml ascorbic acid, with half medium changes every 3 days. Cells were treated with 1-100 $\mu$ M ATP, UTP or PP<sub>i</sub> for the duration of the culture.

To assess bone formation, experiments were terminated by fixing the cells in 2.5% glutaraldehyde for 5 minutes. Cell culture plates were imaged at 800 dpi using a flat-bed scanner (Epson Perfection 4990 Photo) and the total area of bone nodules formed was quantified by image analysis, as described previously (Orriss et al., 2012b). Cell layers were stained with alizarin red for microscopy.

#### ***Analysis of P2 receptor gene expression by real time PCR***

VSMCs were cultured in control or calcification medium for 14 days. RNA was extracted using RNeasy total RNA (Qiagen Ltd, Crawley, UK), according to the manufacturer's instructions. RNA was quantified and reverse transcribed as previously described (Mackenzie et al., 2014). Levels of mRNA expression were measured using the SYBR green detection method (Roche, East Sussex, UK) as previously reported (Staines et al., 2014). Data are presented as the fold change in expression in calcifying VSMCs relative to control cells. Primer sequences are shown in table 1.

#### ***Cell viability assay***

VSMCs and osteoblasts were cultured for 14 days in medium supplemented with ATP, UTP or PP<sub>i</sub> (1-100 $\mu$ M). Cell number and viability was determined using the CytoTox 96® colorimetric cytotoxicity assay (Promega UK, Southampton UK), as described previously (Orriss et al., 2012a). Cell supernatants were collected to determine medium LDH levels (cell viability). To establish total cellular LDH levels, cells were lysed with 1% Triton X-100 in water (lysis buffer, 15 $\mu$ l/ml of medium) for 1 hour. The LDH content of the supernatants and cell lysates were measured colorimetrically (490nm) as per manufacturer's instructions. A standard curve for determination of cell numbers was constructed using cells seeded at 10<sup>2</sup> to 10<sup>6</sup>/well. Cell viability (shown as percentage of dead cells) was calculated by expressing medium LDH as a percentage of the total cellular LDH.

#### ***Quantification of apoptosis by flow cytometry***

VSMCs plated in 24-well trays were cultured in control or calcification medium ( $\pm$  treatment) for 7 days. Apoptosis was assessed via flow cytometry using an annexin V antibody conjugated to fluorescein (Life Technologies, Paisley, UK), as per manufacturer's instructions. Briefly, cells were detached using trypsin (0.25%) and resultant pellet washed in ice cold PBS. This suspension was centrifuged and resuspended in 1X annexin-binding buffer (Life Technologies, Paisley, UK). A sample of this suspension was incubated with the annexin V antibody for 15 minutes, after which, analysed using a BD FACSCanto II Flow Cytometer (Becton, Dickinson and Company, Oxford, UK). Data was processed to calculate percentage apoptosis using Flowing Software (version 2.5.1) (Turku University, Finland).

#### ***Determination of TNAP activity***

VSMCs and osteoblasts VSMCs were cultured with 1-100 $\mu$ M ATP, UTP or PP<sub>i</sub> for 14 days. TNAP activity was measured in *P2Y<sub>2</sub>R*<sup>-/-</sup>, *Enpp1*<sup>-/-</sup> or wildtype (*P2Y<sub>2</sub>R*<sup>+/+</sup> / *Enpp1*<sup>+/+</sup>) cell lysates using a colorimetric assay (Anaspec, CA, USA), as previously described (Orriss et al., 2012a). TNAP activity was normalised to cell protein using the Bradford assay.

#### ***Histology***

Histological analysis was performed on aortas obtained from 24-week old male *P2Y<sub>2</sub>R*<sup>+/+</sup> or *P2Y<sub>2</sub>R*<sup>-/-</sup> mice. Tissues were fixed in 10% neutral buffered formalin and embedded in paraffin wax blocks. Serial sections were cut every 5 $\mu$ m and mounted onto slides. Before staining, the samples were deparaffinised using xylene, then rehydrated through a series of decreasing ethanol solutions and finally water. Slides were stained with haematoxylin and eosin (H&E) to examine cell morphology and alizarin red to visualise vascular calcification.

***Statistical analysis***

Data were analysed using GraphPad Prism 6 software (San Diego, CA). Statistical comparisons were made using one-way or two-way analysis of variance (ANOVA) with a post-hoc Bonferroni correction for multiple comparisons. Results are expressed as means  $\pm$  SEM for 6 replicates and are representative of experiments performed at least three times using cells or tissues obtained from different animals.



## RESULTS

### ***P2 receptor expression is increased in calcifying VSMCs***

Analysis of mRNA expression revealed that mouse VSMCs express multiple P2 receptor subtypes including all the P2X receptors and the majority of the P2Y receptors (**Fig. 1A**). Expression of the P2X1, P2X2, P2X4, P2X5, P2X6 and P2Y<sub>2</sub> receptors was increased up to 3-fold in calcifying VSMCs compared to control VSMCs (**Fig.1A**).

### ***ATP, UTP and PP<sub>i</sub> inhibit VSMC calcification***

ATP ( $\geq 10\mu\text{M}$ ) and UTP ( $\geq 1\mu\text{M}$ ) dose dependently inhibited VSMC calcification by up to 80% (**Fig. 1B, 1C, 1G**). UTP was more potent at exerting inhibitory actions from  $1\mu\text{M}$ . ADP and UDP had no effect at any of the concentrations tested (**Fig. 1D, 1E, 1G**). The ubiquitous mineralisation inhibitor, PP<sub>i</sub> ( $\geq 10\mu\text{M}$ ), reduced VSMC calcification up to 85% (**Fig. 1F, 1G**). Representative images in **Fig. 1G** show the inhibitory effects of  $10\mu\text{M}$  ATP, UTP or PP<sub>i</sub> on calcification in mouse VSMC cultures.

### ***ATP, UTP and PP<sub>i</sub> increase VSMC viability and decrease apoptosis***

In calcifying VSMCs, treatment with ATP, UTP and PP<sub>i</sub> ( $\geq 10\mu\text{M}$ ) decreased the percentage of dead cells present by up to 75%, 60% and 80%, respectively (**Fig. 2A-2C**). ATP and UTP had no effect on osteoblast cell viability (**Fig. 2D, 2E**). PP<sub>i</sub> at the highest dose ( $100\mu\text{M}$ ) caused a 45% increase in osteoblast cell death.

The level of apoptosis was increased up to 2.3-fold in calcifying VSMCs compared to control VSMCs (**Fig 2G-2I**). Treatment with ATP, UTP and PP<sub>i</sub> ( $\geq 10\mu\text{M}$ ) decreased the level of apoptosis in calcifying cells to the level seen in control VSMCs (**Fig 2G-2I**).

### ***The effect of ATP, UTP and PP<sub>i</sub> on TNAP activity in calcifying VSMCs and osteoblasts***

ATP and UTP ( $\geq 1\mu\text{M}$ ) increased VSMC TNAP activity by up to 88% and 50%, respectively (**Fig. 3A, 3B**). PP<sub>i</sub> ( $\geq 1\mu\text{M}$ ), which is a substrate for TNAP, dose dependently stimulated enzyme activity by up to 80% (**Fig. 3C**). In contrast, ATP and UTP ( $\geq 1\mu\text{M}$ ) inhibited osteoblast TNAP activity by up to 50% (**Fig. 3D, 3E**). PP<sub>i</sub> had no effect on TNAP activity in osteoblasts (**Fig. 3F**). It should also be noted that the basal TNAP activity of mineralising osteoblasts was at least 100-fold higher than that of a calcifying VSMCs (**Fig. 3**).

### ***The P2Y<sub>2</sub> receptor does not mediate the effects of ATP & UTP on VSMC calcification***

Deletion of the P2Y<sub>2</sub> receptor had no effect on the basal level of calcification in VSMC or aortic ring cultures (**Fig. 4A, 4B**). The inhibitory effects of ATP and UTP on calcification were unchanged in P2Y<sub>2</sub>R<sup>-/-</sup> VSMCs (**Fig. 4C, 4D**). The selective P2Y<sub>2</sub> receptor antagonist, ARC118935XX, also failed to attenuate the inhibitory effects of UTP (**Fig. 4E**). Isolated aortas from 24-week old P2Y<sub>2</sub>R<sup>+/+</sup> and P2Y<sub>2</sub>R<sup>-/-</sup> mice showed no obvious differences in structure or cell morphology (**Fig. 4F**). Alizarin red staining revealed no spontaneous calcium deposition within any of the arterial layers (**Fig. 4F**). Representative images shown are from the descending aorta. The effects of ATP and UTP on VSMC cell viability and TNAP activity were also unaffected by the deletion of the P2Y<sub>2</sub> receptor (**Fig. 4G-4J**).

### ***Reduced inhibitory effects of ATP and UTP on bone mineralisation in P2Y<sub>2</sub>R<sup>-/-</sup> osteoblasts***

ATP ( $\geq 10\mu\text{M}$ ) and UTP ( $\geq 1\mu\text{M}$ ) dose dependently reduced bone mineralisation in P2Y<sub>2</sub>R<sup>+/+</sup> osteoblasts by up to 80% (**Fig. 5A, 5B, 5E**). The overall level of mineralised bone nodule formation was up to 60% higher in P2Y<sub>2</sub>R<sup>-/-</sup> cells compared to P2Y<sub>2</sub>R<sup>+/+</sup>. In P2Y<sub>2</sub>R<sup>-/-</sup> osteoblasts, the effects of ATP and UTP were only observed at  $100\mu\text{M}$  making them 10-fold and 100-fold less potent at inhibiting bone mineralisation, respectively (**Fig. 5A, 5B, 5E**). Representative whole well scans (unstained) and phase contrast microscopy images (alizarin red stained) of osteoblast cell layers show the reduced inhibitory effects of ATP and UTP in P2Y<sub>2</sub>R<sup>-/-</sup> osteoblasts. They also illustrate the increased level of bone formation in P2Y<sub>2</sub>R<sup>-/-</sup> cells (**Fig. 5E**).

***The inhibitory effects of ATP and UTP on TNAP activity are lost in P2Y<sub>2</sub>R<sup>-/-</sup> osteoblasts***

In P2Y<sub>2</sub>R<sup>+/+</sup> osteoblasts ATP (≥10μM) and UTP (≥1μM) reduce TNAP activity by up to 40%. P2Y<sub>2</sub>R<sup>-/-</sup> osteoblasts display basal TNAP activity levels that are up to 30% higher than wildtype cells. The inhibitory effects of ATP and UTP on TNAP activity were lost in P2Y<sub>2</sub>R<sup>-/-</sup> osteoblasts (Fig. 5C, 5D).

***The inhibitory effects of ATP and UTP on VSMC calcification and bone mineralisation are reduced in Enpp1<sup>-/-</sup> mice***

Since PP<sub>i</sub> mimics many of the actions of ATP and UTP we investigated whether the effects of these extracellular nucleotides involved P2 receptor independent mechanisms. In Enpp1<sup>+/+</sup> cells, ATP and UTP inhibit VSMC calcification from concentrations of 10μM and 1μM, respectively. In Enpp1<sup>-/-</sup> VSMCs, ATP and UTP are 10-fold less potent only exerting inhibitory effects at 100μM and 10μM, respectively (Fig. 6A, 6B).

Consistent with earlier studies, Enpp1<sup>-/-</sup> osteoblasts displayed increased levels of bone mineralisation compared to Enpp1<sup>+/+</sup> cells (Anderson et al., 2005; Orriss et al., 2015). The inhibitory effects of ATP and UTP on bone mineralisation were 10-fold and 100-fold less potent, respectively, in Enpp1<sup>-/-</sup> osteoblasts (Fig. 6C, 6D).

***NPP1 deletion does not blunt the effects of ATP and UTP on cell viability and TNAP activity***

The protective effects of ATP and UTP on cell viability were unchanged in cultures of Enpp1<sup>-/-</sup> VSMCs (Fig. 7A, 7B). Deletion of NPP1 also had no effect on the ATP/UTP-mediated increase in VSMC TNAP activity (Fig. 7C, 7D).

## DISCUSSION

This study compared the effects of extracellular nucleotides on AMC and bone mineralisation. We found that, whilst the functional effect was the same, the cellular mechanisms mediating the observed inhibitory actions of ATP and UTP differed. In VSMCs, ATP and UTP prevent calcification, at least in part, by reducing VSMC apoptosis. In contrast, ATP and UTP have no effect on osteoblast survival and instead block bone mineralisation via the inhibition of TNAP. Furthermore, the P2Y<sub>2</sub> receptor does not appear to mediate the actions of ATP and UTP on AMC but is involved in the effects on bone mineralisation. However, NPP1-mediated generation of PP<sub>i</sub> contributes to the inhibitory effects observed in both VSMCs and osteoblasts (see summary **Fig. 8**).

This study demonstrated, like others, that VSMCs express multiple P2X and P2Y receptor subtypes (Lewis and Evans, 2000; Wang et al., 2002). Consistent with earlier work that described protective effects of extracellular nucleotides on aortic valve calcification (Cote et al., 2012) and VSMC calcification (Villa-Bellosta and Sorribas, 2013), we also found that both ATP and UTP were inhibitory. UTP, which was more potent than ATP, only activates the P2Y<sub>2</sub> and P2Y<sub>4</sub> receptor subtypes. Given the involvement of the P2Y<sub>2</sub> receptor in inhibiting bone mineralisation (**Fig. 5**) (Hoebertz et al., 2002; Orriss et al., 2007), aortic valve calcification and intimal calcification (Cote et al., 2012; Qian et al., 2017), the P2Y<sub>2</sub> receptor appeared to be a strong candidate for mediating the effects in VSMCs. However, the effects of ATP and UTP on VSMC calcification were not lost in P2Y<sub>2</sub>R<sup>-/-</sup> cells or blocked by a selective P2Y<sub>2</sub> receptor antagonist. Furthermore, P2Y<sub>2</sub>R<sup>-/-</sup> VSMCs and aortic rings displayed no differences in the basal level of calcification *in vitro* and isolated aortas from 24-week old animals also showed no signs of AMC. It should be noted that aged animals were not studied here and so it is possible that the condition develops in older P2Y<sub>2</sub>R<sup>-/-</sup> mice. Taken together, these data suggest that ATP and UTP are not acting via the P2Y<sub>2</sub> receptor to regulate AMC.

Our findings contrast to previous work which has implicated the P2Y<sub>2</sub> receptor in the effects of nucleotides on valve calcification (Cote et al., 2012) and arterial intimal calcification (Qian et al., 2017). There are a number of potential explanations for these observed differences. Firstly, the lack of involvement of the P2Y<sub>2</sub> receptor in aortic VSMCs could represent variation in the cellular mechanisms that underpin these different forms of pathological vascular calcification. Secondly, it could be due to differences in the experimental models used since each investigation has utilised distinct *in vitro* and *in vivo* methods to study the different forms of vascular calcification. Finally, given that the expression of functional P2 receptors is highly variable between cell types (Burnstock, 2007), it could also reflect a lower level of P2Y<sub>2</sub> receptor expression in aortic VSMCs compared to valve interstitial cells or osteoblasts. Interestingly, the mRNA expression of several P2 receptors including the P2Y<sub>2</sub> receptor was upregulated in calcifying VSMCs compared to control cells. Since the expression of P2 receptors by osteoblasts is differentiation dependent (Dixon et al., 1997; Noronha-Matos et al., 2012; Orriss et al., 2006), this could reflect the changes in the VSMC phenotype which occur during the development of AMC.

NPP1 hydrolyses extracellular ATP and UTP to produce the key local inhibitor of mineralisation, PP<sub>i</sub>. Since the effects of PP<sub>i</sub> on VSMC calcification mimicked those of ATP and UTP, we used *Enpp1*<sup>-/-</sup> cells to determine the involvement of P2 receptor independent mechanisms. The effects of ATP and UTP were blunted in *Enpp1*<sup>-/-</sup> VSMCs and osteoblasts suggesting that NPP1-mediated hydrolysis to produce PP<sub>i</sub> does contribute to the observed inhibitory actions in both cell types. This is consistent with earlier work that examined the role of ATP-derived PP<sub>i</sub> in VSMC calcification (Prosdocimo et al., 2009; Prosdocimo et al., 2010) and bone formation (Orriss et al., 2013; Orriss et al., 2007). However, since the effects were not completely attenuated it suggests that other mechanisms are also involved. In osteoblasts, this is likely to be due to the P2 receptors that have been implicated in the regulation of bone mineralisation, such as the P2Y<sub>2</sub>, P2X<sub>1</sub> and P2X<sub>7</sub> receptors (Hoebertz et al., 2002; Orriss et al., 2012a; Orriss et al., 2007). In VSMCs, the other mechanisms leading the functional effects are less clear. Although unlikely

to be mediated by the P2Y<sub>2</sub> receptor, involvement of other P2 receptor subtypes (e.g. the P2Y<sub>4</sub> receptor) is probable and presents an area for future study. Furthermore, since VSMCs have been shown to also express functional NPP3 (Prosdocimo et al., 2009), a role for other ecto-nucleotidases cannot be discounted.

Increased VSMC apoptosis has been implicated in the development of vascular calcification (Proudfoot et al., 2000). In this study we found that, at the concentrations which inhibited calcification, ATP, UTP and PP<sub>i</sub> prevented the increase in apoptosis usually seen in calcifying VSMC cultures. The effect of extracellular nucleotides on cell viability was unchanged in *Enpp1*<sup>-/-</sup> VSMCs. This suggests that, although PP<sub>i</sub> itself appears to exert protective actions, ATP and UTP are working directly to promote VSMC survival in calcifying conditions. Apoptosis is thought to promote calcification because the VSMC-derived apoptotic bodies can act as a nucleation site for hydroxyapatite crystal formation (Proudfoot et al., 2000). Thus, the inhibitory effects of extracellular nucleotides (and PP<sub>i</sub>) on AMC could be mediated, at least in part, by their ability to reduce apoptosis. The overall result being that there are less apoptotic bodies on which calcification can be initiated. In contrast, extracellular nucleotides have no effect on osteoblast viability and so the inhibitory effects on bone mineralisation are not mediated via alterations in apoptosis.

These differential effects of extracellular nucleotides on cell survival may also illustrate the differences between physiological bone mineralisation and pathological AMC. Bone mineralisation is not associated with increased levels of osteoblast apoptosis and instead matrix vesicles are the primary nucleation site for hydroxyapatite formation (Anderson et al., 1975). The initiation of AMC appears more heterogeneous with multiple factors acting as nucleation sites for calcification including apoptotic bodies and mineralisation-competent vesicles (Kapustin and Shanahan, 2012; Proudfoot et al., 2000).

TNAP is a crucial enzyme in bone mineralisation, with deficiencies leading to hypophosphatasia a condition characterised by hypomineralisation and rickets (Caswell et al., 1991). Increased TNAP expression and activity has also been associated with AMC (Narisawa et al., 2007; Sheen et al., 2015). Consistent with earlier work, we confirmed that ATP and UTP inhibited TNAP activity in osteoblasts (Orriss et al., 2007). We also demonstrated that this inhibition involved P2Y<sub>2</sub> receptor activation since the effects on TNAP activity were lost in *P2Y<sub>2</sub>R*<sup>-/-</sup> osteoblasts. In contrast, at concentrations where calcification is inhibited by up to 80%, ATP and UTP stimulated VSMC TNAP activity. The reasons for this counterintuitive increase are unclear but since PP<sub>i</sub>, and to a lesser extent ATP and UTP, are hydrolysed by TNAP the higher substrate levels could lead to alterations in enzyme kinetics. It is important to note that, even when stimulated, the TNAP activity of calcifying VSMCs was at least 100-fold lower than that of mineralising osteoblasts *in vitro*. Taken together, our observations suggest two things; firstly, that whilst reduced TNAP activity contributes to the inhibition of bone mineralisation by ATP and UTP, alterations in TNAP activity are unlikely to mediate the functional effects of nucleotides on VSMC calcification. Instead the effects on VSMC apoptosis appear to predominate. Secondly, although TNAP is essential for bone mineralisation its role in AMC appears more complex.

It is widely accepted that the development of AMC shares similarities to physiological bone formation. Whilst the role of NPP1 was similar in both VSMCs and osteoblasts, this study demonstrated that there are also clear differences in the cellular mechanisms mediating the inhibitory effects of extracellular nucleotides. This suggests that some of the underlying processes leading to AMC and bone mineralisation are not the same. It is these differences which need to be fully understood in order to identify a drug target or therapeutic agent that can prevent or regress AMC without exerting negative actions on the skeleton.

## ACKNOWLEDGEMENTS

The authors are grateful for funding from the British Heart Foundation (grant number: PG/15/13/31296) and Arthritis Research UK (grant number: 19205).

## REFERENCES

- Abbracchio MP, Burnstock G. 1994. Purinoceptors: are there families of P2X and P2Y purinoceptors? *Pharmacol Ther* 64(3):445-475.
- Anderson HC, Cecil R, Sajdera SW. 1975. Calcification of rachitic rat cartilage in vitro by extracellular matrix vesicles. *Am J Pathol* 79(2):237-254.
- Anderson HC, Harmey D, Camacho NP, Garimella R, Sipe JB, Tague S, Bi X, Johnson K, Terkeltaub R, Millan JL. 2005. Sustained osteomalacia of long bones despite major improvement in other hypophosphatasia-related mineral deficits in tissue nonspecific alkaline phosphatase/nucleotide pyrophosphatase phosphodiesterase 1 double-deficient mice. *Am J Pathol* 166(6):1711-1720.
- Burnstock G. 2007. Physiology and pathophysiology of purinergic neurotransmission. *Physiol Rev* 87(2):659-797.
- Burnstock G, Arnett TR, Orriss IR. 2013. Purinergic signalling in the musculoskeletal system. *Purinergic Signal* 9(4):541-572.
- Burnstock G, Kennedy C. 1985. Is there a basis for distinguishing two types of P2-purinoceptor? *Gen Pharmacol* 16(5):433-440.
- Burnstock G, Ralevic V. 2014. Purinergic signaling and blood vessels in health and disease. *Pharmacol Rev* 66(1):102-192.
- Caswell AM, Whyte MP, Russell RG. 1991. Hypophosphatasia and the extracellular metabolism of inorganic pyrophosphate: clinical and laboratory aspects. *Crit Rev Clin Lab Sci* 28(3):175-232.
- Cote N, El HD, Pepin A, Guauque-Olarte S, Ducharme V, Bouchard-Cannon P, Audet A, Fournier D, Gaudreault N, Derbali H, McKee MD, Simard C, Despres JP, Pibarot P, Bosse Y, Mathieu P. 2012. ATP acts as a survival signal and prevents the mineralization of aortic valve. *J Mol Cell Cardiol* 52(5):1191-1202.
- Dixon CJ, Bowler WB, Walsh CA, Gallagher JA. 1997. Effects of extracellular nucleotides on single cells and populations of human osteoblasts: contribution of cell heterogeneity to relative potencies. *Br J Pharmacol* 120(5):777-780.
- Fish RS, Klootwijk E, Tam FW, Kleta R, Wheeler DC, Unwin RJ, Norman J. 2013. ATP and arterial calcification. *Eur J Clin Invest* 43(4):405-412.
- Fleisch H, Bisaz S. 1962. Mechanism of calcification: inhibitory role of pyrophosphate. *Nature* 195:911.
- Gartland A, Orriss IR, Rumney RM, Bond AP, Arnett T, Gallagher JA. 2012. Purinergic signalling in osteoblasts. *Front Biosci* 17:16-29.
- Hajjawi MO, MacRae VE, Huesa C, Boyde A, Millan JL, Arnett TR, Orriss IR. 2014. Mineralisation of collagen rich soft tissues and osteocyte lacunae in *Enpp1*<sup>-/-</sup> mice. *Bone* 69C:139-147.
- Hoebertz A, Mahendran S, Burnstock G, Arnett TR. 2002. ATP and UTP at low concentrations strongly inhibit bone formation by osteoblasts: a novel role for the P2Y<sub>2</sub> receptor in bone remodeling. *J Cell Biochem* 86(3):413-419.
- Homolya L, Watt WC, Lazarowski ER, Koller BH, Boucher RC. 1999. Nucleotide-regulated calcium signaling in lung fibroblasts and epithelial cells from normal and P2Y<sub>2</sub> receptor (-/-) mice. *J Biol Chem* 274(37):26454-26460.
- Johnson K, Goding J, Van Etten D, Sali A, Hu SI, Farley D, Krug H, Hessle L, Millan JL, Terkeltaub R. 2003. Linked deficiencies in extracellular PP(i) and osteopontin mediate pathologic calcification associated with defective PC-1 and ANK expression. *J Bone Miner Res* 18(6):994-1004.
- Kapustin AN, Shanahan CM. 2012. Calcium regulation of vascular smooth muscle cell-derived matrix vesicles. *Trends Cardiovasc Med* 22(5):133-137.

Lewis CJ, Evans RJ. 2000. Lack of run-down of smooth muscle P2X receptor currents recorded with the amphotericin permeabilized patch technique, physiological and pharmacological characterization of the properties of mesenteric artery P2X receptor ion channels. *Br J Pharmacol* 131(8):1659-1666.

Mackenzie NC, Staines KA, Zhu D, Genever P, Macrae VE. 2014. miRNA-221 and miRNA-222 synergistically function to promote vascular calcification. *Cell Biochem Funct* 32(2):209-216.

Mackenzie NC, Zhu D, Milne EM, van 't HR, Martin A, Darryl QL, Millan JL, Farquharson C, Macrae VE. 2012. Altered bone development and an increase in FGF-23 expression in *Enpp1(-/-)* mice. *PLoS ONE* 7(2):e32177.

Narisawa S, Harmey D, Yadav MC, O'Neill WC, Hoylaerts MF, Millan JL. 2007. Novel inhibitors of alkaline phosphatase suppress vascular smooth muscle cell calcification. *J Bone Miner Res* 22(11):1700-1710.

Noronha-Matos JB, Coimbra J, Sa-e-Sousa A, Rocha R, Marinhos J, Freitas R, Guerra-Gomes S, Ferreirinha F, Costa MA, Correia-de-Sa P. 2014. P2X7-induced zeiosis promotes osteogenic differentiation and mineralization of postmenopausal bone marrow-derived mesenchymal stem cells. *FASEB J* 28(12):5208-5222.

Noronha-Matos JB, Correia-de-Sa P. 2016. Mesenchymal Stem Cells Ageing: Targeting the "Purinome" to promote osteogenic differentiation and bone repair. *J Cell Physiol* 231(9):1852-1861.

Noronha-Matos JB, Costa MA, Magalhaes-Cardoso MT, Ferreirinha F, Pelletier J, Freitas R, Neves JM, Sevigny J, Correia-de-Sa P. 2012. Role of ecto-NTPDases on UDP-sensitive P2Y(6) receptor activation during osteogenic differentiation of primary bone marrow stromal cells from postmenopausal women. *J Cell Physiol* 227(6):2694-2709.

Orriss IR. 2015. The role of purinergic signalling in the musculoskeletal system. *Auton Neurosci* (191):124-134.

Orriss IR, Arnett TR, Russell RG. 2016. Pyrophosphate: a key inhibitor of mineralisation. *Curr Opin Pharmacol* 28:57-68.

Orriss IR, Gueneri D, Hajjawi MO, Shaw K, Patel JJ, Arnett TR. 2017. Activation of the P2Y2 receptor regulates bone cell function by enhancing ATP release. *J Endocrinol* [epub ahead of print]

Orriss IR, Hajjawi MO, Huesa C, MacRae VE, Arnett TR. 2014. Optimisation of the differing conditions required for bone formation in vitro by primary osteoblasts from mice and rats. *Int J Mol Med* 34(5):1201-1208.

Orriss IR, Key ML, Brandao-Burch A, Patel JJ, Burnstock G, Arnett TR. 2012a. The regulation of osteoblast function and bone mineralisation by extracellular nucleotides: The role of P2X receptors. *Bone* 51(3):389-400.

Orriss IR, Key ML, Hajjawi MO, Arnett TR. 2013. Extracellular ATP released by osteoblasts is a key local inhibitor of bone mineralisation. *PloS one* 8(7):e69057.

Orriss IR, Key ML, Hajjawi MO, Millan JL, Arnett TR. 2015. Acidosis is a key regulator of osteoblast ecto-nucleotide pyrophosphatase/phosphodiesterase 1 (NPP1) expression and activity. *J Cell Physiol* 230:3049-3056.

Orriss IR, Knight GE, Ranasinghe S, Burnstock G, Arnett TR. 2006. Osteoblast responses to nucleotides increase during differentiation. *Bone* 39(2):300-309.

Orriss IR, Taylor SE, Arnett TR. 2012b. Rat osteoblast cultures. *Methods Mol Biol* 816:31-41.

Orriss IR, Utting JC, Brandao-Burch A, Colston K, Grubb BR, Burnstock G, Arnett TR. 2007. Extracellular nucleotides block bone mineralization in vitro: evidence for dual inhibitory mechanisms involving both P2Y2 receptors and pyrophosphate. *Endocrinology* 148(9):4208-4216.

Panupinthu N, Zhao L, Possmayer F, Ke HZ, Sims SM, Dixon SJ. 2007. P2X7 nucleotide receptors mediate blebbing in osteoblasts through a pathway involving lysophosphatidic acid. *J Biol Chem* 282(5):3403-3412.

Prosdocimo DA, Douglas DC, Romani AM, O'Neill WC, Dubyak GR. 2009. Autocrine ATP release coupled to extracellular pyrophosphate accumulation in vascular smooth muscle cells. *Am J Physiol Cell Physiol* 296(4):C828-C839.

Prosdocimo DA, Wyler SC, Romani AM, O'Neill WC, Dubyak GR. 2010. Regulation of vascular smooth muscle cell calcification by extracellular pyrophosphate homeostasis: synergistic modulation by cyclic AMP and hyperphosphatemia. *Am J Physiol Cell Physiol* 298(3):C702-713.

Proudfoot D, Skepper JN, Hegyi L, Bennett MR, Shanahan CM, Weissberg PL. 2000. Apoptosis regulates human vascular calcification in vitro: evidence for initiation of vascular calcification by apoptotic bodies. *Circ Res* 87(11):1055-1062.

Qian S, Regan JN, Shelton MT, Hoggatt A, Mohammad KS, Herring PB, Seye CI. 2017. The P2Y2 nucleotide receptor is an inhibitor of vascular calcification. *Atherosclerosis* 257:38-46.

Rutsch F, Vaingankar S, Johnson K, Goldfine I, Maddux B, Schauerte P, Kalhoff H, Sano K, Boisvert WA, Superti-Furga A, Terkeltaub R. 2001. PC-1 nucleoside triphosphate pyrophosphohydrolase deficiency in idiopathic infantile arterial calcification. *Am J Pathol* 158(2):543-554.

Sali A, Favalaro JM, Terkeltaub R, Goding JW. 1999. Germline deletion of the nucleoside triphosphate pyrophosphohydrolase (NTPPPH) plasma cell membrane glycoprotein (PC-1) produces abnormal calcification of the periarticular tissues. In: Vanduffe L, Lemmens R, editors *Ecto-ATPases and related ectonucleotides*, Maastricht, the Netherlands: Shaker Publishing BV:267-282.

Schuchardt M, Tolle M, Pruffer J, Pruffer N, Huang T, Jankowski V, Jankowski J, Zidek W, van der Giet M. 2012. Uridine adenosine tetraphosphate activation of the purinergic receptor P2Y enhances in vitro vascular calcification. *Kidney Int* 81(3):256-265.

Sheen CR, Kuss P, Narisawa S, Yadav MC, Nigro J, Wang W, Chhea TN, Sergienko EA, Kapoor K, Jackson MR, Hoylaerts MF, Pinkerton AB, O'Neill WC, Millan JL. 2015. Pathophysiological role of vascular smooth muscle alkaline phosphatase in medial artery calcification. *J Bone Miner Res* 30(5):824-836.

Shroff RC, Shanahan CM. 2007. The vascular biology of calcification. *Semin Dial* 20(2):103-109.

Staines KA, Zhu D, Farquharson C, MacRae VE. 2014. Identification of novel regulators of osteoblast matrix mineralization by time series transcriptional profiling. *J Bone Miner Metab* 32(3):240-251.

Taylor SE, Shah M, Orriss IR. 2014. Generation of rodent and human osteoblasts. *BoneKEy Reps* 3:585.

Villa-Bellosta R, Sorribas V. 2013. Prevention of vascular calcification by polyphosphates and nucleotides- role of ATP. *Circ J* 77(8):2145-2151.

Villa-Bellosta R, Wang X, Millan JL, Dubyak GR, O'Neill WC. 2011. Extracellular pyrophosphate metabolism and calcification in vascular smooth muscle. *Am J Physiol Heart Circ Physiol* 301(1):H61-68.

Wang L, Karlsson L, Moses S, Hultgardh-Nilsson A, Andersson M, Bornha C, Gudbjartsson T, Jern S, Erlinge D. 2002. P2 receptor expression profiles in human vascular smooth muscle and endothelial cells. *J Cardiovasc Pharmacol* 40(6):841-853.

Xing Y, Gu Y, Bresnahan JJ, Paul EM, Donahue HJ, You J. 2014. The roles of P2Y2 purinergic receptors in osteoblasts and mechanotransduction. *PLoS ONE* 9(9):e108417.

Young MJ, Adams JE, Anderson GF, Boulton AJ, Cavanagh PR. 1993. Medial arterial calcification in the feet of diabetic patients and matched non-diabetic control subjects. *Diabetologia* 36(7):615-621.

Zhu D, Mackenzie NC, Farquharson C, Macrae VE. 2012. Mechanisms and clinical consequences of vascular calcification. *Front Endocrinol(Lausanne)* 3:95.

Zhu D, Mackenzie NC, Millan JL, Farquharson C, Macrae VE. 2011. The appearance and modulation of osteocyte marker expression during calcification of vascular smooth muscle cells. *PLoS ONE* 6(5):e19595.

## FIGURE LEGENDS

### Figure 1. *The functional effects of extracellular nucleotides on VSMC calcification*

(A) VSMCs express mRNA for multiple P2 receptor subtypes; expression of the P2X1, P2X2, P2X4, P2X5, P2X6 and P2Y<sub>2</sub> receptors is increased up to 3-fold in calcifying cells compared to control cells. (B) ATP and (C) UTP dose dependently ( $\geq 1\mu\text{M}$ ) inhibit VSMC calcification by up to 80%. (D) ADP and (E) UDP have no effect on VSMC calcification. (F) PP<sub>i</sub> ( $\geq 10\mu\text{M}$ ) reduces VSMC calcification by up to 85%. Values are mean  $\pm$  SEM ( $n = 4$  (mRNA only) or 6), \* =  $p < 0.05$ , \*\* =  $p < 0.01$ , \*\*\* =  $p < 0.001$ . (G) Representative light microscopy images of alizarin red stained VSMC cultures treated with extracellular nucleotides. Scale bar = 50 $\mu\text{m}$

### Figure 2. *ATP, UTP and PPi reduce VSMC cell death but have no effect on osteoblast survival*

(A) ATP, (B) UTP and (C) PP<sub>i</sub> dose dependently decrease the proportion of dead cells in calcifying VSMC cultures by up to 80%. No effect of (D) ATP or (E) UTP on osteoblast viability. (F) PP<sub>i</sub> at the highest dose tested caused a small increase in the percentage of dead cells. (G, H, I) Culture of VSMCs in calcifying conditions was associated by an increased level of apoptosis (up to 2.3 fold). ATP, UTP and PP<sub>i</sub> ( $\geq 10\mu\text{M}$ ) reduced the amount of apoptosis to levels seen in non-calcifying cells. Values are mean  $\pm$  SEM ( $n = 6$ ), \* =  $p < 0.05$ , \*\* =  $p < 0.01$ , \*\*\* =  $p < 0.001$ .

### Figure 3. *Opposing effects of extracellular nucleotides on TNAP activity in VSMCs and osteoblasts*

(A) ATP, (B) UTP and (C) PP<sub>i</sub> ( $\geq 1\mu\text{M}$ ) increased VSMC TNAP activity by up to 88%. (D) ATP and (E) UTP reduced osteoblast TNAP activity by up to 50%. (F) No effect of PP<sub>i</sub> on osteoblast TNAP activity. Note that the basal levels of TNAP activity are approximately 100-fold higher in mineralising osteoblasts than calcifying VSMCs. Values are mean  $\pm$  SEM ( $n = 6$ ), \* =  $p < 0.05$ , \*\* =  $p < 0.01$ , \*\*\* =  $p < 0.001$ .

### Figure 4. *Lack of involvement of the P2Y<sub>2</sub> receptor in regulating AMC*

(A) Cultured VSMCs and (B) aortic rings from  $P2Y_2R^{-/-}$  animals displayed no differences in the basal level of calcification. The inhibitory effects of (C) ATP and (D) UTP were unchanged in  $P2Y_2R^{-/-}$  VSMCs. (E) The selective P2Y<sub>2</sub> receptor antagonist, ARC118925XX, did not block the inhibitory effects of UTP on VSMC calcification. (F) Histological analysis of isolated aortas from 24-week old  $P2Y_2R^{-/-}$  mice showed no difference in cell morphology or the presence of AMC. Scale bar = 100 $\mu\text{m}$ . The protective effects of (G) ATP and (H) UTP on cell viability were unaffected by deletion of the P2Y<sub>2</sub> receptor. The stimulatory actions of (I) ATP and (J) UTP on TNAP activity were unchanged in  $P2Y_2R^{-/-}$  VSMCs. Values are mean  $\pm$  SEM ( $n = 6$ ), \* =  $p < 0.05$ , \*\* =  $p < 0.01$ , \*\*\* =  $p < 0.001$ .

### Figure 5. *The role of the P2Y<sub>2</sub> receptor in the inhibitory effects of ATP and UTP on bone mineralisation*

The inhibitory effects of (A) ATP and (B) UTP were reduced 10-fold and 100-fold, respectively, in  $P2Y_2R^{-/-}$  osteoblasts; mineralised bone nodule formation was 60% higher in  $P2Y_2R^{-/-}$  osteoblasts. The inhibitory actions of (C) ATP and (D) UTP on TNAP activity were lost in  $P2Y_2R^{-/-}$  osteoblasts. Basal TNAP activity was increased up to 40% in  $P2Y_2R^{-/-}$  cells. Values are mean  $\pm$  SEM ( $n = 6$ ), \* =  $p < 0.05$ , \*\* =  $p < 0.01$ , \*\*\* =  $p < 0.001$ . Comparison of  $P2Y_2R^{+/+}$  to  $P2Y_2R^{-/-}$ : # =  $p < 0.05$ , ## =  $p < 0.01$ , ### =  $p < 0.001$ . (E) Representative whole well scans (unstained) and phase contrast microscopy images (alizarin red stained) showing the effects of ATP and UTP on bone mineralisation in  $P2Y_2R^{-/-}$  and  $P2Y_2R^{+/+}$  osteoblasts. Areas of unmineralised matrix are highlighted by the arrows. Scale bars: whole well scans = 0.5cm, microscopy images = 200 $\mu\text{m}$ .



**Figure 6. The actions of NPP1 contribute towards the inhibitory effects of ATP and UTP on VSMC calcification and bone mineralisation**

The inhibitory effects of (A) ATP and (B) UTP on calcification were reduced 10-fold in *Enpp1*<sup>-/-</sup> VSMCs. The decrease in bone mineralisation caused by (C) ATP and (D) UTP was 10-fold and 100-fold lower, respectively, in *Enpp1*<sup>-/-</sup> osteoblasts. Values are mean ± SEM (*n* = 6), \* = *p*<0.05, \*\* = *p*<0.01, \*\*\* = *p*<0.001. Comparison of *Enpp1*<sup>+/+</sup> to *Enpp1*<sup>-/-</sup>: # = *p*<0.05, ### = *p*<0.001.

**Figure 7. The effect of ATP and UTP on VSMC survival and TNAP activity does not involve NPP1**

The protective effects of (A) ATP and (B) UTP on cell viability were unchanged in *Enpp1*<sup>-/-</sup> VSMCs. The stimulatory actions of (C) ATP and (D) UTP on TNAP activity were the same in *Enpp1*<sup>+/+</sup> and *Enpp1*<sup>-/-</sup> cells. Values are mean ± SEM (*n* = 6), \* = *p*<0.05, \*\*\* = *p*<0.001.


**Figure 8. Summary of the differing cellular mechanisms by which ATP and UTP exert their effects on AMC and bone mineralisation**

ATP and UTP inhibit AMC and bone mineralisation but there are some differences in the mechanisms mediating these actions. In VSMCs (top), ATP and UTP act via P2 receptors to decrease apoptosis and, as a consequence, calcification. In osteoblasts (OBs, bottom panel), ATP and UTP act via the P2Y<sub>2</sub> receptor to inhibit alkaline phosphatase (TNAP) activity. However, for both cell types hydrolysis of ATP and UTP to produce PP<sub>i</sub> by NPP1 also contributes to the inhibitory effects observed.


**Table 1: Primer sequences for real time PCR**

<b>Gene</b>	<b>Primer sequence (5'-3')</b>	
<b>P2X<sub>1</sub></b>	<b>S</b>	tgt acg ggg aga aga acc tg
	<b>AS</b>	tcc caa aca cct tga aga gg
<b>P2X<sub>2</sub></b>	<b>S</b>	cgt ctt cat cgt gca gaa aa
	<b>AS</b>	cac ttt gtg ttc cga cat gg
<b>P2X<sub>3</sub></b>	<b>S</b>	tac caa gtc ggt ggt tgt ga
	<b>AS</b>	cca ccc cac aaa gta gga ga
<b>P2X<sub>4</sub></b>	<b>S</b>	gca ccc tcc acc atc tct aa
	<b>AS</b>	aaa cct ctt gcc aga agc aa
<b>P2X<sub>5</sub></b>	<b>S</b>	ggg ctt tct tct gtg acc tg
	<b>AS</b>	gtt ggc ctc aac ctc aac at
<b>P2X<sub>6</sub></b>	<b>S</b>	agc cat ggc ata aaa act gg
	<b>AS</b>	gtg aag ttc ttg gcc tga gc
<b>P2X<sub>7</sub></b>	<b>S</b>	ggc act gga gga aaa ttt ga
	<b>AS</b>	tga gca agt caa tgc aca ca
<b>P2Y<sub>1</sub></b>	<b>S</b>	agg aaa gct tcc agg agg ag
	<b>AS</b>	cgt gtc tcc att ctg ctt ga
<b>P2Y<sub>2</sub></b>	<b>S</b>	gtc agc agt gac gac tca aga c
	<b>AS</b>	tca gag gat atc agc ccc ttt a
<b>P2Y<sub>4</sub></b>	<b>S</b>	agg aag cag cag aac acc at
	<b>AS</b>	caa gga gtc tgc act ggt ca
<b>P2Y<sub>6</sub></b>	<b>S</b>	ttc cat ctt gca tga gac aga
	<b>AS</b>	gct tga aat cct cac ggt aga
<b>P2Y<sub>13</sub></b>	<b>S</b>	ggc cac tag atg tca cct ttt c
	<b>AS</b>	gat ggt ggg gtg gta act aga a


**Figure 1**



**Figure 2**



**Figure 2**



**Figure 2**






Figure 3




**Figure 4**





**Figure 5**



**Figure 6**




Figure 7




Figure 8





Figure 3



**Figure 6**




Figure 7





Figure 3





**Figure 6**




Figure 7

