

This is the peer-reviewed, manuscript version of an article published in *Global Food Security*. The version of record is available from the journal site:

<https://doi.org/10.1016/j.gfs.2018.03.003>.

© 2018. This manuscript version is made available under the CC-BY-NC-ND 4.0 license

<http://creativecommons.org/licenses/by-nc-nd/4.0/>.

The full details of the published version of the article are as follows:

TITLE: Assessing the chemical and microbiological quality of farmed tilapia in Egyptian fresh fish markets

AUTHORS: Eltholth, M; Fornace, K; Grace, D; Rushton, J; Haesler, B

JOURNAL: Global Food Security

PUBLISHER: Elsevier

PUBLICATION DATE: June 2018

DOI: 10.1016/j.gfs.2018.03.003

1 Title: Assessing the chemical and microbiological quality of farmed tilapia in Egyptian fresh
2 fish markets

3 Running title: Quality of farmed tilapia in Egyptian fresh fish markets

4

5

6

7

8

1 ABSTRACT

2 Fish make important contributions to food and nutrition security in low and middle income
3 countries; however, they are also prone to contamination with a range of chemical and
4 biological hazards. The presence of people's perception and health hazards has implications
5 for consumer acceptability and hence the potential contribution of fish to nutrition and health.
6 The aim of this study was to assess the chemical and microbiological quality of farmed tilapia
7 in Egypt. We conducted a systematic literature review resulting in 38 papers meeting inclusion
8 criteria. We also conducted a survey of seven hazardous chemicals in fish sampled from farms
9 (300 samples from 100 farms) and of 5 biological hazards as well as total bacterial counts in
10 fish sampled from retailers (300 samples from 100 retailers). The results showed that the level
11 of contamination with heavy metals and pesticides was lower than the national and
12 international permissible limits. On the other hand level of contamination of a considerable
13 proportion of samples with microbial pollutants was higher than the permissible limits. Results
14 from the literature indicated that, the level of contamination of wild tilapia was higher than
15 farmed tilapia, again in contradiction to common perceptions. Our results indicate that the risk
16 of human exposure to heavy metals and pesticides via consumption of farmed tilapia is
17 negligible compared to microbial hazards. These findings suggest that post-harvest
18 contamination is the major health risk in the tilapia fish value chain and we make
19 recommendations for addressing this.

20 Key words:

21 Farmed Tilapia; Heavy metals; Pesticide residues; Egypt

22

23

1 **1. INTRODUCTION**

2 Globally, aquaculture accounts for about 50% of fish consumed and is one of the fastest
3 growing food sectors (Subasinghe et al., 2009) with a growth rate of approximately 6.2% in
4 2011 (FAO, 2013). In low and middle income countries, most farmed fish (>80%) is produced
5 in fresh water by small scale producers (Hastein et al., 2006). While fish is an important source
6 of low fat protein rich food and omega-three/omega-six fatty acids that protect against adverse
7 health effects, such as coronary heart disease and stroke (Domingo, 2007), there are also
8 increasing concerns about foodborne hazards, chemical and microbial, that might be present in
9 fish. These concerns can also result in decreasing demand for farmed fish (Smallwood and
10 Blaylock 19991). This could negatively affect fish farmers and retailers and decrease
11 consumption and use of animal source food.

12 Egypt is the largest aquaculture producer in Africa and among the top 10 producer countries
13 worldwide; in 2015 the aquaculture production was 1.5 million tonnes¹. Published data on
14 public health hazards associated with farmed tilapia in the Egyptian fresh fish value chains are
15 scarce. Given the unprecedented production and consumption of tilapia in Egypt of over half a
16 million tonnes in 2011 (Macfadyen et al., 2012), it is critical to get an understanding of potential
17 contamination of this important food. A characterisation of farmed tilapia production,
18 marketing and consumption patterns in the Nile Delta identified various potential points of
19 potential contamination (Eltholth et al., 2015). Agriculture drainage canals, which contain
20 water that has been used for agricultural activities and constitute the main water supply for
21 most fish farms in this area were identified as sources of pesticide residues, runoff derived
22 fertilizers and metals that may contaminate farmed fish (Authman et al., 2013, Authman et al.,
23 2012, Authman, 2011, Authman and Abbas, 2007).

24
25 The aim of this study was to assess the chemical and microbiological quality of farmed tilapia
26 in the Nile Delta, Egypt. The objectives were 1) to measure chemical and biological
27 contamination levels at farm and retail levels, respectively and 2) to compare detected
28 contamination with values published in the scientific literature. Outcomes of this study are
29 important inputs for risk assessment and appraisal of strategies for management of human
30 exposure to chemical and/or microbiological health hazards via consumption of farmed tilapia.

¹ <https://dailynewsegypt.com/2017/02/20>

2. MATERIALS AND METHODS

2.1. Study site and sampling

For the assessment of chemical pollutants, 300 whole tilapia of at least 100g were collected from 100 fish farms (three each) in Kafrelsheikh governorate, the main fish producing area in Egypt, in which about 55% of the farmed fish is produced (Macfadyen et al., 2011). Farms were selected randomly (Eltholth et al., 2015). Visits were scheduled with the owner, manager, or a worker who was authorised to talk to the enumerators. Upon visiting the farm, three fish from one pond, given that all ponds within the farm were connected, were collected in sterile plastic bags. Samples were transported to Kafrelsheikh University Central Laboratory of Environmental Studies (KUCLES) on ice in an ice box as soon as possible.

For the assessment of microbiological pollutants, tilapia samples were collected from 100 fresh fish retailers (20 wholesalers, 56 retailers and 24 street vendors) from Kafrelsheikh, Gharbia and Menoufia governorates in the Nile Delta. The target fish retailers were those serving the consumers in the study area; they were traced from consumers interviewed as described elsewhere (Eltholth et al., 2015). From each retailer three whole tilapia of at least 100g were taken from tilapia boxes offered for sale to customers. To make sure that tilapia were locally produced, retailers were asked about the source of tilapia. All samples were collected in sterile plastic bags and transported to the Central Diagnostic and Research Laboratory, Faculty of Veterinary Medicine Kafrelsheikh University on ice in an ice box.

2.2. Laboratory analysis

Samples were prepared and analysed for chemical and microbial contaminants according to the standard methods. Detailed description of the analysis for chemical and microbial contaminants is available in appendix 1.

2.3. Literature review

As consumers are more concerned about chemical pollutants of farmed tilapia and their perception about safety of farmed tilapia. We aimed to compare detected contamination with values published in the scientific literature, a systematic literature review was carried out. The following databases were searched: PubMed and Google Scholar. For the initial identification of primary studies the search terms “heavy metals”, “arsenic”, “lead”, “cadmium”, “mercury”, “muscles”, “pesticides”, “residues”, “organophosphorus” and “organochlorines” were applied in combination with “Egypt” AND “tilapia” using AND. The searches of Google Scholar were restricted to articles published since 2000 to December 2013. Titles and abstracts were screened

1 for primary identification of relevant studies according to the following criteria: (i) published
2 in a scientific journal, (ii) in English language and (iii) addressed at least one of the chemical
3 pollutants under the scope of this study. The following studies were excluded: review articles,
4 experimental studies or trials for assessing the impact of pollution on tilapia, and studies
5 investigating the contamination of fish species other than *Tilapia niloticus*. Additionally,
6 studies investigating the contamination of tilapia organs other than muscles were excluded, as
7 only muscle tissue is commonly consumed. Relevant articles were imported to the Endnote
8 database, duplicates were removed and the full text was obtained. Next, the articles were read
9 in full and the concentration of pollutants under the scope of this study was recorded. Simple
10 descriptive statistics using excel was used to summarize the findings of relevant studies.

11 2.4. Ethical approval

12 The present study received approval from the Ethics and Welfare Committee of the Royal
13 Veterinary College, London, UK (reference number URN 2012 1191) and Kafrelsheikh
14 University, Egypt. Oral informed consent was obtained from each study participant after
15 reading written consent form. The interviewers confirmed the participants' oral consent by
16 ticking the relevant boxes on the hardcopies. The consent form mainly explained about the
17 purpose of the study, the risks and benefits of participation in the study, conditions of
18 confidentiality and the right to refusal or withdrawal from the study.

19 3. RESULTS

20 3.1. Chemical analysis of tilapia samples from fish farms

21 Six pesticides and one heavy metal were detected (aldrin, dieldrin, endrin, heptachlor,
22 heptachlor-epoxide (beta) and lindane pesticides and mercury respectively). All pesticide
23 results were well within the maximum permissible limits (MPL) defined by Codex
24 Alimentarius Commission (CAC, 2009). The concentration of mercury in all contaminated
25 samples was lower than the MPL (0.50 ppm) stated by Egyptian Organization for
26 Standardization and Quality (EOS, 2010). Arsenic, lead and cadmium were not detected in any
27 of the samples. Results for the chemical hazards in farmed tilapia are in table 1.

28 **Table 1: The concentrations of pesticide residues in tilapia from fish farms in Kafrelsheikh governorate, Egypt**

Concentration (ppb*)	Pesticides					
	Aldrin	Dieldrin	Endrin	Heptachlor	Heptachlor- Epoxide	Lindane

Minimum	7.48	9.2	0.34	0.8	0.34	1.2
Maximum	35.42	33.6	12.4	6.96	2.91	5.9
Mean	19.18	16.78	2.37	2.68	1.29	3.04
Standard deviation	7.64	6.18	2.69	1.48	0.56	1.33
MPL**	300	300	300	200	200	200

*ppb=part per billion, **MPL= Maximum permissible limits adapted from (Yahia and Elsharkawy, 2014)

3.2. Bacteriological analysis of fish samples from retail sale

The mean total counts of aerobic plate count (APC), *E. coli*, *S. aureus* and *V. parahaemolyticus* in positive samples are listed in table 2. The proportions of samples with count higher than the MPL stated by EOS (EOS 2005) were 13.7%, 8.0%, 7.7%, 3.3%, 13% and 12.3% for APC, *E. coli*, *L. monocytogenes*, *Salmonella*, *S. aureus* and *V. parahaemolyticus*, respectively. Out of the tested samples, only 64.3% complied with all bacteriological standards: the level of APC and tested pathogens were under the MPL stated by the EOS. The mean APC (6.2×10^5 cfu/g) for samples collected from the whole sale market was lower than that from the retail markets and the latter was lower than that from street vendors.

Table 2: Bacteriological characterisation of fresh tilapia samples collected from retail sale in the Nile Delta of Egypt

Foodborne pathogens	Bacterial count		Egyptian standard cfu/g*	Samples > Egyptian standards (%)
	Mean	SD		
Total aerobic count (10^5 cfu/g)	6.2	3.80	1×10^6	13.7
<i>E. coli</i> (10^3 cfu/g)	0.73	0.54	100	8.0
<i>L. monocytogenes</i> **	NA	NA	0.0	7.7
<i>Salmonella</i> spp.**	NA	NA	0.0	3.3
<i>S. aureus</i> (10^3 cfu/g)	1.17	1.03	1000	13.0
<i>V. parahaemolyticus</i> (10^3 cfu/g)	0.89	0.91	0.0	12.3

*MPL stated by EOS (EOS, 2005) for fresh chilled fish, ** The count for *L. monocytogenes* and *Salmonella* spp. Was not presented as enrichment steps were used.

3.3.Literature review for chemical pollutants in tilapia

The initial search revealed 6108 publications. After exclusion of duplications and applying the inclusion criteria, 38 papers were retained. From these articles data were extracted and summarised in appendix 2 and 3 for pesticide residues and heavy metals, respectively. Not all data from relevant papers were extracted, only data concerned with the pollutants under the scope of this study. Of the studies included, 8.7% and 14.3% were concerned with the detection of pesticide residues and heavy metals in cultured fish, respectively. The majority of relevant studies were concerned with contamination of water and wild tilapia in Egyptian lakes and the level of pollution in the Nile tributaries and agricultural draining canals. The level of pesticide residues reported in all studies was lower than the MPL, Table 3. The concentration of heavy metals was higher than the MPL in most of cases. Results from these studies suggest that the level of contamination of wild tilapia, particularly from lakes and drainage canals, was higher than that of farmed tilapia. Also the average level of contamination, particularly with heavy metals, was higher than that detected in the current study.

Table 3: Summary of the results from scientific literature for heavy metals in tilapia muscles collected from different sources in Egypt.

Concentration (ppm)	Lead	Cadmium	Mercury
Minimum	0.002	0.01	0.01
Maximum	48.7	10.85	67.10
Mean	4.49	1.12	8.69
SD	8.09	2.32	19.75
MPL*	0.30	0.05	0.50

MPL= maximum permissible limits for heavy metals in fish stated by Egyptian Organization for Standardization and Quality (EOS, 2010).

4. DISCUSSION

This study assessed the prevalence and level of contamination with selected chemical and microbiological pollutants in farmed tilapia produced and marketed as fresh fish in the Nile delta of Egypt. The results of this study showed that the level of contamination of farmed tilapia, in the Nile delta, with pesticide residues and heavy metals was within the MPL. However the level of contamination with certain microbiological contaminants was higher than the MPL which could be due to post-harvesting contamination.

4.1. Chemical pollutants in farmed tilapia

Six pesticide residues were detected and the level was within the MPL according to CAC (CAC, 2009). According to the Egyptian standards “*the maximum levels should be set at a strict level which is reasonably achievable by following good agricultural, fishery and manufacturing practices and taking into account the risk related to the consumption of the food*” (EOS, 2010). In general, the level of detected chemical pollutants was lower than that from previous studies particularly for tilapia samples collected from lakes and river Nile tributaries. These results contradict the perception of many consumers that wild caught tilapia are safer than farmed one (Eltholth et al., 2015). The low level of heavy metals in tilapia tissues in this study may be due to the frequent changing of water in fish ponds and the use of branded fish feed by more than 90% of fish farmers (Eltholth et al., 2015). Apart from water sources, management, type of fish feed and feeding system may influence the level of contamination of fish tissues with metals and/or pesticide residues (Trocino et al., 2012). The metal concentration in different parts of tilapia varies according to the metabolic activities of the organs and the routes of exposure. Its concentration in muscle tissues is lower than other organs (Authman et al., 2012, Authman et al., 2013). This variation of concentration and distribution of chemical pollutants indicate that the risk of human exposure to chemical pollutants via consumption of tilapia will depend on the way of processing, cooking and consumption behaviour. The main threats to human health from heavy metals are from lead, cadmium, mercury and arsenic (Järup, 2003). Serious health effects on both adults and children such as cancer and damage to the nervous system have been associated with the consumption of fish contaminated with these heavy metals (Castro-González and Méndez-Armenta, 2008, Vieira et al., 2011). However, these effects are typically seen upon consumption of fish contaminated at much higher levels, than those detected in this study where no heavy metal exceeded MPL. Variations in the concentrations of chemical pollutants in farmed and wild tilapia may be associated with the agricultural cycle in Egypt and the use of fertilizers and pesticides.

4.2. Bacteriological pollutants in farmed tilapia

The mean APC for more than 85% of samples was within the MPL defined by Egyptian standards (10^6 cfu/g) for marketed fresh fish for all samples (EOS, 2005). Pre and post-harvesting contamination and handling of tilapia were identified as potential sources of contamination (Eltholth et al., 2015). Methods of harvesting and handling of tilapia from the farm to retail sale could contribute to the bacterial load of tilapia given the lack of a proper cold chain. This impact will be more during summer season where ambient temperature would be higher than 30 °C. Fish could be contaminated post-capture from fishing tools, contaminated water, contaminated ice, soiled surfaces and boxes, as well as by poor hygienic handling practices (Mhango et al., 2010). Environmental factors such as ambient temperature and relative humidity may play a role in the contamination of tilapia and the level of contamination could vary seasonally. Studies reported that, Gram-positive bacteria such as *Staphylococcus spp.* were increased in sea foods during storage at 2–4°C (Rantsiou et al., 2005, Nimrat et al., 2006). While storing fish at 0°C as soon as possible after capturing will maintain the quality of fish by delaying the spoilage process (Farag, 2012). Microbial pollutants would increase the spoilage of fish and consequently decreasing the shelf life. Bacterial pollutants not only affect the quality of tilapia but also may posing food safety problems.

Very few samples were contaminated with *E. coli*, *L. monocytogenes*, *Salmonella spp.*, *S. aureus* and *V. parahaemolyticus*. Results of the bacteriological analysis were better than that from previous studies (Emire and Gebremariam, 2010, Hamed et al., 2013). *Salmonella spp.* was detected in 10% of seafood samples collected from markets in Alexandria, Egypt (Bakr et al., 2011) which has similar or even lower ambient temperature than the area for the current study. In fish markets with similar environmental conditions the prevalence of *Staphylococcus spp.* in whole and gutted tilapia samples from street vendors in Gaborone, Botswana was 62% and 41%, respectively (Mhango et al., 2010). *Vibrio spp.* was detected in 52% of seafood samples collected from markets in Alexandria, Egypt (Bakr et al., 2011, Kaysner and DePaola Jr, 2000). As *V. parahaemolyticus* in outbreak investigations linked with sea food consumptions. The presence of pathogenic bacteria should be of concern to tilapia handlers, both on retail and households, and consumers but this would depend on the way of processing, cooking and consumption behaviour.

Although, our findings and data from literature indicated that farmed tilapia is safer than wild caught fish, consumers' perceived wild caught fish as safer and pay more for it. This might be due to fears that farmed tilapia are fed on poultry manure and dead poultry. Also some retailers claiming that, they are selling good quality fish from natural resources. The lack of knowledge

and traceability affect consumers' behaviour and consequently has negative impacts on the farmed tilapia value chain (Eltholth et al., 2015). Farmed fish can be of good quality but for value chain to develop need to a) manage hazards particularly post-harvest contamination; b) improve trust; and c) better align hazard perception with reality.

5. CONCLUSIONS

The results of this study show a relatively low level of chemical contaminants but a higher level of biological contaminants in farmed tilapia in the Nile Delta, suggesting the priority is reduction in microbial hazards. This could be supported by promotion of best management practices and use of hazard analysis critical control points (HACCP). Microbiological criteria can be used for assessing the HACCP and other hygienic measures along the farmed tilapia production chain. Further information on consumption amounts and the impact of processing and cooking on hazards is needed in order to estimate the risk to human health: however this initial study suggests farmed tilapia does not present a major or exceptional risk to human health. It also indicated that, consumers' perception in the study area is poorly aligned with reality.

6. ACKNOWLEDGEMENT

This study is an outcome from a project “*Rapid Integrated Assessment of Nutrition and Health Risks Associated with Fish Value Chains in Egypt*” conducted during September 2012 to May 2013. This project was funded by the Australian Centre for International Agricultural Research (ACIAR) and the CGIAR Research Program Agriculture for Nutrition and Health led by the International Food Policy Research Institute. The authors would like to thank the staff members of the Central Diagnostic and Research Laboratory, Faculty of Veterinary Medicine and the Central Laboratory of Environmental Studies Kafrelsheikh University. BH acknowledges financial support from the Leverhulme Centre for Integrative Research on Agriculture and Health (LCIRAH).

7. REFERENCES

- ABBASSY, M., IBRAHIM, H. & ABDEL-KADER, H. 2003. Persistent organochlorine pollutants in the aquatic ecosystem of Lake Manzala, Egypt. *Bull Environ Contam Toxicol*, 70, 1158-1164.
- ABDALLAH, M. A. M. & MORSY, F. A. E. 2013. Persistent organochlorine pollutants and metals residues in sediment and freshwater fish species cultured in a shallow lagoon, Egypt. *Environ Technol*, 34, 2389-2399.
- ABDEL-SATAR, A. & GENEID, Y. A. 2009. Evaluation of heavy metal status in ecosystem of lake Manzalah, Egypt. *Global Journal of Environmental Research*, 3, 194-204.

- ABDEL-SATAR, A. M. & YACOUB, A. 2005. Heavy metals and macro nutrients concentration in *Oreochromis niloticus* and *Tilapia zillii* fish species inhabiting some Egyptian Lakes and El-Salam Canal. *Egyptian Journal of Aquatic Biology and Fisheries*, 9, 97-116.
- ABDO, M. H. 2006. Levels of some trace metals in the water and fish flesh of common species in Abu Za'baal ponds, Egypt. *Egyptian Journal of Aquatic Research*, 32, 13.
- ABDOU, K. A. ASSESSMENT OF SOME CHEMICALS LEVELS AS INDICATORS FOR ENVIRONMENTAL POLLUTION IN WADI EL RAYAN PROTECTED AREA LAKES, FAYOUM, EGYPT. The Fifth Annual International Scientific Conference of Egyptian Society of Environmental Toxicology, 2009.
- ABOU-ARAB, A., GOMAA, M., BADAWEY, A. & NAGUIB, K. 1995. Distribution of organochlorine pesticides in the Egyptian aquatic ecosystem. *Food Chem*, 54, 141-146.
- ABUMOURAD, I. M., AUTHMAN, M. & ABBAS, W. T. 2013. Heavy Metal Pollution and Metallothionein Expression: A Survey on Egyptian Tilapia Farms. *Journal of Applied Sciences Research*, 9.
- AHMED, D. F. I. 2013. Effect of industrial waste discharges including heavy metals in Burullus lake on some physiological parameters and antioxidants in *Tilapia niloticus* and *Siluriformes* fish. *New York Science Journal*, 6, 8.
- AHMED, N. S. & EL-SAAD, E. 2010. Effect of heat treatment on the nutritive value and residues of some synthetic pesticides in fresh Bolti fish. *Arab Universities Journal of Agricultural Sciences*, 18, 329-335.
- ALI, M. H. H. & ABDEL-SATAR, A. M. 2005. Studies of some heavy metals in water, sediment, fish and fish diets in some fish farms in El-Fayoum province, Egypt. *Egypt. J. Aquat. Res.*, 31, 13.
- ALY, O. A. & BADAWEY, M. I. 1984. Organochlorine residues in fish from the River Nile, Egypt. *Bull Environ Contam Toxicol*, 33, 246-252.
- ARAFA, M. & ALI, A. 2008. Effect of some heavy metals pollution in Lake Mariout on *Oreochromis niloticus* fish. *The Egyptian J Comparative Pathol Clinic Pathol*, 21, 191-201.
- AUTHMAN, M. 2008. *Oreochromis niloticus* as a biomonitor of heavy metal pollution with emphasis on potential risk and relation to some biological aspects. *Global Veterinaria*, 2, 104-109.
- AUTHMAN, M., ABBAS, W. T. & GAAFAR, A. Y. 2012. Metals concentrations in Nile tilapia *Oreochromis niloticus* from illegal fish farm in Al-Minufiya Province, Egypt, and their effects on some tissues structures. *Ecotoxicol Environ Saf.*
- AUTHMAN, M. M. 2011. Environmental and experimental studies of aluminium toxicity on the liver of *Oreochromis niloticus* (Linnaeus, 1758) fish. *Life Sci. J*, 8, 764-776.
- AUTHMAN, M. M. & ABBAS, H. H. 2007. Accumulation and Distribution of Copper and Zinc in Both Water and Some Vital Tissues of Two Fish Species (*Tilapia zillii* and *Mugil cephalus*) of Lake Qarun, Fayoum Province, Egypt. *Pakistan Journal of Biological Sciences*, 10, 2106-2122.

- AUTHMAN, M. M., ABBAS, H. H. & ABBAS, W. T. 2013. Assessment of metal status in drainage canal water and their bioaccumulation in *Oreochromis niloticus* fish in relation to human health. *Environ Monit Assess*, 185, 891-907.
- AZAB, M., DARWISH, A., MAHMOUD, H. A. & SDEEK, F. A. 2013. Residue levels of organochlorine pesticides in some ecosystem components of Manzala Lake. *Environ Monit Assess*, 1-12.
- BADAWY, M. I. & EL-DIB, M. A. 1984. Residues of organochlorine pesticides in fish from the Egyptian Delta lakes. *Environ Int*, 10, 3-8.
- BAKR, W. M., HAZZAH, W. A. & ABAZA, A. F. 2011. Detection of *Salmonella* and *Vibrio* species in some seafood in Alexandria. *Journal of American Science*, 7, 663-668.
- CAC. 2009. FAO/WHO Food stds codex alimentarius pesticide residues in food - MRLs & Extraneous MRLs [last updated March]; 2009 [Online]. Available: http://www.codexalimentarius.net/mrls/pestdes/jsp/pest_q-e.jsp.
- CASTRO-GONZÁLEZ, M. & MÉNDEZ-ARMENTA, M. 2008. Heavy metals: Implications associated to fish consumption. *Environ Toxicol Pharmacol*, 26, 263-271.
- CHEESBROUGH, M. 1987. Medical laboratory manual for tropical countries. Vol. I. Medical laboratory manual for tropical countries. Vol. I. 2nd edition.
- DOMINGO, J. L. 2007. Omega-3 fatty acids and the benefits of fish consumption: is all that glitters gold? *Environ Int*, 33, 993-998.
- EL-BARBARI, M. & SAMI, E. Determination of heavy metal residues in marine and freshwater fishes from markets in Sharkia Governorate. Proceedings of the 5th Global Fisheries and Aquaculture Research Conference, Faculty of Agriculture, Cairo University, Giza, Egypt, 1-3 October 2012., 2012. Massive Conferences and Trade Fairs, 199-208.
- EL-KATTAN, Y. & NAHLA, A. 2008. Levels of some heavy metals in River Nile water and *Oreochromis niloticus* fish at Menoufia Governorate. *Egypt. J. Comp. Path. & Clinic. Path.*, 21, 12.
- EL-MEKKAWI, H., DIAB, M., ZAKI, M. & HASSAN, A. 2009. Determination of Chlorinated Organic Pesticide Residues in Water, Sediments, and Fish from Private Fish Farms at Abbassa and Sahl Al-Husainia, Shakia Governorate. *Australian Journal of Basic and Applied Sciences*, 3, 4376-4383.
- EL-NEMAKI, F. A., ALI, N. A., ZEINHOM, M. M. & RADWAN, O. A. Impacts of different water resources on the ecological parameters and quality of tilapia production at El-Abbassa fish farms in EGYPT. 8th International Symposium on Tilapia in Aquaculture, 2008.
- EL-SAYED, E.-S. A., EL-AYYAT, M. S., NASR, E.-S. & KHATER, Z. Z. K. 2011. Assessment of heavy metals in water, sediment and fish tissues, from Sharkia province, Egypt El-Sayed A. El-Sayed, Mohamed S. El-Ayyat 2, El-Sayed Nasr and. *Egypt. J. Aquat. Biol. & Fish.*, 15, 19.
- ELGHOBASHY, H. A., ZAGHLOUL, K. H. & METWALLY, M. A. 2001. Effect of some water pollutants on the Nile tilapia, *Oreochromis niloticus* collected from the River Nile and some Egyptian lakes. *Egyptian Journal of Aquatic Biology and Fisheries*, 5, 251-279.

- ELTHOLTH, M., FORNACE, K., GRACE, D., RUSHTON, J. & HÄSLER, B. 2015. Characterisation of production, marketing and consumption patterns of farmed tilapia in the Nile Delta of Egypt. *Food Policy*, 51, 131-143.
- EMIRE, S. A. & GEBREMARIAM, M. M. 2010. INFLUENCE OF FROZEN PERIOD ON THE PROXIMATE COMPOSITION AND MICROBIOLOGICAL QUALITY OF NILE TILAPIA FISH (*OREOCHROMIS NILOTICUS*). *Journal of Food Processing and Preservation*, 34, 743-757.
- EOS 2005. Egyptian Organization for Standardization and Quality, Chilled fish. ES No. 3494.
- EOS 2010. Egyptian Organization for Standardization and Quality, Maximum level for heavy metal contaminants in food. ES No. 7136.
- ESSA, H. H. & RATEB, H. Z. 2011. RESIDUES OF SOME HEAVY METALS IN FRESHWATER FISH (*OREOCHROMIS NILOTICUS* AND *LABEO NILOTICUS*) IN ASSIUT CITY MARKETS. *Ass. Univ. Bull. Environ. Res.*, 14, 9.
- FAO 2013. FAO Fisheries and Aquaculture Department has published the Global Aquaculture Production Statistics for the year 2011. 2013 ed.:
<ftp://ftp.fao.org/FI/news/GlobalAquacultureProductionStatistics2011.pdf>.
- FARAG, H. 2012. Sensory and chemical changes associated with microbial flora of *Oreochromis niloticus* stored in ice. *International Food Research Journal*, 19.
- FENG, P., WEAGANT, S. D. & GRANT, M. A. 2002. Enumeration of *Escherichia coli* and the coliform bacteria. *Bacteriological Analytical Manual*, 8, 102-135.
- FISHAR, M. & ALI, M. 2005. Accumulation of trace metals in some benthic invertebrate and fish species relevant to their concentration in water and sediment of lake qarun, Egypt.
- GABR, H. R. & GAB-ALLA, A.-A. 2007. Comparison of Biochemical Composition and Organoleptic Properties Between Wild and Cultured Finfish. *Journal of Fisheries & Aquatic Science*, 2.
- GAD, N. S. & TOUFEEK, M. A. 2010. Distribution and accumulation of some trace metals in water and fish from Damietta Branch of River Nile. *African J. Biol. Sci.*, 6, 21.
- HAMED, Y., ABDELMONEIM, T., ELKIKI, M., HASSAN, M. & BERNDTSSON, R. 2013. Assessment of Heavy Metals Pollution and Microbial Contamination in Water, Sediments and Fish of Lake Manzala, Egypt. *Life Science Journal*, 10.
- HASSANIN, S. I. METALS RESIDUES, HISTOLOGICAL ALTERATIONS AND COOKING METHODS OF FISH cultured in WASTEWATER PONDS. From the pharaohs to the future. Eighth International Symposium on Tilapia in Aquaculture. Proceedings. Cairo, Egypt, 12-14 October, 2008., 2008. AQUAFISH Collaborative Research Support Program, 569-588.
- HASTEIN, T., HJELTNES, B., LILLEHAUG, A., UTNE SKARE, J., BERNTSEN, M. & LUNDEBYE, A. 2006. Food safety hazards that occur during the production stage: challenges for fish farming and the fishing industry. *Rev Sci Tech*, 25, 607-25.

- HITCHINS, A. D. & JINNEMAN, K. 2003. Detection and enumeration of *Listeria monocytogenes* in foods. Bacteriological analytical manual (BAM) chapter,[online].
- HOLT, J. G., KRIEG, N. R., SNEATH, P., STALEY, J. & WILLIAMS, S. 1994. Bergey's Manual of Determinative Bacteriology. 9th edition. William and Wilkins Baltimore Maryland, USA, 527-533.
- JÄRUP, L. 2003. Hazards of heavy metal contamination. British medical bulletin, 68, 167-182.
- JONES JR, J. B., WOLF, B. & MILLS, H. A. 1991. Plant analysis handbook. A practical sampling, preparation, analysis, and interpretation guide, Micro-Macro Publishing, Inc.
- KAUD, H. & EL-DAHSHAN, A. 2010. Bioaccumulation and histopathological alterations of the heavy metals in *Oreochromis niloticus* fish. Nature and science, 8, 147-156.
- KAYSNER, C. & DEPAOLA JR, A. 2000. Outbreaks of *Vibrio parahaemolyticus* gastroenteritis from raw oyster consumption: assessing the risk of consumption and genetic methods for detection of pathogenic strains.
- KHALLAF, E., GALAL, M. & AUTHMAN, M. 1998. Assessment of heavy metals pollution and their effects on *Oreochromis niloticus* in aquatic drainage water. Journal of the Egyptian German Society of Zoology, 26, 39-74.
- KHORSHED, M. Survey of some trace elements in some marketable Egyptian edible fish species. Proceedings of the 2nd Global Fisheries and Aquaculture Research Conference, Cairo International Convention Center, 24-26 October 2009., 2009. Massive Conferences and Trade Fairs, 97-112.
- MACFADYEN, G., ALLAH, A., KENAWY, D., AHMED, M., HEBICHA, H., DIAB, A., HUSSEIN, S., ABOUZIED, R. & EL NAGGAR, G. 2011. Value-chain analysis of Egyptian aquaculture. Penang, Malaysia: The WorldFish Center.
- MACFADYEN, G., NASR-ALLA, A. M., AL-KENAWY, D., FATHI, M., HEBICHA, H., DIAB, A. M., HUSSEIN, S. M., ABOU-ZEID, R. M. & EL-NAGGAR, G. 2012. Value-Chain Analysis—an assessment methodology to estimate Egyptian aquaculture sector performance. Aquaculture.
- MALHAT, F. 2011. Distribution of heavy metal residues in fish from the River Nile tributaries in Egypt. Bull Environ Contam Toxicol, 87, 163-165.
- MANSOUR, S. & SIDKY, M. 2002. Ecotoxicological studies. 3. Heavy metals contaminating water and fish from Fayoum Governorate, Egypt. Food Chem, 78, 15-22.
- MANSOUR, S. & SIDKY, M. 2003. Ecotoxicological Studies. 6. The first comparative study between Lake Qarun and Wadi El-Rayan wetland (Egypt), with respect to contamination of their major components. Food Chem, 82, 181-189.
- MATURIN, L. & PEELER, J. 2001. Aerobic plate count, chap. 3. Bacteriological analytical manual, 8th ed. AOAC International, Arlington, VA.
- MHANGO, M., MPUCHANE, S. & MPUCHANE, B. 2010. Incidence of indicator organisms, opportunistic and pathogenic bacteria in fish. African Journal of Food, Agriculture, Nutrition and Development, 10.

- MOHAMED, F. A. 2008. Bioaccumulation of selected metals and histopathological alterations in tissues of *Oreochromis niloticus* and *Lates niloticus* from Lake Nasser, Egypt. *Global Veterinaria*, 2, 205-218.
- NASR, N., ARIEF, M., ABDEL-ALEEM, A. & MALHAT, F. 2009. Persistent Organic Pollutants (POPs) in Egyptian Aquatic Environment. *J. Appl. Sci. Res*, 5, 1929-1940.
- NIMRAT, S., SIRIBOONLAMOM, S., ZHANG, S., XU, Y. & VUTHIPHANDCHAI, V. 2006. Chilled storage of white shrimp (*Litopenaeus vannamei*) spermatophores. *Aquaculture*, 261, 944-951.
- OMAR, W. A., ZAGHLOUL, K. H., ABDEL-KHALEK, A. A. & ABO-HEGAB, S. 2013. Risk Assessment and Toxic Effects of Metal Pollution in Two Cultured and Wild Fish Species from Highly Degraded Aquatic Habitats. *Arch Environ Contam Toxicol*, 65, 753-764.
- OSMAN, A. G. 2012. Biomarkers in Nile Tilapia *Oreochromis niloticus niloticus* (Linnaeus, 1758) to Assess the Impacts of River Nile Pollution: Bioaccumulation, Biochemical and Tissues Biomarkers. *Journal of Environmental Protection*, 3.
- RANTSIOU, K., IACUMIN, L., CANTONI, C., COMI, G. & COCOLIN, L. 2005. Ecology and characterization by molecular methods of *Staphylococcus* species isolated from fresh sausages. *International journal of food microbiology*, 97, 277-284.
- RASHED, M. 2001. Cadmium and lead levels in fish (*Tilapia nilotica*) tissues as biological indicator for lake water pollution. *Environ Monit Assess*, 68, 75-89.
- SAEED, S. M. 2013. Impact of environmental parameters on fish condition and quality in Lake Edku, Egypt. *Egypt. J. Aquat. Biol. & Fish*, 17, 12.
- SAEED, S. M., SHAKER, I. & M, R. Assessment of heavy metals pollution in water and sediments and their effect on *Oreochromis niloticus* in the Northern Delta Lakes, Egypt. 8th International Symposium on Tilapia in Aquaculture, 2008. 475-490.
- SALIM, A., HASSANIN, M. & ZOHAIR, A. 2003. A simple procedure for reducing lead content in fish. *Food and chemical toxicology*, 41, 595-597.
- SANKAR, G., SARAVANAN, P., KRISHNAMURTHY, P., CHANDRAKALA, N. & RAJENDRAN, K. 2012. Isolation and identification of *Vibrio* spp. in diseased *Channa punctatus* from aquaculture fish farm. *IJMS*, 41, 2.
- SASSD, S. Heavy metals accumulation in common fish species inhabiting Lake Edku and their relation to lipid content and liver size. *Proceedings of the 4th Global Fisheries and Aquaculture Research Conference, the Egyptian International Center for Agriculture, Giza, Egypt, 3-5 October 2011.*, 2011. *Massive Conferences and Trade Fairs*, 139-150.
- SHEREIF, M. & MANCY, K. 1995. Organochlorine pesticides and heavy metals in fish reared in treated sewage effluents and fish grown in farms using polluted surface waters in Egypt. *Water Science and Technology*, 32, 153-161.

- Smallwood, D.M., and J.R. Blaylock. 1991. "Food Safety and the Demand for Food: Methodological Models and Applications," in *Economics of Food Safety*, J.A. Caswell, ed., pp. 3-27. Amsterdam, The Netherlands: Elsevier Science Publishing Co., Inc.
- SUBASINGHE, R., SOTO, D. & JIA, J. 2009. Global aquaculture and its role in sustainable development. *Reviews in Aquaculture*, 1, 2-9.
- TROCINO, A., XICCATO, G., MAJOLINI, D., TAZZOLI, M., TULLI, F., TIBALDI, E., MESSINA, C. M. & SANTULLI, A. 2012. Levels of dioxin-like polychlorinated biphenyls (DL-PCBs) and metals in European sea bass from fish farms in Italy. *Food Chem*, 134, 333-338.
- VANDERZANT, C. & SPLITTSTOESSER, D. 1992. *Compendium of methods for the Microbiological examination of foods*, 3rd edn. eds. Washington DC: American Public Health Association. ISBN 0-87553173-3.
- VIEIRA, C., MORAIS, S., RAMOS, S., DELERUE-MATOS, C. & OLIVEIRA, M. 2011. Mercury, cadmium, lead and arsenic levels in three pelagic fish species from the Atlantic Ocean: intra-and inter-specific variability and human health risks for consumption. *Food and chemical toxicology*, 49, 923-932.
- YACOUB, A. M. & GAD, N. S. 2012. Accumulation of some heavy metals and biochemical alterations in muscles of *Oreochromis niloticus* from the River Nile in Upper Egypt. *INTERNATIONAL JOURNAL OF ENVIRONMENTAL SCIENCE AND ENGINEERING (IJESE)*, 3, 1-10.
- YAHIA, D. & ELSHARKAWY, E. E. 2014. Multi pesticide and PCB residues in Nile tilapia and catfish in Assiut city, Egypt. *Science of The Total Environment*, 466, 306-314.
- YAMASHITA, N., URUSHIGAWA, Y., MASUNAGA, S., WALASH, M. I. & MIYAZAKI, A. 2000. Organochlorine pesticides in water, sediment and fish from the Nile River and Manzala Lake in Egypt. *International Journal of Environmental Analytical Chemistry*, 77, 289-303.
- YEHIA, H. M. & SEBAEE, E. S. 2012. Bioaccumulation of heavy metals in water, sediment and fish (*Oreochromis niloticus* and *Clarias anguillaris*), in Rosetta branch of the River Nile, Egypt. *African Journal of Biotechnology*, 11, 14204-14216.
- ZAGHLOUL, K. H., HASHEESH, W. S., ZAHRAN, I. A. & MARIE, M.-A. S. 2007. Ecological and biological studies on the Nile tilapia *Oreochromis niloticus* along different sites of lake Burullus Egypt. *J. Aquat. Biol. & Fish.*, 11- 31.

Appendix 1 laboratory analysis

1. Chemical analysis

At the Kafrelsheikh University Central Laboratory of Environmental Studies (KUCLES), after cleaning the tilapia and removal of scales, two separate tissue samples of 10 g each were taken from each fish from the dorsal muscles. One sample (10 g) was for the detection of heavy metals and the other for pesticide residues. Samples were immediately frozen at -20°C until analysis. The following protocols were used for analysis: For the detection of organophosphorus (OP) and organochloride (OC) residues a protocol as described by Darko et al. (2008) was adopted. Each sample was ground in a blender to obtain a homogenous composite. Pesticide residues were extracted using liquid extraction with 150 mL of acetone and n-hexane (20:80 v/v). The tissue sample was placed into a beaker containing 50 mL from the solvent mixture and was shaken for 20 min, followed by filtration. This process was repeated twice, each time with 50 mL of the solvent mixture. The collected filtrate then was passed through anhydrous sodium sulfate. The extracts were concentrated to 1 mL using vacuum rotary evaporator. The extract for each sample was then dissolved in hexane (10 mL) and passed through pre-conditioned octadecyl C-18 columns at a rate of 2 ml/min to clean up. The column was washed with 30% methanol (1 mL) followed by ultrapure water (1 mL) and then allowed to dry. The trapped sample (analyte) in the column was eluted five times with 0.5 mL aliquots of hexane to recover the pesticides residues. Hexane in the sample was then evaporated off leaving the residues alone in the vial. The dried sample was dissolved in hexane (1 mL), thoroughly mixed and then transferred to auto sampler vials ready for gas chromatography. The OC and OP residues were analyzed using Gas chromatography–mass spectrometry (GC-MS, THERMO SCIENTIFIC). All chemicals used were HPLC grade.

For the detection of arsenic, lead and cadmium, fish samples were dried at 65°C for 48 h in a desiccator. One gram of each sample was digested using dry ashing method in a muffle furnace at 450° C for 5h, then the ash was extracted using 20% hydrochloric acid as described in detail elsewhere (Jones Jr et al., 1991). All heavy metals were measured by atomic absorption spectrometry (Unicam 969, Analytical Technology Inc., Cambridge, United Kingdom).

Mercury was detected directly from tilapia tissue samples using NIC Mercury Analyzer MA-3000. The following standard solutions were prepared and used: 1) L-cysteine 100mg/L: 100 mg of L-cysteine HSC₂CH (NH₂) COOH was added to a 1000 ml flask with water and 2 ml of nitric acid was then added to make a total of 1000 ml of solution. The flask was then shaken to stir the solution and stored in a cool dark place, 2) Mercury standard solution 10 mg/L: 1 ml

of mercury standard solution for atomic absorption (1000 ppm) was to add L-cysteine solution (100 mg/L) to make a total of 100 ml and 3) Mercury standard solution 0.1mg/L: L-cysteine (100 mg/L) was added to 1ml of the previous prepared solution to make a total of 100 ml. The manufacturer's protocol was followed for preparing the standard solutions and the measurement methods.

2. Bacteriological analysis

Samples were prepared for bacterial isolation and enumeration by cutting out, with sterile instruments, 25 g of dorsal muscles. Samples were then homogenized in 225 ml of sterile buffered peptone water. The homogenate samples were inoculated on standard plate count media and media for the selective isolation of bacteria. The isolated colonies were subjected to bacteriological characterization procedures. First, selective media were used for the identification of potential isolated colonies, while further identification was conducted by using morphological and biochemical tests. Aerobic plate count, *Salmonella* spp., *E. coli*, *S. aureus*, *L. monocytogenes*, and *V. parahaemolyticus* tests were performed as described below. All media used were from Oxoid Ltd, UK. For the APC, homogenised samples were serially diluted and plated using the pour-overlay method on solid medium using standard plate count agar. Petri dishes were then incubated in inverted position for 48 ± 2 h at 37°C. (Maturin and Peeler, 2001) Presumptive *E. coli* was enumerated on eosin methylene blue (EMB) agar. (Feng et al., 2002) After incubation, typical colonies with metallic sheen were counted after which, 3-5 colonies were selected for biochemical confirmation. *Listeria monocytogenes* was enriched on Fraser Broth at 35°C for up to 48 h and then inoculated on PALCAM *Listeria* agar at 37°C for up to 48 h. (Hitchins and Jinneman, 2003) *Salmonella* spp. were detected by a pre-enrichment with Rappaport-Vassiliadis peptone broth at 42°C for 24 h, then plating on xylose lysine deoxycholate (XLD) agar at 42°C for 24 h (ISO 6579, 2002). For the detection of *V. parahaemolyticus*, alkaline peptone water (APW) was used as a diluent and thiosulphate citrate bile salt (TCBS) agar 3% for 24 h at 37°C as selective media (Sankar et al., 2012). Baird Parker agar (BPA) was used for the detection and enumeration of *S. aureus*, incubated at 37°C for 24–48 hours (Vanderzant and Splittstoesser, 1992). For biochemical characterisation of isolates, the following biochemical tests were used: catalase, oxidase, sugar fermenter, indol production, methyl red, Voges-Proskauer, Simmon's citrate, urease, blood agar and TSI (Cheesbrough, 1987, Holt et al., 1994).

Appendix 2: The concentrations of pesticide residues in tilapia muscle tissues from different sources in Egypt (ND= not detected, NM= not measured)

Source of tilapia	Pesticide residues (Mean concentration)							Reference
	Unit	Aldrin	Dieldrin	Endrin	Heptachlor	Heptachlor-Epoxide (beta)	Lindane	
Fish markets (Fresh)	ng/g	ND	1.69	1.89	1.49	ND	0.58	(Ahmed and El-Saad, 2010)
Fish markets in (Grilled)	ng/g	ND	0.66	0.71	0.85	ND	ND	
Assiut_Elwasta	µg/kg	0.61	ND	NM	NM	NM	0.89	(Yahia and Elsharkawy,
Assiut_Mankbad	µg/kg	0.22	ND	NM	0.18	NM	0.47	2014)
Lake Manzala	ng/g	NM	NM	NM	NM	NM	0.91	(Abbassy et al., 2003)
Lake Manzala	µg/kg	0.033	NM	0.026	0.003	0.059	NM	(Azab et al., 2013)
Lake Manzala	µg/kg	8.6	NM	NM	1.35	NM	2.508	(Abou-Arab et al., 1995)
River Nile	µg/kg	3.85	NM	NM	2.63	NM	2.2	
River Nile	µg/kg	ND	ND	23.94	ND	ND	8.93	(Aly and Badawy, 1984)
Fish Farms	ng/g	ND	ND	NM	NM	NM	1.054	(El-Mekkawi et al., 2009)
Fish Farms	ng/g	13.038	10.125	NM	NM	NM	9.5	
Lake Manzala	ng/g	ND	ND	ND	0.14	ND	1.49	(Yamashita et al., 2000)
River Nile	ng/g	ND	ND	ND	0.10	ND	0.18	
Lake Idku	ng/g	0.023	0.130	0.028	NM	NM	0.025	(Abdallah and Morsy, 2013)
Lake Idku	µg/kg	ND	ND	<0.15	ND	ND	4.83	(Badawy and El-Dib, 1984)
Lake Maryut	µg/kg	ND	ND	<0.15	ND	ND	4.53	
Fresh water canals	ng/g	0.802	1.423	1.47	0.59	0.639	0.914	(Nasr et al., 2009)
Drain water canals	ng/g	0.614	1.597	1.75	2.576	1.786	2.01	
Lake Qarun	mg/kg	0.004	ND	ND	ND	ND	0.008	(Mansour and Sidky, 2003)

Appendix 3: The concentrations of heavy metals in tilapia muscle tissues from different sources in Egypt (ND= not detected, NM= not measured)

Source of tilapia	Heavy metals (Mean concentration)				Reference
	Unit	Cadmium	Lead	Mercury	
Illegal fish farm in Sabal drainage canal	mg/kg	4.35	3.02	2.35	(Authman et al., 2012)
Fish farms	mg/kg	1.21	1.52	3.50	(Kaoud and El-Dahshan, 2010)
Fish farms in Kafrelsheikh	µg/kg	2.33	6.27	NM	(Abumourad et al., 2013)
Fish farms in AL-Abasa	µg/kg	2.27	18.8	NM	
Fish farms in EL-Fayoum	µg/kg	4.27	8.63	NM	
Suez fish farm	mg/kg	NM	1.65	NM	(Shereif and Mancy, 1995)
Fish farm in Lake Manzala	mg/kg	NM	11.25	NM	
Fish ponds with water supply from Bahr El-Baquar drain	mg/kg	0.317	0.42	0.02	(HASSANIN, 2008)
Shanawan drainage canal, Al-Minufiya	mg/kg	5.30	48.70	67.1	(Khallaf et al., 1998)
Private fish farm at Fayoum Governorate	mg/kg	0.10	6.38	NM	(Mansour and Sidky, 2002) 1997/1998
Lake Qarun	mg/kg	0.16	1.65	NM	
Private fish farm at Fayoum Governorate	mg/kg	0.01	0.01	NM	(Mansour and Sidky, 2002) 1998/1999
Lake Qarun	mg/kg	0.08	0.06	NM	
El-Shoura fish farm in El-Fayoum Governorate	mg/kg	2.20	6.50	NM	(Ali and Abdel-Satar, 2005)
Sabal drainage canal, Al-Minufiya Province	mg/kg	3.40	31.95	39.13	(Authman, 2008)
Lake Edku	µg/g	ND	0.24	NM	(Abdallah and Morsy, 2013)
Lake Edku	µg/g	0.19	0.59	NM	(Saeed et al., 2008)
Lake Borollus	µg/g	0.01	0.05	NM	
Lake Manzala	µg/g	10.36	10.1	NM	
Lake Edku	µg/g	0.21	0.68	NM	(Sassd, 2011)
Lake Edku	µg/g	0.27	0.52	NM	(Saeed, 2013)
Lake Brullus	mg/kg	NM	0.30	NM	(Zaghloul et al., 2007)
Lake Brullus	mg/kg	ND	10.75	0.13	(Ahmed, 2013)

Bahar tera	mg/kg	ND	9.60	0.03	
River Nile at Monofya governorate	mg/kg	0.22	1.03	0.33	(El-Kattan and Nahla, 2008)
Ismalia canal	mg/kg	0.01	0.19	NM	(Elghobashy et al., 2001)
River Nile (Shubra El-Khcima, Cairo sector)	mg/kg	0.04	1.22	NM	
Maazala Lake	mg/kg	0.35	3.2	NM	
Lake Edku	mg/kg	0.11	1.57	NM	
Lake Borollus	mg/kg	0.03	0.22	NM	
Lake Qarun	mg/kg	0.01	0.33	NM	
Lake Maryut	mg/kg	0.53	1.12	NM	
Lake Maryut	mg/kg	0.24	2.77	NM	(Arafa and Ali, 2008)
Fish market survey	mg/kg	0.01	0.06	0.02	(Khorshed, 2009)
River Nile/Upper Egypt	mg/kg	NM	14.16	NM	(Yacoub and Gad, 2012)
River Nile	mg/kg	0.79	5.82	0.01	(Osman, 2012)
Fish markets survey	µg/kg	0.98	0.13	4.88	(Essa and Rateb, 2011)
Sabal draiage ncanal (Al-Menoufiya Province)	mg/kg	10.85	10.21	8.66	(Authman et al., 2013)
Lake Manzalah	µg/g	1.33	5.36	NM	(Abdel-Satar and Geneid, 2009)
Lake Manzalah	mg/kg	0.64	3.37	NM	(Abdel-Satar and Yacoub, 2005)
Elsalam canal	mg/kg	0.66	3.48	NM	
Lake Edku	mg/kg	0.27	0.52	NM	(Saeed, 2013)
Abu Za'baal ponds	µg/g	3.60	21.30	NM	(Abdo, 2006)
Lake Nasser	µg/g	1.25	1.75	NM	(Mohamed, 2008)
Sharkia province	µg/g	0.81	3.34	NM	(El-Sayed et al., 2011)
Lake Manzala	µg/g	0.11	3.81	NM	(Hamed et al., 2013)
High Dam Lake, Aswan	mg/kg	0.02	0.13	NM	(Rashed, 2001)
Lake Qarun	µg/g	1.23	6.72	NM	(Fishar and Ali, 2005)
Lake Qarun	mg/kg	0.06	0.09	NM	(Mansour and Sidky, 2003)
Lake Qarun	mg/kg	NM	3.76	NM	(Omar et al., 2013)
Qarun fish farms	mg/kg	NM	2.38	NM	(Omar et al., 2013)
Wadi El-Rayan protected area lakes	mg/kg	0.11	0.157	0.01	(Abdou, 2009)

Farmed tilapia	mg/kg	NM	2.41	NM	(Gabr and Gab-Alla, 2007)
Wild tilapia	mg/kg	NM	2.32	NM	(Gabr and Gab-Alla, 2007)
Ponds with irrigation water	µg/g	0.13	1.54	NM	(El-Nemaki et al., 2008)
Ponds with agriculture drainage water	µg/g	0.04	0.90	NM	(El-Nemaki et al., 2008)
River Nile tributaries	µg/g	0.76	1.51	NM	(Malhat, 2011)
River Nile-Rosetta branch	mg/kg	1.51	5.94	NM	(Yehia and Sebaee, 2012)
River Nile-Damietta Branch	mg/kg	NM	14.65	NM	(Gad and Toufeek, 2010)
Fish markets/Sharkiah governorate	mg/kg	0.02	0.15	0.37	(El-Barbari and Sami, 2012)
Fish markets	mg/kg	NM	0.52	NM	(Salim et al., 2003).
