

This is the author's accepted manuscript of the following article:

Gostelow, R., Hazuchova, K., Scudder, C., Forcada, Y., Church, D. and Niessen, S. J. M. (2017)
'Prospective evaluation of a protocol for transitioning porcine lente insulin-treated diabetic cats to human recombinant protamine zinc insulin', *Journal of Feline Medicine and Surgery*.

The final publication is available at SAGE Journals via
<https://doi.org/10.1177/1098612X17697482>.

The full details of the published version of the article are as follows:

TITLE: Prospective evaluation of a protocol for transitioning porcine lente insulin-treated diabetic cats to human recombinant protamine zinc insulin

AUTHORS: Ruth Gostelow, Katarina Hazuchova, Christopher Scudder, Yaiza Forcada, David Church, Stijn J M Niessen

JOURNAL TITLE: Journal of Feline Medicine and Surgery

PUBLICATION DATE: March 15, 2017 (online)

PUBLISHER: SAGE Publications

DOI: 10.1177/1098612X17697482

Title: Prospective evaluation of a protocol for transitioning porcine lente insulin-treated diabetic cats to human recombinant protamine zinc insulin

Authors: Gostelow, R.,^a Hazuchova, K.,^a Scudder, C.,^a Forcada, Y.,^a Church, D.,^a Niessen, SJM.^{a,b}

Institutions and Affiliations: ^a Diabetic Remission Clinic, Department of Clinical Science and Services, The Royal Veterinary College, London, UK. ^b Newcastle Medical School, Newcastle upon Tyne, Tyne and Wear, UK.

Corresponding Author: R. Gostelow BVetMed(Hons) MVetMed DipACVIM DipECVIM-ca MRCVS. Email: rgostelow@rvc.ac.uk. Department of Clinical Science and Services, The Royal Veterinary College, Hawkshead Lane, North Mymms, Hertfordshire, AL9 7TA, UK. Telephone: 01707 666 366.

Keywords: Diabetes mellitus, feline, insulin, remission

Abstract

Objectives: Evaluate a nadir-led protocol for transitioning porcine lente insulin suspension (PLIS)-treated diabetic cats onto human recombinant protamine zinc insulin (PZIR).

Methods: Recently-diagnosed (<5 months) diabetic cats, treated with twice-daily (BID) PLIS for ≥ 6 weeks, were recruited. Fructosamine, 24-hour blood glucose curve (BGC), quality of life assessment (DIAQoL-pet score), and Diabetic Clinical Score (DCS) were assessed on enrolment (during PLIS treatment), and 2, 4 and 12 weeks after transitioning to PZIR (starting dose 0.2-0.7U/kg BID). Short duration of insulin-action was defined as <9 hours. Linear mixed effects modelling assessed for change in fructosamine, mean blood glucose (MBG) during BGCs, DIAQoL-pet score, DCS, and BID insulin dose. McNemar's tests compared the proportion of cats with hypoglycaemia at week 0 (PLIS-treated) and week 4 (PZIR-treated).

Results: Twenty-two cats were recruited. Median PLIS dose at enrolment was 0.5U/kg (interquartile range (IQR) 0.3–0.7U/kg) BID, equalling median PZIR starting dose (0.5, IQR 0.3-0.7U/kg BID). Transitioning was followed by significant decreases in fructosamine ($p=0.00007$), insulin dose ($p=0.02$), DCS ($p=8.1 \times 10^{-8}$) and DIAQoL-pet score ($p=0.003$), indicating improved quality of life. MBG did not alter significantly ($p=0.1$). Five cats (22.7%) achieved diabetic remission. Hypoglycaemia was recorded in 30 of 190 12-hour BGCs (15.8%) and five cats experienced clinical hypoglycaemia. Proportion of cats with hypoglycaemia did not differ between PLIS (week 0) and PZIR (week 4) ($p=1.0$). Duration of action was analysed in 19 cats. Six cats (31.6%) showed short duration of action on PLIS, compared to 2 cats (10.5%) after 4 weeks on PZIR. All six cats with short PLIS duration showed duration of ≥ 9 hours on PZIR.

Conclusions and Relevance: All cats transitioned successfully. Transitioning was associated with significantly improved clinical signs and quality of life, with some cats achieving remission. Transition to PZIR should be considered for cats with short duration of action on PLIS.

Introduction

The major goals in managing feline diabetes mellitus (DM) are to reduce or eliminate diabetic clinical signs while avoiding hypoglycaemia and other complications.¹ Treatment can also result in diabetic remission, which can substantially improve the quality of life of both cats and their owners.² Subcutaneous insulin q12h and a low-carbohydrate, high-protein diet are the mainstays of treatment.^{3,4} Although several different insulin types have been used for feline DM, a number of recent guidelines and a systematic review on diabetic remission recommend the use of long-acting insulin preparations when treating diabetic cats.³⁻⁵ Long-acting insulins include protamine zinc insulin (PZI) and the recombinant human insulin analogues glargine and detemir, and have been associated with a longer duration of action than porcine lente insulin suspension (PLIS),⁶ which has an intermediate duration of action. This longer duration of action could help eliminate diabetic clinical signs, and could also increase a cat's chance of diabetic remission because early, effective glycaemic control might reduce the deleterious effects of glucotoxicity on pancreatic beta-cells.^{5,7,8} However, until recently, PLIS (Caninsulin®, MSD Animal Health) had been the only veterinary-licensed insulin in the United Kingdom and Europe.

The licensing of a human recombinant protamine zinc insulin solution (PZIR) (ProZinc®, Boehringer Ingelheim) has recently changed the above situation, and information on transitioning cats from PLIS onto PZIR is therefore necessary. A previous study examining the efficacy of PZIR in diabetic cats included a small proportion of cats (n=13) that were previously treated with other insulin types (mainly ultralente and neutral protamine Hagedorn).⁹ However, no previous studies have specifically assessed the progress of PLIS-treated cats following transition to PZIR, or evaluated how best to conduct the transition. The aim of this study was therefore to prospectively evaluate the efficacy of a standardised nadir-based protocol to transition PLIS-treated cats onto PZIR using the manufacturer-recommended PZIR starting dose, and to document the effects in individual cats following this transition.

Materials and Methods

This was a prospective, uncontrolled cohort study conducted at the Royal Veterinary College (RVC) Diabetic Remission Clinic (DRC) between December 2013 and December 2015. Ethical approval was granted by the RVC Ethics and Welfare Board, and the Veterinary Medicines Directorate (Animal Test Certificate Number ATC-S-036). To be eligible, cats must have been diagnosed with DM in the last five months. This reasonably short duration of DM was chosen to increase the likelihood that some cats might achieve diabetic remission following transitioning because remission is reported to be

52 more likely early in the course of a cat's DM.⁸ Diagnosis of DM was based on a history of compatible clinical signs (polyuria,
53 polydipsia, polyphagia) and persistent hyperglycaemia (> 15 mmol/l) at the time of their first trial appointment. These
54 criteria were similar to those used in a previous study examining the effect of PZIR therapy in diabetic cats.⁹ Recruited
55 cats must have been receiving BID PLIS (Caninsulin®, MSD Animal Health) for a minimum of 6 weeks prior to their initial
56 trial visit. This minimum duration of PLIS treatment aimed to allow cats to achieve stable glycaemic control using PLIS by
57 the time of their trial recruitment visit. This initial treatment period also allowed cats with concurrent disease or
58 suspected insulin resistance to be identified prior to their first trial visit. All cats were transitioned onto a high-protein,
59 low-carbohydrate diet a minimum of 10 days before their first trial appointment.

60

61 Final eligibility was based on the results of underlying disease screening at the initial visit. This involved clinical history
62 and physical examination by residency-trained or board-certified internal medicine specialists (RG, CS, KH and SN). Cats
63 then underwent complete blood cell count, serum biochemistry, serum total thyroxine (TT4), serum fructosamine,
64 insulin-like growth factor-1 (IGF-1) and feline pancreatic lipase immunoreactivity (fPLI) measurement using assays
65 previously validated for cats. Cystocentesis samples were submitted for urinalysis and bacterial culture, and cats also
66 received abdominal ultrasonography and echocardiography by a residency-trained specialist.

67

68 Each cat also underwent a 24-hour blood glucose curve (BGC) to assess glycaemic control on their current PLIS regime.
69 Curves were conducted using the Guardian® REAL-Time Continuous Glucose Monitoring System (CGMS) (Medtronic,
70 Watford, UK), starting at least 12 hours after all laboratory tests and diagnostic imaging were complete. Curves were
71 completed using 2-hourly readings with a veterinary handheld glucometer (AlphaTRAK®2, Zoetis) if the CGMS failed
72 during the 24-hour monitoring period. Each cat's clinical signs at recruitment were graded using a validated clinical
73 scoring system (Diabetic Clinical Score) (Table 1).¹⁰ The Diabetic Clinical Score (DCS) scores major clinical signs of poor
74 diabetic control from 0 (absent) to 3 (severe) to generate a final score out of 12. Finally, cat and owner quality of life was
75 estimated through owners completing the validated DIAQoL-pet quality of life questionnaire for diabetic cats.¹¹ The
76 DIAQoL-pet generates an Average-Weighted Impact Score (AWIS) to reflect pet and owner quality of life, with more
77 negative values reflecting a more negative impact of DM. Cats were excluded following the initial visit if they met one or
78 more of the following exclusion criteria below: Long-acting glucocorticoids administration within the previous 60 days,
79 or other systemic/inhaled glucocorticoids within the previous 30 days; administration of megestrol acetate or

progesterogens within the previous 6 months; an IGF-1 concentration of >1000ng/ml, indicating likely hypersomatotropism¹²; concurrent hyperthyroidism. Cats that had undergone successful radioactive iodine therapy or thyroidectomy remained eligible; clinical signs of pancreatitis accompanied by consistent pancreatic changes on abdominal ultrasonography and/or an fPLI concentration of > 5.4µg/l; concurrent disease of which management would have affected ability to comply with trial commitments, including International Renal Interest Society stage 3 and 4 chronic kidney disease (CKD); patient aggression or fearfulness.

Re-examination Protocol

Enrolled cats had their insulin therapy changed to PZIR at the manufacturer's recommended starting dose of 0.2–0.7 U/kg q12h at the time of discharge from their first appointment. Doses were chosen based on cats' previous sensitivity to PLIS. All doses used throughout the study, including starting doses, were multiples of 0.5 U. Cats were re-examined at the Diabetic Remission Clinic 2, 4 and 12 weeks after starting PZIR therapy. At each visit, cats underwent assessment of serum fructosamine, DCS, 24 h water intake and a 24 h BGC. Results of 24 h BGCs at the 0, 2, 4 and 12 week re-examinations were used to calculate a mean blood glucose (MBG) concentration for each cat at these time points. Owners completed a repeated DIAQoL-pet questionnaire at 4 and 12 weeks. Cats also received a 12 h BGC at the 1 and 8 week time points, which could be performed at their primary care clinic, or at home using a handheld glucometer (AlphaTRAK 2; Zoetis), and could receive additional BGCs at home, or in primary practices, if clinically indicated.

PZIR dose was altered following each BGC over the 12 week trial using a nadir-based protocol (Table 2). All hypoglycaemic events, and episodes of diabetic remission, following transition to PZIR were recorded. Diabetic remission was defined as maintenance of normoglycaemia without receiving antihyperglycaemic medication (with the exception of low-carbohydrate diet) for a minimum of 28 days. This definition was based on the findings of a recent systematic review examining feline diabetic remission.⁵ Hypoglycaemia was defined as a blood glucose (BG) concentration of <3 mmol/l (<54 mg/dl), based on a definition used by recent guidelines on the treatment of feline DM.⁴ Any clinical signs compatible with hypoglycaemia, and other potential adverse events, were also recorded.

A basic comparison of duration of insulin action was carried out between week 0, when cats were treated with PLIS, and week 4, when treated with PZIR. The week 4 time point was chosen a priori for calculation of PZIR duration of action in order to provide sufficient time for cats to stabilise on PZIR, yet also being early enough in the trial's follow-up to avoid

substantial numbers of trial withdrawals or the development of other complicating factors, such as cats entering diabetic remission. Duration of insulin action was calculated as the time from insulin administration until the time at which BG returned to its pre-insulin value following the patient's BG nadir. This duration of action was calculated for each 12 h period in a cat's 24 h BGC, and a mean value was taken. When BG did not return to its pre-insulin value, the duration of action for that 12 h period was assigned a value of 12 h, which is referred to as 'long duration of action' in the remainder of this study. In contrast, duration of action of <9 h was defined as 'short duration of action' on the basis that clinical signs are often expected when insulin duration is below this cut-off.¹ Any BGCs in which insulin was withheld to check for diabetic remission were excluded from duration of action analysis.

117

118 *Statistical Analysis*

Continuous and ordinal data were assessed for normality using Shapiro–Wilk normality tests, and examination of descriptive statistics and histogram plots. Average values are expressed as mean \pm SD for normally distributed data, and as median (interquartile range [IQR]) for non-normally distributed data.

122

Linear mixed-effects modelling with compound symmetry covariance structure assessed for change in MBG, serum fructosamine, q12h insulin dose, DCS and DIAQoL-pet score over the 12 week study period. Twice-daily insulin dose and DCS were logarithmically transformed in order to meet the model's assumptions. For parameters that showed significant change, Fisher's least significant difference post-hoc comparison was used to assess pairwise differences between each time point and week 0.

128

The proportion of cats affected by short and long duration of insulin action at the week 0 and week 4 time points was compared using McNemar's tests. Cats with a short duration of action with either insulin type were evaluated to assess whether short duration of action was also present using the alternative insulin. A McNemar's test was also used to assess the proportion of cats that had at least one hypoglycaemic reading detected during week 0 and week 4 BGCs.

133

134 **Results**

Thirty-nine cats were examined for trial inclusion between October 2013 and September 2015. Seventeen cats were excluded with the most common reasons being identification of neoplasia or mass lesions (n=4), likely hypersomatotropism (n=3), excessive fearfulness or aggression (n=3) and cats being non-insulin dependent based on in-

138 hospital blood glucose monitoring (n=2). Other reasons for exclusion were hyperthyroidism, gastrointestinal disease,
139 suspected forebrain disease, hypertrophic cardiomyopathy with congestive heart failure, and owners' deciding against
140 study enrolment (all n=1).

141

142 All remaining 22 recruited cats completed the study. They consisted of 12 neutered males and 10 neutered females, and
143 included 17 domestic shorthaired cats, two domestic longhaired cats and one cat of each of the following breeds:
144 Abyssinian, Norwegian Forest and Singapura. Mean age was 10.6 ± 2.3 years and mean bodyweight was 4.9 ± 1.4 kg with a
145 median body condition score of 6 (4–7 out of 9). Cats had been diagnosed with DM a median of 71 (49–93) days prior to
146 enrolment and had been receiving BID porcine lente insulin for a median of 65 (47–91) days before enrolment. The
147 median PLIS dose received by cats at time of enrolment was 0.5 U/kg (0.3–0.7 U/kg) BID and median starting dose of PZIR
148 was 0.5 U/kg (0.3–0.7 U/kg) BID. Seventeen owners (77%) used home blood glucose monitoring (HBGM) as part of their
149 cat's diabetic management.

150

151 Table 3 shows change in indicators of diabetic control over the 12 week study, and change in q12h insulin dose over the
152 trial period is illustrated in Figure 1. A significant change was detected in serum fructosamine ($P = 0.00007$), q12h insulin
153 dose ($P = 0.02$), DIAQoL-pet AWIS ($P = 0.003$) and DCS ($P = 8.1 \times 10^{-8}$) over the study period. Post-hoc analysis revealed
154 a significant decrease in both serum fructosamine concentration and DCS at every time point compared with week 0,
155 whereas q12h insulin dose and DIAQoL-pet AWIS had both significantly decreased compared with baseline by week 12
156 (Table 3). MBG did not significantly change over the study period ($P = 0.1$). Over the trial period, 12 cats (55.4%)
157 experienced an increase in body weight, nine cats (41%) experienced a decrease in body weight and body weight
158 remained stable in one cat. Median water intake while hospitalised decreased from 28.6 ml/kg/day (IQR 19.6–39.0
159 ml/kg/day) at week 0 to 10.4 ml/kg/day (IQR 7.5–16.8 ml/kg/day) by week 12. Four cats (18.2%) entered diabetic
160 remission during the study period, and one other cat had stopped PZIR by week 12 and went on to achieve successful
161 remission.

162

163 The study population underwent a total of 190 12-hour periods of blood glucose monitoring during the trial and
164 hypoglycaemia was detected during 30 (15.8%) 12-hour BGCs. Five cats experienced a hypoglycaemic episode associated
165 with clinical signs. One clinical hypoglycaemic episode was detected during a BGC, whereas four were confirmed by
166 owners performing a BG check on their cat at home. Clinical signs included subdued demeanour (n=1), incoordination

167 (n=3), incoordination and blindness (n=1) and collapse (n=1). Three cases resolved with feeding and two cases recovered
168 following intravenous glucose treatment. One of the five cats with symptomatic hypoglycaemia subsequently achieved
169 remission. Seven out of 22 cats (31.8%) became hypoglycaemic during their week 0 BGC using PLIS, whereas four out of
170 19 cats (21.1%) that had a BGC at week 4 using PZIR experienced hypoglycaemia during their week 4 curve ($p=1.0$). One
171 cat developed a transient subcutaneous swelling at the site of PZIR administration. This swelling resolved without
172 treatment.

173

174 Three cats were excluded from the analysis of duration of insulin action due to cessation of insulin by the 4-week time
175 point (n=2) or missing the 4-week re-examination due to owner illness (n=1). Six cats showed a short duration of action
176 on PLIS compared to 2 cats after 4 weeks on PZIR ($p=0.18$) (Figure 2). Six cats had long duration of action when treated
177 with PZIR, compared to 3 cats when treated with PLIS ($p=0.5$). All six cats that had a short duration of insulin action on
178 PLIS, had a duration of PZIR action of ≥ 9 hours 4 weeks after transitioning. Both cats with short duration of action on PZIR
179 had a duration of action ≥ 9 hours on PLIS (Figure 2).

180

181 Discussion

182 This study examined the efficacy of transitioning BID PLIS-treated diabetic cats onto BID PZIR using a standardised nadir-
183 led insulin dosing protocol. Transitioning was associated with an improvement in both fructosamine concentration and
184 DM-related clinical signs within two weeks of starting PZIR. Pet and owner quality of life, measured by the DIAQoL-pet
185 tool, had significantly improved within 12 weeks of starting PZIR treatment, and 22.7% of cats achieved diabetic remission
186 after starting PZIR. Individual cats that suffered from short duration of action on PLIS, showed a resolution of this problem
187 4 weeks' following transition onto PZIR. Hypoglycaemia was the most common side effect, as expected, although was
188 rarely symptomatic, and was treated successfully. Hypoglycaemia was not more common with either insulin type when
189 two, a priori, set time points were compared ($t=0$ and $t=4$ weeks post-transition).

190

191 The findings of this study support that PZIR is an effective therapy for feline DM, and that PZIR treatment can be
192 associated with a rapid improvement in fructosamine concentration and clinical signs of DM in cats previously treated
193 with PLIS, when used in a nadir-based protocol alongside a low-carbohydrate, high-protein diet. These findings agree
194 with two previous trials which examined the efficacy of PZIR,⁹ or related bovine- and porcine-derived PZI,¹³ in both insulin-
195 naïve and previously-treated diabetic cats. These studies found that approximately 85% of the whole study population

196 achieved good glycaemic control within 45 days of starting PZI therapy, and 65–70% of previously-treated feline diabetics
197 showed good diabetic control after 45 days of PZI therapy. However, both of these studies contained only one cat treated
198 with PLIS q12h, so were unable to assess specifically the outcome of transitioning PLIS-treated cats onto PZI. This new
199 information will be of particular interest to veterinarians looking for alternative insulin types for feline diabetics in the
200 UK and Europe where PLIS has, until recently, been the only licensed veterinary insulin.

201

202 The current study is the first to include quality-of-life measurement as a treatment outcome and found that cat and
203 owner quality of life, measured using a validated quality-of-life tool, was unchanged from baseline after 1 month of PZIR
204 treatment, and had significantly improved by the end of the 12 week study. This improvement could be owing to owners
205 and cats becoming accustomed to the treatment of DM, and that some cats had achieved diabetic remission by this time
206 point. Remission would be expected to substantially increase the wellbeing of cats and their owners. Change in patients'
207 clinical signs of DM has been assessed by previous studies on PZI treatment in diabetic cats.^{9,13} However, this is also the
208 first study to use a validated clinical scoring system¹⁰ as a treatment outcome. This aimed to more consistently assess
209 changes in clinical signs in a standardised fashion, representing an improvement on the methodology of previous trials.

210

211 PZI is proposed to offer better glycaemic control than PLIS in diabetic cats.^{3,4} This recommendation is founded on previous
212 studies, which have shown PZI to have a longer duration of action compared to lente insulin in healthy cats,⁵ as well as
213 compared to other intermediate-duration insulin types in diabetic cats.¹⁴ To our knowledge, no previous study has
214 compared the duration of action of PZIR and PLIS in diabetic cats. Assessing duration of insulin action can be particularly
215 challenging in diabetic cats because duration of action cannot be reliably estimated when cats have poorly-controlled
216 DM.¹ Also, diabetic cats typically receive insulin therapy every 12 hours so measuring insulin duration of >12 hours would
217 not be possible without disrupting their usual treatment regime. This limitation prevented a precise comparison of insulin
218 duration in the current study, which instead assessed duration of action characteristics by focussing on those cats that
219 had a short duration of action when treated with either insulin type. All cats with a short duration of insulin action on
220 PLIS showed a resolution of this problem when treated with PZIR. However, two cats were found to have a short duration
221 of action when transitioned to PZIR, but did not experience this problem on PLIS. Furthermore, nine other cats showed a
222 decreased duration of action with PZIR compared with PLIS, but remained above the limit for short duration of action
223 (Figure 2). These findings support previous observations that individual diabetic cats can be unpredictable in their
224 response to different insulin types,¹ or their response can be difficult to reliably ascertain due to documented day-to-day

225 variability of BGCs in feline diabetics.¹⁵ The employed CGMS had a maximum measureable blood glucose limit of >22.2
226 mmol/l (400 mg/dl) and this could have artificially reduced the estimated duration of action in some cats.

227

228 The only adverse events recorded in this study were hypoglycaemia, as expected, and one episode of transient swelling
229 at the site of PZIR administration. The latter has been previously reported in PZIR- treated cats.⁹ Hypoglycaemia was
230 detected in 15.8% of all 12-hour BGCs in the current study. This is slightly greater than the proportion of 9.0% detected
231 in a trial examining the use of bovine/porcine PZI,¹³ but less than the proportion of 22.2% reported in a previous study
232 examining PZIR.⁹ A higher proportion of cats in the current study experienced confirmed clinical hypoglycaemia compared
233 to these two previous trials. However, most cat owners in the current study used HBGM to monitor their cat's DM,
234 whereas neither previous trial utilised HBGM. Four out of five clinical hypoglycaemic episodes in the current study were
235 diagnosed by owners performing a one-off BG measurement at home. This frequent use of HBGM likely increased the
236 number of confirmed clinical hypoglycaemic episodes compared to previous trials because owners were likely to
237 promptly check their cat's blood glucose if concerned about their pet's demeanour. This is in contrast to a previous PZIR
238 study where BG measurement could not be performed at several times of suspected hypoglycaemia.⁹ Finally, the current
239 study predominantly carried out BGCs using CGMSs, which have been shown to detect hypoglycaemic BGC nadirs that
240 might be missed during curves using handheld glucometers.¹⁶ This is likely to have ensured that most hypoglycaemic
241 episodes in this trial were successfully documented, especially when combined with the frequent use of HBGM. The
242 current study is therefore likely to provide an accurate estimate of the incidence of hypoglycaemia among diabetic cats
243 transitioned to PZIR.

244

245 This study did not document an improvement in MBG concentration over the duration of the study, despite an
246 improvement in serum fructosamine concentration, diabetic clinical score and several cats entering diabetic remission.
247 As mentioned, the CGMS used to perform BGCs in this study had a maximum measureable blood glucose concentration
248 of 22.2mmol/l (400mg/dl) and this might have lowered the MBG values in some cats. Also, the BGCs, from which MBG
249 was calculated, were all performed in hospital and so stress-related hyperglycaemia from hospitalisation might have
250 affected blood glucose measurements.¹⁷ The study did aim to minimise stress-related hyperglycaemia by using a CGMS
251 to perform BGCs and ensuring that all BGCs were performed more than 12 hours after hospital admission. Although no
252 significant change was detected in MBG, the fact that serum fructosamine concentration improved over the study period

suggests that these in-hospital curves might not reflect cat's diabetic control between re-examination visits. The earlier mentioned day-to-day variability of blood glucose curves, is also likely associated with this apparent discrepancy.¹⁶

This study has some limitations. Although several markers of glycaemic control improved in included cats following their transition from PLIS to PZIR, the current trial was not designed to demonstrate unequivocally that one insulin preparation is superior to the other. For instance, it is theoretically possible that included cats would also have experienced an improvement in their diabetic control had they remained on PLIS therapy for longer. Despite this, cats in the current study showed a rapid, significant improvement in diabetic clinical signs after starting PZIR, whereas a previous trial found that three months of PLIS therapy was needed to produce a significant improvement in clinical signs when starting diabetic cats on PLIS.¹⁸

Other factors might also have played a role in the documented improvement post-transition in the current trial. These factors include the use of a nadir-based protocol, support from a dedicated DM clinic during the trial, and the fact that cats were transitioned onto a low-carbohydrate diet as part of the study protocol. Dietary carbohydrate restriction is likely to promote good glycaemic control in diabetic cats.¹⁹ In this study, cats' transition to a low-carbohydrate, high-protein diet was carried out before trial recruitment to allow the benefits of feeding a low-carbohydrate, high-protein diet to take effect before their first study visit. Although the minimum time for this diet change was 10 days prior to enrolment, recruited cats were fed a low-carbohydrate high-protein diet for a median of 29 days before their first trial appointment. A previous study found that insulin-treated diabetic cats experienced a significant decrease in MBG within 4 weeks of transitioning to a low-carbohydrate diet.¹⁹ It is therefore likely that the benefits of changing to a low-carbohydrate, high-protein diet in the current study were already apparent by many cats' first trial appointment and that subsequent improvements in glycaemic control were more related to other factors, including the change from PLIS to PZIR. Overall, an unbiased direct comparison of PZIR and PLIS would ultimately require an adequately powered, well designed, randomised clinical trial, which is currently lacking in the veterinary literature, as discussed in a recent systematic review of feline diabetes treatment.⁵ However, the transition process described in this study provides valuable information, which can now be used to inform effective and safe transition protocols for cats requiring transition from PLIS to PZIR, for one reason or another.

281 **Conclusions**

282 Diabetic cats treated with intermediate-acting PLIS can be transitioned successfully onto long-acting PZIR using a nadir-
283 led protocol and the manufacturer-recommended starting dose. Overall, transitioning to PZIR was associated with an
284 improvement in fructosamine, diabetic clinical signs, and pet and owner quality of life, and a proportion of cats entered
285 diabetic remission after starting PZIR therapy. Incidence of hypoglycaemia was similar to previously described with PZI
286 use in cats. This documented incidence of hypoglycaemia, the potential for diabetic remission, and the efficacy of PZIR at
287 modest dosages could nevertheless mean that conservative PZIR dosing should be used in cats where close monitoring
288 cannot be performed. In conclusion, this study provides a framework for transitioning diabetic cats from PLIS to PZIR, if
289 needed. Additionally, veterinary-licensed human recombinant PZI proved to be an effective treatment for feline DM and
290 was associated with increased duration of action in those PLIS-treated cats with short duration of action. Licensed long-
291 acting PZIR therefore provides a viable alternative treatment option to licensed intermediate-acting PLIS in feline
292 diabetics, especially since recent recommendation emphasise the preferential use of long-acting insulin types.

293

294 **Acknowledgments**

295 With thanks to the staff of the Royal Veterinary College Clinic Investigation Centre, Dr. Yu-Mei Chang for statistical
296 assistance, and to cats and owners who participated in this study.

297

298 **Conflicts of Interest**

299 This study was supported by Boehringer Ingelheim.

300

301 **Funding**

302 This research was supported by Boehringer Ingelheim. The Royal Veterinary College Diabetic Remission Clinic also
303 receives support from Nestlé Purina PetCare and Zoetis. The primary author's PhD was supported by the Beryl Evetts
304 and Robert Luff Animal Welfare Trust.

References

1. Reusch C. Feline Diabetes Mellitus. In: Feldman EC, Nelson R, Reusch C, Scott-Moncrieff C and Behrend E, (eds.). *Canine and feline endocrinology*. 4th ed. ed. St. Louis, Missouri: Saunders, 2015, p. 258 - 314.
2. Gottlieb S, Rand JS. Remission in cats: including predictors and risk factors. *Vet Clin North Am Small Anim Pract* 2013; 43: 245–249.
3. Rucinsky R, Cook A, Haley S, Nelson R, Zoran DL and Poundstone M. AAHA diabetes management guidelines. *Journal of the American Animal Hospital Association*. 2010; 46: 215-24.
4. Sparkes AH, Cannon M, Church D, et al. ISFM Consensus Guidelines on the Practical Management of Diabetes Mellitus in Cats. *Journal of feline medicine and surgery*. 2015; 17: 235-50.
5. Gostelow R, Forcada Y, Graves T, Church D and Niessen S. Systematic review of feline diabetic remission: separating fact from opinion. *Vet J*. 2014; 202: 208-21.
6. Marshall RD, Rand JS and Morton JM. Glargine and protamine zinc insulin have a longer duration of action and result in lower mean daily glucose concentrations than lente insulin in healthy cats. *Journal of veterinary pharmacology and therapeutics*. 2008; 31: 205-12.
7. Zini E, Osto M, Franchini M, et al. Hyperglycaemia but not hyperlipidaemia causes beta cell dysfunction and beta cell loss in the domestic cat. *Diabetologia*. 2009; 52: 336-46.
8. Marshall RD, Rand JS and Morton JM. Treatment of newly diagnosed diabetic cats with glargine insulin improves glycaemic control and results in higher probability of remission than protamine zinc and lente insulins. *Journal of feline medicine and surgery*. 2009; 11: 683-91.
9. Nelson RW, Henley K and Cole C. Field safety and efficacy of protamine zinc recombinant human insulin for treatment of diabetes mellitus in cats. *Journal of veterinary internal medicine / American College of Veterinary Internal Medicine*. 2009; 23: 787-93.
10. Venzon Varela F, Gostelow R, Forcada Y, Church D and Niessen SJ. The Diabetic Clinical Score (DCS): evaluation of a simple standardised quantification tool to allow rapid description of clinical signs in diabetic cats. *The Journal of small animal practice*. 2016: 31-2.
11. Niessen SJ, Powney S, Guitian J, et al. Evaluation of a quality-of-life tool for cats with diabetes mellitus. *Journal of veterinary internal medicine / American College of Veterinary Internal Medicine*. 2010; 24: 1098-105.
12. Niessen SJ, Petrie G, Gaudiano F, et al. Feline acromegaly: an underdiagnosed endocrinopathy? *J Vet Intern Med*. 2007; 21: 899-905.
13. Nelson RW, Lynn RC, Wagner-Mann CC and Michels GM. Efficacy of protamine zinc insulin for treatment of diabetes mellitus in cats. *Journal of the American Veterinary Medical Association*. 2001; 218: 38-42.
14. Moise NS and Reimers TJ. Insulin therapy in cats with diabetes mellitus. *Journal of the American Veterinary Medical Association*. 1983; 182: 158-64.
15. Alt N, Kley S, Haessig M and Reusch CE. Day-to-day variability of blood glucose concentration curves generated at home in cats with diabetes mellitus. *Journal of the American Veterinary Medical Association*. 2007; 230: 1011-7.

- 348 16. Dietiker-Moretti S, Müller C, Sieber-Ruckstuhl N, et al. Comparison of a continuous glucose
349 monitoring system with a portable blood glucose meter to determine insulin dose in cats
350 with diabetes mellitus. *Journal of veterinary internal medicine / American College of*
351 *Veterinary Internal Medicine*. 2011; 25: 1084-8.
- 352 17. Rand JS, Kinnaird E, Baglioni A, Blackshaw J and Priest J. Acute stress hyperglycemia in cats
353 is associated with struggling and increased concentrations of lactate and norepinephrine.
354 *Journal of veterinary internal medicine / American College of Veterinary Internal Medicine*.
355 2002; 16: 123-32.
- 356 18. Martin GJ and Rand JS. Control of diabetes mellitus in cats with porcine insulin zinc
357 suspension. *The Veterinary record*. 2007; 161: 88-94.
- 358 19. Bennett N, Greco DS, Peterson ME, . Comparison of a low carbohydrate-low fiber diet and
359 a moderate carbohydrate-high fiber diet in the management of feline diabetes mellitus. *J*
360 *Feline Med Surg* 2006; 8: 73–84.
361

362 Table 1: Diabetic Clinical Score (DCS) used to grade the severity of diabetes-associated clinical signs in
 363 participating cats

Clinical Sign Being Scored	Severity (compared to prior to onset of diabetes mellitus)	Assigned Score
Unintended weight loss over the past 2 months (Assessed using bodyweight records or measurements)	None, intended weight loss or weight gain	0
	Mild (<5% loss)	1
	Moderate (5-10% loss)	2
	Severe (>10% loss)	3
Increased drinking and/or urination (Assessed by questioning owner)	None	0
	Mild – some increase noted	1
	Moderate – increased filling of water bowl	2
	Severe – constantly seen to drink	3
Increased appetite (Assessed by questioning owner)	Normal or decreased appetite	0
	Mild – finishes food eagerly	1
	Moderate – finishes food eagerly and begs for more	2
	Severe – obsessed with food	3
Decreased activity/ attitude (Assessed by questioning owner)	Normal or increased activity	0
	Mild – slightly less active	1
	Moderate – certainly less active	2
	Severe – mainly lying around	3
	Total score	/12

364

Table 2: Nadir-based protocol used to alter PZIR dose according to the results of blood glucose curves during the 12-week study period

Blood Glucose Curve Results	Action Taken
Pre-insulin BG < 10mmol/l	Withhold insulin for 12 hours to check for remission
Pre-insulin BG > 10mmol/l + nadir of < 3mmol/l + nadir 3 - 6 mmol/l + nadir > 6 mmol/l	Reduce dose by 0.5-1 U/injection Keep dose unchanged Increase dose by 0.5-1 U/injection
Hypoglycaemia associated with clinical signs	Reduce total dose by 50%

Abbreviations: BG, Blood Glucose.

Table 3: Change in indicators of diabetic control among recruited cats over the 12-week study period. The table also indicates whether these changes reached statistical significance in linear mixed effects modelling.

Study Timepoint (week)	BID Insulin Dose (U/kg)	Mean BG (mmol/l) (mg/dl)	Serum fructosamine (μmol/l)	Diabetic Clinical Score (out of 12)	DIAQoL-pet AWIS
0	0.48 (0.32 – 0.75)	13.2 (9.1 – 18.8) 237.6 (163.8 – 338.4)	469 (358 – 588)	3 (1 – 6)	-1.7 (-3.4 - -1.0)
2	0.52 (0.30 – 0.65)	11.2 (6.2 – 16.1) 201.6 (111.6 – 289.8)	428 (349 – 505) *	3 (0 – 4) *	-
4	0.46 (0.19 – 0.67)	11.5 (7.2 – 16.3) 207.0 (129.6 – 293.4)	401 (309 – 454) **	0 (0 – 2) ***	-1.8 (-2.8 - -0.8)
12	0.39 (0.18 – 0.61) **	10.8 (8.3 – 15.4) 194.4 (149.4 – 277.2)	338 (266 – 486) ***	0 (0 – 2) ***	-1.0 (-2.2 - -0.4) **

Abbreviations: BID, Twice daily; BG, Blood glucose; AWIS, Average weighted impact score. All average values are reported as median (interquartile range). *, ** and *** indicate a significant difference compared to week 0 value in post-hoc analysis (* indicates p<0.5, ** indicates p<0.01, *** indicates p<0.001)

Figure 1. Scatterplot showing change in twice-daily insulin dose over study period. Red symbols indicate values when treated with porcine lente insulin suspension q12h, and blue symbols indicate values when treated with protamine zinc insulin solution q12h. **Significant difference compared with week 0 (P <0.01). Vertical line and error bars indicate median and interquartile range.




Figure 2: Paired scatterplot showing duration of action of porcine lente insulin suspension (PLIS) q12h and protamine zinc insulin solution (PZIR) q12h in individual cats. Dashed line indicates short duration of insulin action (<9 h). Red symbols indicate duration when treated with PLIS q12h, and blue symbols indicate duration when treated with PZIR q12h. Pairs of values from individual cats have joining lines. Two pairs of cats had the same duration of action before and after transitioning (two decreasing from 12 h to 10.9 h and two increasing from 7 h to 12 h), resulting in only 17 data pairs being visible on the graph.

