

This article has been published in a revised form in *Animal* [[10.1017/S1751731117002117](https://doi.org/10.1017/S1751731117002117)]. This version is free to view and download for private research and study only. Not for re-distribution, re-sale or use in derivative works. © The Animal Consortium 2017.

The full details of the published version of the article are as follows:

TITLE: Effect of a photoperiodic green light programme during incubation on embryo development and hatch process

AUTHORS: Tong, Q; McGonnell, I M; Demmers, T G M; Roulston, N; Bergoug, H; Romanini, C E; Verhelst, R; Guinebrière, M; Etteradossi, N; Berckmans, D; Exadaktylos, V

JOURNAL TITLE: *Animal*

PUBLICATION DATE: April 2018

PUBLISHER: Cambridge University Press: STM Journals

DOI: 10.1017/S1751731117002117

25 No light-dark cycle was used in the control group. Four batches of eggs (n=300 per
26 group per batch) from fertile Ross 308 broiler breeders were used in this experiment.
27 The beak length and crown-rump length of embryos incubated under green light were
28 significantly longer than that of control embryos at day 10 and day 12, respectively
29 ($P<0.01$). Furthermore, green light exposed embryos had a longer third toe length
30 compared to control embryos at day 10, day 14 and day 17 ($P=0.02$). At group level
31 (n=4 batches), light stimulation had no effect on chick weight and quality at take-off, the
32 initiation of hatch and hatch window. However, the individual hatching time of the light
33 exposure focal chicks (n=33) was 3.4 h earlier ($P=0.49$) than the control focal chicks
34 (n=36) probably due to the change in melatonin rhythm of the light group. The results of
35 this study indicate that green light accelerates embryo development and alters hatch
36 related hormones (thyroid and corticosterone) which may result in earlier hatching.

37

38 **Key words:** broiler incubation, green light, embryo growth, circadian rhythm, hatch
39 process

40

41 **Implications**

42 Chickens are incubated commercially in darkness throughout the entire 21 days, which
43 is different from nature incubation. Light exposure is important for embryos development
44 and circadian rhythm establishment. Our results recommend that application of green
45 light during the first 18 days of incubation accelerated hatch of individual chick which
46 may be due to the acceleration of embryo development and the alteration of hormone
47 profiles.

48

49 **Introduction**

50 Broiler chickens are often incubated commercially in complete darkness. Under natural
51 conditions, however, avian embryos would certainly receive some light stimulation
52 during development. Day-night (light-dark) rhythms, mediated by high concentrations of
53 the hormone melatonin (MT) at night and low levels by day, are a universal feature of
54 living organisms (Schultz and Kay, 2003). Avian embryos, developing outside the
55 maternal organism and without direct endocrine signals from the mother, may develop
56 early internal circadian systems to cope with the new life outside the egg. The circadian
57 rhythm in birds is indeed established early on in embryonic development and requires
58 external cues (Nichelmann *et al.*, 1999, Zeman *et al.*, 1999). Chicken pinealocytes, the
59 brain cells that release melatonin, were shown to be very sensitive to light with an
60 intensity of as little 10 lux in vitro and showed a circadian activity pattern (Faluhelyi and
61 Csernus, 2007). Melatonin production starts early on in the incubation period: in chicken
62 embryos from day 10, but no regular rhythm is detected until between day 16 and day
63 18 provided an external day/night rhythm is imposed. The amplitude of the rhythm
64 increases considerably during the last 2 days of incubation at the time of internal and
65 external pipping and the beginning of lung respiration of the embryos (Starck and
66 Ricklefs, 1998). However, in artificial incubation, eggs are handled under (almost)
67 continuous darkness and a previous study showed that the hormone production of the
68 pineal does not become rhythmic between day 15 and day 18 (high levels of MT release
69 were experienced and showed apparently random alterations) (Csernus *et al.*, 2007).
70 This suggests that no circadian rhythm develops during incubation in completely dark
71 conditions, because of the lack of appropriate environmental signals stimulating the
72 embryo. Establishment of the embryonic circadian rhythm can have impact on the

73 functioning of the circadian clock of pre- and peri-natal chick. The circadian rhythm
74 established in the embryo determines the timing of hatching and the hatchlings day-
75 night rhythm until at least three days after hatching (Zeman *et al.*, 1999, Zeman and
76 Gwinner, 1993). The rhythm of melatonin levels, which is established in the embryo, can
77 also affect behavioural rhythmicity (Archer *et al.*, 2009). Melatonin has been shown to
78 act as an anti-stress agent. Melatonin can suppress stress-induced increases in rat
79 plasma corticosterone concentrations, and animals subjected to stress showed altered
80 circadian patterns in plasma melatonin with elevated corticosterone concentrations
81 (Mocchegiani *et al.*, 1999, Barriga *et al.*, 2002). Melatonin is able to modulate stress at
82 both the central and peripheral levels by exerting its inhibitory role in the hypothalamic-
83 pituitary-adrenal (HPA) axis in chicks and by suppressing corticosterone production
84 (Saito *et al.*, 2005).

85 Melatonin rhythms during incubation can affect the behaviour, stress and hatch
86 performance of poultry. However, there has been relatively little systematic work
87 assessing the pattern and type of light stimulation needed to produce these effects or
88 the mechanism underlying them. In the wild, an incubating hen generally comes off her
89 nest once a day between 11:00 and 14:00 h, occasionally as early as 9:25 and as late
90 as 15:00, presumably when the eggs would be least subject to heat loss and are
91 completely exposed to daylight. The time spent off the nest seems fairly consistent and
92 averages 38.7 ± 1.2 min (Duncan *et al.*, 1978). In addition, when turning the eggs, the
93 hen usually stands up, thereby exposing the eggs to more light and to lower ambient
94 temperature. The temporary exposure to light means that the full spectrum of radiation
95 may potentially reach the surface of the avian eggs. Depending on the nest
96 environment, eggs will experience light from the heating infra-red wavelengths to the

97 potentially mutagenic ultra-violet light. However, base colour pigments of eggshell are
98 likely to control the light that reaches the embryo by blocking light of harmful infrared
99 and UV wavelengths but admitting beneficial wavelengths (Maurer *et al.*, 2011).

100 Previous studies have investigated the impact of light in the incubator environment.
101 Exposing eggs to green light (1340-1730 lux) from 5 to 15 days of incubation increased
102 embryo growth and hatchability by 4.8% (Shafey and Al-mohsen, 2002). An overall
103 improvement in the embryo/hatchling survival rate was also observed following a near-
104 infrared (630-1000 nm) light-emitting diode (LED) exposure. There were also increases
105 in mean body weight, crown–rump length, liver weight and decrease in hatchling
106 residual yolk weight (as a function of post-hatch survival time and increased nutrient
107 utilisation during development) (Yeager *et al.*, 2005). Continuous green light (1340-1730
108 lux) during the first 18 days of incubation accelerated hatching times by about 24 hours
109 in meat-type breeder eggs (Hybro). Far-red (670nm) LED-exposure once per day from
110 0-20 days of incubation resulted in chickens pipping (breaking the shell) 2.92 hours
111 earlier and there was a 2.91 hours shorter duration between pip and hatch.

112 Incandescent light (12L:12D) accelerates hatching times ($P < 0.01$) without affecting
113 hatchability, weight at hatching, liver or heart growth in turkey eggs (Fairchild and
114 Christensen, 2000, Shafey and Al-mohsen, 2002, Yeager *et al.*, 2005). Therefore, light
115 intensity can play an important role in the speed of development of the avian embryo
116 and hatchability, at least in the domestic broiler chicken. Furthermore, light exposure
117 and the source of light could prove to be important factors in achieving synchrony or
118 significantly reducing the spread of hatch under artificial rearing conditions. Bird and
119 human perception of colour differs. Human beings have 3 type of cones and birds have
120 four which are responsible for seeing frequency (colours). Therefore, chicks can see

121 much broader spectrum than human (Osorio *et al.*, 1999). A broader range of studies is
122 available for the effect of wavelength, but the average wavelength of the most sensitive
123 light for chicken embryos is about 550-560nm (Rogers *et al.*, 1998) and green light
124 seems to be the most effective in stimulating embryonic growth and development, as
125 well as post-hatch growth in chickens (Halevy *et al.*, 2006, Rozenboim *et al.*, 2013).
126 It would appear that light can have significant impact on the developing chick embryo,
127 however it is far from clear what the optimal light intensity for embryonic development in
128 the wild is, thus it is difficult to predict which lighting regimes have the most benefit in the
129 incubator setting. Based on the natural incubation behaviour of mother hens, tightly
130 controlled light intensities and wavelengths and natural patterns of illumination were
131 applied in this study. The purpose was to examine the effect of green light during
132 incubation upon the hatch process and to determine whether a melatonin rhythm would
133 be established by such light stimulation, and the subsequent impact on thyroid and
134 corticosterone hormone production which potentially regulates the hatching process.

135

136 **Material and methods**

137 *Eggs and incubation*

138 Animal experiments were performed with the ethics approval from the Royal Veterinary
139 College Animal Ethics Committee. Four incubation trials were conducted. In each trial two
140 incubators swapped between the control and light groups were used and each incubator
141 was able to set 300 eggs in 2 trays (Petersime NV, Zulte, Belgium). In total, four batches
142 of fertile Ross 308 eggs (600 each batch) were obtained from a local supplier (Henry
143 Stewart & Co. Ltd, Lincolnshire, UK). The eggs were incubated using a standard
144 incubation profile and the incubation conditions were continuously monitored and

145 controlled by the incubator controller (BIO-IRIS, Petersime™). All incubation conditions
146 (machine temperature, humidity, CO₂ concentration and ventilation rate) were identical in
147 the two incubators.

148

149 *Light protocol*

150 A light-dark cycle (12L:12D) was provided using green light (460-580 nm) in the first 18
151 days of incubation and no light-dark cycle in the last three days. The light cycle consisted
152 of 12 hours darkness followed by 12 hours light period, which was controlled using a
153 mechanical timer. The light period consisted of 4 hours of low intensity (100-130 lux) green
154 light, followed by one hour continuous illumination at a high intensity at 1200-1400 lux and
155 the remaining 7 hours at the low intensity (100-130 lux) (Shafey and Al-mohsen, 2002,
156 Shafey *et al.*, 2005). In the light group, to produce an even spread of illumination on the
157 surface of each egg and no extra heat produced by the light source, the monochromatic
158 light was provided by a total of 204 green low power LEDs (0.5W Power PLCC4 SMT,
159 AVAGO TECHNOLOGIES) mounted in a frame which was placed above the eggs. The
160 light intensity at the egg surface was measured using a light meter (Testo luminous
161 intensity measuring instrument 545, GmbH & CO. Germany). The light intensity could be
162 guaranteed for the top tray, so only the eggs from top tray were sampled. However, the
163 bottom tray was also set to ensure the other incubation parameters, such as heat
164 production and CO₂ concentration were not negatively affected. No light-dark cycle was
165 provided in the control incubator.

166

167 *Monitoring of hatch window and hatching time*

168 The onset of hatch (IP) and hatch window (HW), of each incubation, was determined by
169 the incubator controller (Petersime BIO-IRIS™) (Tong *et al.*, 2015). In total, 10 focal eggs
170 in each incubator from the first two experiments and 20 focal eggs of each incubator from
171 the last two experiments were randomly selected, weighted and individually labelled to
172 determine the individual hatching time. During transfer to the hatchings baskets, each
173 focal egg was placed in a specially constructed area within the top basket (using 8 x 8 x
174 8 cm³ metallic mesh grid). The hatching time of each focal egg was determined using
175 eggshell temperature (EST) as previously described by Romanini *et al.* (2013). The
176 identification of hatching time is based on a registered EST drop (2-5 °C) when the chick
177 emerges from the egg.

178

179 *Hatch performance, embryo and blood parameters assessment*

180 All eggs were candled on 18 days and 12 hours of incubation time and those with evidence
181 of a living embryo were transferred from the turning trays to hatching baskets within half
182 an hour. Both incubators were stopped after 512 hours (21 days and 8 hours) of incubation.
183 Hatchability (the percentage of fertile eggs that hatch), early death (ED) from day 0 to day
184 7, middle death (MD) from day 8 to day 15, late death (LD) from day 16 to day 21, and
185 mortality were determined at the end of incubation based on breakout results. All the
186 hatched chicks from the top basket of each group were scored for quality using a
187 standardised method (Tona *et al.*, 2003) and chicks with full score (100%) were
188 considered as first class chicks. Samples of three eggs or chicks were selected randomly
189 from the top tray in each incubator and were collected at eight incubation stages: day 10,
190 day 12, day 14, day 16, day 17, day 18, day 19 (internal pipping, IP), day 20 (external
191 pipping, EP) and take-off (focal chick). Embryos or chicks were euthanized and blood was

192 collected from allantoic vein or the left ventricle, respectively. The chicks and organs (heart,
193 liver and stomach) were dissected and weighed. The length of the beak and third toe as
194 well as the crown-rump length (CRL) were measured.

195 The blood of three embryos or chicks was sampled at every three hours over an 18 hours
196 period (from 18:00 of day 17 to 12:00 of day 18), at internal pipping on day 19, external
197 pipping on day 20 and take-off on day 21. Different eggs or chicks were sacrificed at each
198 time point of sampling. The blood sample was collected into a heparin tube and
199 centrifuged at 3000 rpm for 10 min. The plasma was stored at -20°C until hormones
200 measurement. The blood samples from embryos or chicks collected at the same time
201 point were pooled to measure MT and other hatching related hormones. Plasma
202 hormones levels were measured using a commercial chicken melatonin (MT) ELISA kit,
203 chicken Tri-iodothyronine (T3) ELISA kit, chicken thyroxine (T4) ELISA Kit (CUSABUO
204 BIOTECH CO., Ltd, Wuhan, China) and Corticosterone HS EIA kit (IDS Ltd, Boldon,
205 England).

206

207 *Statistical analysis*

208 Data was analysed using SPSS (PASW statistics 20) and was presented as the mean ±
209 standard error of the mean (SEM). A linear mixed model was used to analyse the effect
210 of light treatments (control and light) on hatchability, mortality (ED, MD and LD), HW, chick
211 quality and chick weight:

$$212 Y = \mu + \text{light treatment} + \text{incubator} + \text{batch} + \epsilon$$

213 A second linear mixed model was used to analyse the effect of light treatments and
214 incubation stage on embryonic parameters, blood values and plasma hormones
215 concentrations. The model was:

216 $Y = \mu + \text{light treatment} + \text{incubation stage} + \text{interaction (treatment} \times \text{incubation stage)}$
217 $+ \text{incubator} + \text{batch} + \epsilon$

218 The μ is the overall mean and ϵ is the residual error term. Light treatment, incubation stage,
219 interaction, incubator were fixed effects; batch was a random effect. The interaction was
220 removed from the original model when it was not significant. When the effect was
221 statistically different ($P \leq 0.05$), the means were further compared using Least Significant
222 Difference (LSD) test or nonparametric statistics (only for chick quality).

223

224 **Results**

225 *Hatch performance and embryo development*

226 There was no effect of incubator on hatchability and different stages of mortality taking
227 into account the batch effect. However, the hatchability of the control group was
228 significantly higher than the light group because of the combination of mortality from three
229 stages (Figure 1; $P = 0.03$) (78.2 versus 73.7%). No effect of light treatment was found on
230 chick quality and chick weight for four batches. The egg weight at setting for the control
231 and light groups were $59.96 \pm 1.7\text{g}$ and $59.58 \pm 1.7\text{g}$, respectively. The light treatment had
232 an effect on the heart weight ($P = 0.001$), beak length ($P = 0.006$), third toe length ($P = 0.024$)
233 and the crown-rump length ($P = 0.005$) at specific stages but not throughout the incubation
234 period. The differences in the heart weight, beak length, third toe length and the crown-
235 rump length between the control and light groups at each incubation stage are shown in
236 Figure 2.

237

238 *Hatch window and hatching time*

239 There was no effect of the light treatment and incubator on the initiation and the length of
240 the hatch window of the entire batch (Table 1). Both groups started pipping around 467
241 hours of incubation time. No difference in the hatching time of the first focal chicks in the
242 control and light groups was observed, which was the same as the IP detected at the
243 group level. However, the majority of focal chicks in the light group hatched earlier than
244 the control focal chicks ($P=0.049$). The average hatching time of the focal chicks in the
245 control and light groups were 487.4 ± 1.2 hours and 484.0 ± 1.1 hours respectively. The focal
246 chicks of the light group hatched 3.4 hours earlier than the focal chicks of the control group.

247

248 *Melatonin and hatch related hormones*

249 Overall the mixed effects models, taking into account batch effect, showed that there were
250 no effects of light treatment and incubator on MT, T3, T4 and the T3/T4 ratio. However,
251 the individual sampling time point had a significant impact on the plasma hormone levels
252 ($P<0.01$). The plasma MT in the control and light groups from day 17 to day 21 is shown
253 in Figure 3. There was no statistically significant difference in MT between groups at any
254 time point in the 18 hours period between days 17-18, however there were differences in
255 the pattern. In the control (no light group) MT levels increase rapidly in the first 3 hours of
256 the dark period (from 85.32 to 100.68 pg/ml) before levelling off at 6 hours and
257 experiencing a more gradual increase up to a peak at 12 hours (109.07pg/ml),
258 experiencing an overall 1.28-fold change. In contrast, the light group experienced a much
259 more gradual increase in the first 6 hours (from 82.42 to 88.82 pg/ml), followed by a rapid
260 increase at 9 hours, which was the peak plasma level (109.23 pg/ml). The total increase
261 was similar at 1.33 times of the base level. The control group showed a rapid decrease in
262 plasma MT levels between 12-15 hours (from 109.07 to 89.14 pg/ml) before returning to

263 a similar level at 0 hour (which was 85.32 pg/ml). In contrast the light exposed embryos
264 experienced a gradual drop in MT levels from 9 hours, i.e. from the beginning of the dark
265 period and continuing to drop to 94.55 pg/ml at 18 hours of the light period. This level had
266 not returned to the level seen at 0 hour (which was 82.42 pg/ml) suggesting there would
267 be further gradual decrease in plasma MT during the remainder of the light period. Data
268 obtained during days 19-21 was at a period when both groups of eggs were in dark
269 conditions. At the beginning of IP the levels of plasma MT had increased in both groups,
270 with the control being higher than the light group (149.18 versus 130.34 pg/ml). These
271 levels increased rapidly during the next 24 hours, particularly in the control group ($P=0.03$),
272 however the data did not show any significant differences between groups. At take-off
273 (day 21), plasma MT levels decreased significantly and returned to the levels seen at day
274 19 in both groups.

275 Figure 4 shows the 18 hours pattern of the plasma CORT levels. In the control group,
276 CORT varied between 6 and 7.5 ng/ml throughout the dark and light periods; in the light
277 group CORT decreased slightly in the first 3 hours of the dark period (from 7.61 to 6.10
278 ng/ml), followed by a steady increase until peaking at 18 hours (8.94 ng/ml). Plasma
279 CORT levels of the control chicks were numerically higher than those of the light
280 stimulated chicks in the 12 hours dark period. This was reversed in the 12 hours light
281 period, where the light group had significantly higher CORT than the control group in the
282 first 3 hours of illumination ($P=0.01$). The plasma CORT levels increased in both groups
283 from day 18 to EP. The higher trend was kept in the light group at IP and EP, but not
284 statistically different between groups and then returned to the same level (11ng/ml) as at
285 take-off.

286 Plasma T3 followed a similar trend to CORT (Figure 5). There was no significant difference
287 between the control and light groups in the dark period. In both groups, T3 levels
288 increased rapidly in the first 6 hours of the dark period, reaching the same levels (0.97
289 ng/ml) before dropping slightly in the next 3 hours of the dark period. The light group
290 increased further in the last 3 hours of the dark period, peaking at 12 hours (1.02 ng/ml),
291 followed by being stable in the light period. The average T3 level in the light embryos was
292 significantly higher than the control embryos ($P=0.02$) during this light period, indicating
293 that light stimulation had an effect on T3 levels. The T3 levels increased significantly (more
294 than doubled) from day 18 to IP in both groups, followed by another dramatic increase at
295 EP which was the peak level (2.90 and 3.76 ng/ml in the control and light groups,
296 respectively). However, the levels at take-off dropped significantly to the similar levels at
297 days 17-18 in both groups. There were no differences in T3 concentrations between the
298 control and light groups at IP and Take-off, but the light stimulated chicks had numerically
299 higher T3 concentration than the control chicks at EP. The plasma T4 concentration is
300 presented in Figure 6 and it showed a similar trend to MT between the control and light
301 groups. The plasma T4 levels of the control embryos had an increase in the first 3 hours
302 of the dark period (from 55.03 to 62.24 ng/ml) before being stable over the next 6 hours
303 of the dark period and experiencing a slight increase up to a peak at 12 hours (65.24
304 ng/ml). In contrast, the light group had a slight increase in the first 6 hours, followed by
305 rapid increase at 9 hours, which was the peak plasma level (70.03 ng/ml), and
306 experiencing a large drop at 12 hours. The control group showed a rapid decrease in
307 plasma T4 levels between 12-15 hours (from 65.24 to 52.44 ng/ml) and had an increase
308 at 18 hours before returning to the same level at 0 hour (58.27 ng/ml). However, the light
309 exposed embryos experienced a small increase in T4 levels from the beginning of the light

310 period, followed by a minor drop into the same levels of the control group at 18 hours.
311 Plasma T4 increased significantly from days 17-18 to IP in both groups with the control
312 being higher than the light group (92.79 versus 80.00 ng/ml). These levels kept increasing
313 until peaking at EP, particularly in the control group ($P < 0.01$). In contrast to T3, the control
314 EP chicks tended to have higher T4 than that of the light stimulated chicks. However, the
315 data did not show any significant differences between groups. At take-off (day 21), plasma
316 T4 levels decreased significantly in the control group ($P < 0.05$) before returning to the
317 same levels of the light group (86 ng/ml). The plasma T3/T4 ratio of the control and light
318 groups for the observed part of the incubation period is shown in Figure 7. The T3/T4 ratio
319 during the 18 hours of days 17-18 was stable at 0.02 in both groups. However, this ratio
320 started increasing before the onset of pipping and reached the maximum at EP in light
321 group (0.04), whereas it remained at the same elevated level in the control group (0.03).
322 These ratios at take-off in both groups returned to the same figure observed at days 17-
323 18.

324

325 **Discussion**

326 Green light stimulation during incubation has been reported to accelerate chick embryo
327 development (Zhang *et al.*, 2014). The results shown in this study also indicated that
328 broilers incubated under a 12L:12D cycle using green light for the first 18 days not only
329 hatch earlier, but grow faster as significant increases in beak length, third-toe length and
330 CRL were found at some incubation stages. However, overall body weight did not increase.
331 Poultry are usually incubated commercially in complete darkness due to concerns about
332 potential adverse effects of light stimulation on performance and economics, for instance,
333 decreased hatchability due to secondary heating (Archer *et al.*, 2009). Previous published

334 studies have reported different effects of light stimulation during incubation on hatchability
335 due to the spectral characteristics of light and the photoperiod. Archer and Mench (2014)
336 demonstrated that there was no effect on hatchability when different photoperiods were
337 applied using full-spectrum fluorescent light with the intensity of 550 lux throughout
338 incubation. In addition, Zhang *et al.*, (2014) reported no effect of continuous green light of
339 15 lux during incubation on hatchability and hatching. However, in the current study a
340 significant decrease in the hatchability was found in the green light incubation. This may
341 be due to the absence of light-dark cycle from day 19 or the reduced heart weight
342 observed in light exposure embryos. This has not been studied or reported before.
343 Another study showed that the heart rate of chicken embryos responded to the injected
344 melatonin (Höchel and Nichelmann, 2001). Therefore, the changed pattern of melatonin
345 in the light incubation may also affect the cardiac function of chicken embryo. It has been
346 reported that the rhythm of melatonin synthesis in embryos can be synchronised by
347 ambient light-dark (LD) cycles and requires a photoperiod longer than 8 hours for its
348 proper functioning (Zeman *et al.*, 1999). Theoretically, the production of melatonin by the
349 pineal gland is elevated by darkness and suppressed by the presence of light, respectively,
350 which establishes the circadian rhythms. In the current study, although there were no
351 significant differences, there were still different trends in the patterns of plasma MT levels
352 in the 17-18 days old chicken embryos which were incubated under darkness compared
353 to 12L:12D. The plasma MT levels of 12 hours light exposed embryos experienced an
354 increase until 9 hours of darkness when the increase in MT tailed off and subsequently
355 the MT dropped during the light period. This confirms the theory that the light exposure
356 suppressed the MT release during the light period. The significant rhythmic patterns of
357 plasma MT have also been reported on day 19 or day 20 (Zeman *et al.*, 1999, Archer and

358 Mench, 2014) under 12D:12L incubation. Archer and Mench (2014) have demonstrated
359 that the plasma MT levels during the light period at day 19 broilers incubated under
360 12L:12D were almost identical to dark incubated embryos, but it elevated during the dark
361 period while the control birds still maintained at the same level. Their results confirm that
362 the light period is required to trigger the increase during the dark period. Without this dark-
363 light rhythm there is only a somewhat steady MT concentration in the continuous dark at
364 the same level as in the light condition.

365 The rhythm of melatonin release may also influence the rhythms of other hormones
366 (Starck and Ricklefs, 1998). However, little is known of the ontogeny of circadian patterns
367 of secretion of other hormones in birds. A daily rhythm in plasma CORT has only been
368 reported in adult birds and the peak occurs at the transition of dark to light (Sato and
369 George, 1973). The present study showed that plasma CORT concentrations of 17-18
370 days old embryos decreased during the dark period and increased during light period.
371 This indicated that light stimulated the secretion of CORT in the embryo which is in
372 agreement with previous reports that CORT response to light is opposite that of melatonin
373 in broiler and human (Cutolo *et al.*, 2006, de Jong *et al.*, 2001). The underlying mechanism
374 is the ability of MT is able to inhibit the hypothalamic-pituitary-adrenal (HPA) axis and thus
375 suppresses the concentrations of CORT and ultimately thus reduces stress, potentially a
376 significant factor for the rest/sleep phase in the dark period.

377 It has been shown that administration of CORT to chicken embryos was followed by an
378 increase of the plasma T3 levels and T3/T4 ratio (Jenkins and Porter, 2004). In this study,
379 both plasma T3 and CORT showed the same increasing trend in light stimulated embryos
380 conforming that CORT may affect T3 or vice versa through light stimulation. This
381 suggested that thyroid hormone concentrations (T3 & T4) were affected by the circadian

382 rhythms of MT via photoperiod. The different opposed rhythmic patterns of T3 and T4,
383 with plasma T3 concentrations decreasing during the night and increasing during the day
384 and plasma T4 concentrations peaking during the night and lower at day, were also found
385 in chicks during the first week post hatch, and were deemed to be controlled by the feeding
386 pattern (Newcomer, 1974, Klandorf *et al.*, 1978). The same patterns of plasma T3 and T4
387 levels in chicken embryos were also found in our study when they were exposed to a
388 photoperiod of 12D:12L.

389 Light stimulation causing the alteration of hormones may in turn affect hatching
390 behaviours. Our results showed that the concentrations of MT, T3, T4 and CORT
391 increased significantly from the onset of pipping. It has been reported that the high plasma
392 T3 levels occur when the pulmonary respiration initiates (Decuypere and Bruggeman,
393 2007). Here, the T3 levels of embryos increased significantly from IP to EP in both control
394 and light groups. Moreover, the control embryos had higher plasma MT and T4 levels,
395 whereas the light exposed embryos had higher plasma CORT, T3 levels and T3/T4 ratio
396 at EP. These different patterns suggested that the first 18 days light stimulation seems
397 likely to have a consequent impact on hormones levels during hatch. However, all returned
398 to the same levels at take-off. Light exposed chicks hatched about 3.4 h earlier in this
399 study probably due to the elevated levels of CORT, T3 and T3/T4 ratio which are
400 considered important for stimulating a variety of developmental and metabolic process
401 necessary for hatching (Carsia *et al.*, 1987). However, it is still unclear that whether the
402 early hatch is directly related to the change in hormones or the strengthened rhythms.

403 This study was conducted to evaluate the effect of a 12 hours light, 12 hours dark
404 photoperiod of green light during incubation on embryo growth, hatch performance and
405 the hatch process. At the group level, light stimulation had no effect on chick weight and

406 quality at take-off, the initiation of hatch and hatch window, but with a reduced hatchability.
407 However, the individual hatching time of the light stimulated focal chicks was 3.4 hours
408 earlier than the control focal chicks. The results of this study indicated that green light
409 accelerated embryo development and altered plasma MT, CORT, T3 and T4
410 concentrations. Further work is required to understand if and how such altered hormone
411 profiles impact upon hatching time.

412

413 **Acknowledgments**

414 This research is a part of the BioBusiness Project (FP7-PEOPLE-ITN-2008) which is
415 supported by the EU Commission and Marie Curie Initial Training Network. We are
416 grateful to Dr. Yumei Chang for statistical advice and the support of Biological Services
417 Unit at the Royal Veterinary College.

418 **References**

- 419 Archer GS and Mench JA 2014. The effects of the duration and onset of light stimulation during
420 incubation on the behavior, plasma melatonin levels, and productivity of broiler chickens.
421 Journal of Animal Science 92, 1753-1758.
- 422 Archer GS, Shivaprasad HL and Mench JA 2009. Effect of providing light during incubation on
423 the health, productivity, and behavior of broiler chickens. Poultry Science 88, 29-37.
- 424 Barriga C, Marchena JM, Lea RW, Harvey S and Rodriuez AB 2002. Effect of stress and
425 dexamethasone treatment on circadian rhythms of melatonin and corticosterone in ring
426 dove (*Streptopelia risoria*). Molecular and Cellular Biochemistry 232, 27-31.
- 427 Carsia RV, Morin ME, Rosen HD and Weber H 1987. Ontogenic corticosteroidogenesis of the
428 domestic fowl: response of isolated adrenocortical cells. Proceedings of the Society for
429 Experimental Biology and Medicine, New York, 184, 436-445.
- 430 Csernus VJ, Nagy AD and Faluhelyi N 2007. Development of the rhythmic melatonin secretion in
431 the embryonic chicken pineal gland. General and Comparative Endocrinology 152, 148-
432 153.
- 433 Cutolo M, Sulli A., Pizzorni C, Secchi ME, Soldano S, Seriola B, Straub RH, Otsa K and
434 Maestroni GJ 2006. Circadian rhythms - Glucocorticoids and arthritis. Basic and Clinical
435 Aspects of Neuroendocrine Immunology in Rheumatic Diseases 1069, 289-299.
- 436 de Jong IC, Van Voorst AS, Erkens JHF, Ehlhardt D A and Blokhuis HJ 2001. Determination of
437 the circadian rhythm in plasma corticosterone and catecholamine concentrations in
438 growing broiler breeders using intravenous cannulation. Physiology & Behavior 74, 299-
439 304.
- 440 Decuypere E and Bruggeman V 2007. The endocrine interface of environmental and egg factors
441 affecting chick quality. Poultry Science 86, 1037-1042.

442 Duncan IJH, Savory CJ and Wood-Gush DGM 1978. Observations on the reproductive
443 behaviour of domestic fowl in the wild. *Applied Animal Ethology* 4, 29-42.

444 Fairchild BD and Christensen VL 2000. Photostimulation of turkey eggs accelerates hatching
445 times without affecting hatchability, liver or heart growth, or glycogen content. *Poultry*
446 *Science* 79, 1627-1631.

447 Faluhelyi N and Csernus V 2007. The effects of environmental illumination on the in vitro
448 melatonin secretion from the embryonic and adult chicken pineal gland. *General and*
449 *Comparative Endocrinology* 152, 154-158.

450 Halevy O, Piestun Y, Rozenboim I and Yablonka-Reuveni Z 2006. In ovo exposure to
451 monochromatic green light promotes skeletal muscle cell proliferation and affects
452 myofiber growth in posthatch chicks. *American Journal of Physiology Regulatory,*
453 *Integrative Comparative Physiology* 290, 1062-1070.

454 Höchel J and Nichelmann M 2001. Ontogeny of heart rate responses to exogenous melatonin in
455 Muscovy duck and chicken embryos. *Life Science* 69, 2295-2309.

456 Jenkins SA and Porter TE 2004. Ontogeny of the hypothalamo-pituitary-adrenocortical axis in
457 the chicken embryo: a review. *Domestic Animal Endocrinology* 26, 267-275.

458 Klandorf H, Sharp PJ and Duncan IJH 1978. Variations in Levels of Plasma Thyroxine and
459 Triiodothyronine in Juvenile Female Chickens during 24-Hr and 16-Hr Lighting Cycles.
460 *General and Comparative Endocrinology* 36, 238-243.

461 Maurer G, Portugal SJ and Cassey P 2011. Review: an embryo's eye view of avian eggshell
462 pigmentation. *Journal of Avian Biology* 42, 494-504.

463 Mocchegiani E, Perissin L, Santarelli L, Tibaldi A, Zorzet S, Rapozzi V, Giacconi R, Bulian D and
464 Giraldi T 1999. Melatonin administration in tumor-bearing mice (intact and
465 pinealectomized) in relation to stress, zinc, thymulin and IL-2. *International*
466 *Immunopharmacology* 21, 27-46.

467 Newcomer WS 1974. Diurnal Rhythms of Thyroid-Function in Chicks. *General and Comparative*
468 *Endocrinology* 24, 65-73.

469 Nichelmann M, Hochel J and Tzschentke B 1999. Biological rhythms in birds--development,
470 insights and perspectives. *Comparative Biochemistry and Physiology* 124, 429-437.

471 Osorio D, Vorobyev M and Jones CD 1999. Colour vision of domestic chicks. *The Journal of*
472 *Experimental Biology* 202, 2951-2959.

473 Rogers LJ, Andrew RJ and Burne TH 1998. Light exposure of the embryo and development of
474 behavioural lateralisation in chicks, I: olfactory responses. *Behavioural Brain Research*
475 97, 195-200.

476 Romanini CE, Exadaktylos V, Tong Q, McGonnel I, Demmers TG, Bergoug H, Eterradosi N,
477 Roulston N, Garain P, Bahr C and Berckmans D 2013. Monitoring the hatch time of
478 individual chicken embryos. *Poultry Science* 92, 303-309.

479 Rozenboim I, El Halawani ME, Kashash Y, Piestun Y and Halevy O 2013. The effect of
480 monochromatic photostimulation on growth and development of broiler birds. *General*
481 *and Comparative Endocrinology* 190, 214-219.

482 Saito S, Tachibana T, Choi YH, Denbow DM and Furuse M 2005. ICV melatonin reduces acute
483 stress responses in neonatal chicks. *Behavioural Brain Research* 165, 197-203.

484 Sato T and George JC 1973. Diurnal Rhythm of Corticotropin-Releasing Factor Activity in
485 Pigeon Hypothalamus. *Canadian Journal of Physiology and Pharmacology* 51, 743-747.

486 Schultz TF and Kay SA 2003. Circadian clocks in daily and seasonal control of development.
487 *Science* 301, 326-328.

488 Shafey TM, Al-Batshan HA, Ghannam MM and Al-Ayed MS 2005. Effect of intensity of eggshell
489 pigment and illuminated incubation on hatchability of brown eggs. *British Poultry Science*
490 46, 190-198.

491 Shafey TM and Al-Mohsen TH 2002. Embryonic growth, hatching time and hatchability
492 performance of meat breeder eggs incubated under continuous green light. Asian-
493 Australasian Journal of Animal Sciences 15, 1702-1707.

494 Starck JM and Ricklefs RE 1998. Avian growth and development : evolution within the altricial-
495 precocial spectrum. Oxford University Press, London.

496 Tona K, Bamelis F, De Ketelaere B, Bruggeman V, Moraes VM, Buyse J, Onagbesan O and
497 Decuypere E 2003. Effects of egg storage time on spread of hatch, chick quality, and
498 chick juvenile growth. Poultry Science 82, 736-741.

499 Tong Q, MCGonnell IM, Romanini CE, Bergoug H, Roulston N, Exadaktylos V, Berckmans D,
500 Bahr C, Guinebretière M, Etteradossi N, Garain P and Demmers T 2015. Effect of
501 species-specific sound stimulation on the development and hatching of broiler chicks.
502 British Poultry Science 56, 143-148.

503 Yeager R L, Franzosa JA, Millsap DS, Angell-Yeager JL, Heise SS, Wakhungu P, Lim JW,
504 Whelan HT, Eells JT and Henshel DS 2005. Effects of 670-nm phototherapy on
505 development. Photomedicine and Laser Surgery 23, 268-272.

506 Zeman M and Gwinner E 1993. Ontogeny of the Rhythmic Melatonin Production in a Precocial
507 and an Altricial Bird, the Japanese-Quail and the European Starling. Journal of
508 Comparative Physiology a-Sensory Neural and Behavioral Physiology 172, 333-338.

509 Zeman M, Gwinner E, Herichova I, Lamosova D and Kost'al L 1999. Perinatal development of
510 circadian melatonin production in domestic chicks. Journal of Pineal Research 26, 28-34.

511 Zhang L, Zhang HJ, Wang J, Wu SG, Qiao X, Yue HY, Yao JH and Qi GH 2014. Stimulation
512 with monochromatic green light during incubation alters satellite cell mitotic activity and
513 gene expression in relation to embryonic and posthatch muscle growth of broiler
514 chickens. Animal 8, 86-93.

515

516 **Table 1** Mean (\pm SEM) time of onset of internal pipping (IP) and length of hatch window (HW) of
517 the control group and light group from four batches.

Group	IP ^a	HW (hour)
Control	467.3 \pm 0.7	23.0 \pm 1.9
Light	467.3 \pm 0.7	22.3 \pm 1.9
<i>P</i> -value	1.0	0.78

518 ^a hours of incubation time

519 **Figure captions**

520

521 **Figure 1** Mean of hatchability and mortality in the control group and the light group over four
522 batches (n=1200 eggs at setting of each group, fertility varied from 82-96%). ED, early death from
523 day 0 to day 7; MD, middle death from day 8 to day 15; LD, late death from day 16 to day 21.
524 *Means significantly different between the control and light groups at $P<0.05$.

525

526 **Figure 2** The heart weight, beak length, third toe length and crown-rump length of the control
527 and light exposed embryos or chicks from day 10 to day 21 of incubation time. Dashed lines
528 indicate the control group and solid lines indicate the light group. Asterisk indicates significant
529 difference between groups at a given incubation stage ($*P<0.05$; $**P<0.01$). Data are presented
530 as mean \pm SEM (12 samples of each group at each incubation stage).

531

532 **Figure 3** Plasma melatonin concentrations over an 18 hours period from day 17 to day 18,
533 internal pipping (IP) at day 19, external pipping (EP) at day 20 and take-off at day 21 of broilers
534 incubated under the control and light groups. Solid line indicates the light group with a
535 photoperiod of 12 h of light and 12 h of darkness for the first 18 days of incubation and no light-
536 dark cycle for the last 3 days of incubation; the horizontal bars indicate the dark (closed) and
537 light (open) periods of the photoperiod in the light group. Dashed line indicates the control group
538 where eggs were incubated under darkness throughout the incubation. Data are presented as
539 mean \pm SEM of four samples and each was pooled blood from three embryos or chicks at each
540 time point. ^{ab/xy} Means significant difference among the given time points within a group
541 ($P<0.05$).

542

543 **Figure 4** Plasma corticosterone concentrations over an 18 hours period from day 17 to day 18,
544 internal pipping (IP) at day 19, external pipping (EP) at day 20 and take-off at day 21 of broilers
545 incubated under the control and light groups. Solid line indicates the light group with a
546 photoperiod of 12 h of light and 12 h of darkness for the first 18 days of incubation and no light-
547 dark cycle for the last 3 days of incubation; the horizontal bars indicate the dark (closed) and
548 light (open) periods of the photoperiod in the light group. Dashed line indicates the control group
549 where eggs were incubated under darkness throughout the incubation. Data are presented as
550 mean \pm SEM of four samples and each was pooled blood from three embryos or chicks at each
551 time point. Asterisk indicates significant difference between groups at a given time point (**
552 $P < 0.01$).

553
554 **Figure 5** Plasma Triiodothyronine (T3) concentrations over an 18 hours period from day 17 to
555 day 18, internal pipping (IP) at day 19, external pipping (EP) at day 20 and take-off at day 21 of
556 broilers incubated under the control and light groups. Solid line indicates the light group with a
557 photoperiod of 12 h of light and 12 h of darkness for the first 18 days of incubation and no light-
558 dark cycle for the last 3 days of incubation; the horizontal bars indicate the dark (closed) and
559 light (open) periods of the photoperiod in the light group. Dashed line indicates the control group
560 where eggs were incubated under darkness throughout the incubation. Data are presented as
561 mean \pm SEM of four samples and each was pooled blood from three embryos or chicks at each
562 time point. ^{abc/xyz} Means significant difference among the given time points within a group
563 ($P < 0.05$).


564
565 **Figure 6** Plasma Thyroxine (T4) concentrations over an 18 hours period from day 17 to day 18,
566 internal pipping (IP) at day 19, external pipping (EP) at day 20 and take-off at day 21 of broilers
567 incubated under the control and light groups. Solid line indicates the light group with a
568 photoperiod of 12 h of light and 12 h of darkness for the first 18 days of incubation and no light-

569 dark cycle for the last 3 days of incubation; the horizontal bars indicate the dark (closed) and
570 light (open) periods of the photoperiod in the light group. Dashed line indicates the control group
571 where eggs were incubated under darkness throughout the incubation. Data are presented as
572 mean \pm SEM of four samples and each was pooled blood from three embryos or chicks at each
573 time point. ^{ab} Means significant difference among the given time points within a group ($P < 0.05$).


574
575 **Figure 7** Plasma T3/T4 ratio over an 18 hours period from day 17 to day 18, internal pipping (IP)
576 at day 19, external pipping (EP) at day 20 and take-off at day 21 of broilers incubated under the
577 control and light groups. Solid line indicates the light group with a photoperiod of 12 h of light
578 and 12 h of darkness for the first 18 days of incubation and no light-dark cycle for the last 3 days
579 of incubation; the horizontal bars indicate the dark (closed) and light (open) periods of the
580 photoperiod in the light group. Dashed line indicates the control group where eggs were
581 incubated under darkness throughout the incubation. Data are presented as mean \pm SEM of four
582 samples and each was pooled blood from three embryos or chicks at each time point. ^{ab/xy}
583 Means significant difference among the given time points within a group ($P < 0.05$).

584


585 Figure 1




586 Figure 2




587 Figure 3




588 Figure 4



589 Figure 5



590 Figure 6



591 Figure 7

