

This is the peer reviewed version of the following article:

Lamb, C. R., Pope, E. H.W. and Lee, K. C.L. (2016), RESULTS OF COMPUTED TOMOGRAPHY IN DOGS WITH SUSPECTED WOODEN FOREIGN BODIES. Vet Radiol Ultrasound. doi:[10.1111/vru.12457](https://doi.org/10.1111/vru.12457)

This article may be used for non-commercial purposes in accordance with [Wiley Terms and Conditions for Self-Archiving](#).

The full details of the published version of the article are as follows:

TITLE: RESULTS OF COMPUTED TOMOGRAPHY IN DOGS WITH SUSPECTED WOODEN FOREIGN BODIES

AUTHORS: Christopher R. Lamb, Elizabeth H.W. Pope, Karla C.L. Lee

JOURNAL TITLE: Veterinary Radiology & Ultrasound

PUBLISHER: Wiley

PUBLICATION DATE: November 2016 (online)

DOI: 10.1111/vru.12457

1 **Results of computed tomography in dogs with suspected wooden foreign bodies**

2 Christopher R. Lamb, Elizabeth H.W. Pope, Karla C.L. Lee

3 From the Department of Clinical Science and Services, The Royal Veterinary College, University of  
4 London.

5 Address correspondence to: C.R. Lamb, Department of Clinical Science and Services, The Royal  
6 Veterinary College, Hawkshead Lane, North Mymms, Hertfordshire AL9 7TA, U.K. Email:  
7 [clamb@rvc.ac.uk](mailto:clamb@rvc.ac.uk)

8

9 Sources of funding: unfunded

10 Conflicts of interest: none declared

11 Key words: computed tomography, dog, foreign bodies, penetrating injuries, wood

12 Running head: CT of wooden foreign bodies

**13 Abstract**

14 Detection of wooden foreign bodies in dogs can be challenging. A retrospective, cross-sectional study  
15 was done to describe computed tomographic (CT) signs associated with wooden foreign bodies, and to  
16 estimate the accuracy of CT for detection of wooden foreign bodies. Patient records and CT images were  
17 reviewed of 72 dogs that had a history of suspected stick injury and CT of the affected body part, or  
18 possible wooden foreign object reported on CT, and had surgical exploration during the same period of  
19 hospitalization. Duration of clinical signs was acute in 48 (67%) dogs and chronic in 24 (33%). Wood was  
20 removed from 55 dogs, including a piece of a tree or shrub in 33 (60%) instances, kebab stick in 8 (15%),  
21 piece of bamboo garden cane in 2 (4%), cocktail stick in 2 (4%), thorn in 1 (2%), and unidentified wood in  
22 the remaining 9 instances. Based on review of CT images with knowledge of the surgical findings,  
23 sensitivity of CT for wooden foreign bodies was 79% (95% CI 65-89%), specificity 93% (78-98%), positive  
24 likelihood ratio 11.5 (2.9- 44.1), and negative likelihood ratio 0.23 (0.13-0.41). Wooden foreign bodies  
25 were predominantly rectangular or linear, with median length 48mm (range 2-270mm), median  
26 thickness 3mm (range 1-22mm), and median attenuation 111HU (range <sup>-</sup>344-<sup>+</sup>640HU). A CT finding of  
27 gas in soft tissues was significantly associated with acute cases, whereas suspected foreign material,  
28 cavitory lesions, fat stranding, and periosteal reaction on adjacent bones were associated with chronic  
29 cases.

## 30 Introduction

31 Wooden stick injuries are a well-recognized problem in dogs, frequently occurring because of dogs  
32 chasing and retrieving wooden sticks and sustaining oropharyngeal penetration injuries.<sup>1-4</sup>  
33 Oropharyngeal stick injuries frequently cause acute signs including pain, blood in the saliva, and  
34 dysphagia. In such cases, surgical exploration and lavage of the wound and removal of foreign material  
35 usually resolves the problem.<sup>1, 3, 4</sup> When the initial trauma is not recognized or treated and stick injuries  
36 become chronic, an abscess and/or discharging sinus may develop.<sup>5-8</sup> Stick foreign bodies can also reach  
37 sites distant from the point of penetration by migration through tissues, which complicates diagnosis  
38 and management.<sup>9-12</sup> Sticks, wooden fragments and other wooden foreign bodies have been reported in  
39 many anatomic sites in dogs, including the orbit, larynx, cervical region, axilla, thorax, abdomen, and  
40 limbs.<sup>8, 11-17</sup>

41 Detection of wooden foreign bodies in dogs can be challenging. In dogs with acute oropharyngeal  
42 injuries, radiography is indicated to look for gas in tissues, which indicates that perforation of the  
43 pharyngeal wall or esophagus has occurred, and retained pieces of wood. More comprehensive imaging  
44 may be necessary because of the possibility of retained foreign material in dogs with persistent or  
45 recurrent signs of localized infection, unexplained pleuritis or peritonitis, suspected abscess or draining  
46 sinus. Radiography<sup>3, 6</sup>, sinography<sup>7, 18</sup>, ultrasonography<sup>10, 19, 20</sup>, computed tomography (CT)<sup>21-26</sup>, and  
47 magnetic resonance (MR) imaging<sup>6, 23, 27, 28</sup> have been used in dogs with suspected penetrating or  
48 migrating foreign bodies. The choice of modality in any individual case will depend on various factors  
49 including site of injury, type of suspected foreign body, and availability. In humans with wooden foreign  
50 bodies, different reports have emphasized ultrasound<sup>29, 30</sup>, CT<sup>31-33</sup> and MR<sup>34</sup> without a clear priority.  
51 Computed tomography is preferred in many instances because it enables an efficient, cross-sectional  
52 examination of any part of the body with relatively high spatial resolution. In a canine cadaver study,  
53 non-contrast CT was more accurate (sensitivity 75%, specificity 96%) than ultrasound or MR imaging for

54 identification of wooden foreign bodies in the manus.<sup>35</sup> Existing reports describe a variety of  
55 appearances for wood in CT images, including hypo- or hyperattenuating foci compared to adjacent soft  
56 tissues, structures that contain gas or have gas on their surface, with shapes, such as angular or linear,  
57 that do not conform to normal anatomy<sup>21-26,35</sup>; however, there is a lack of data regarding the accuracy of  
58 CT for wooden foreign bodies in live veterinary patients.

59 The aims of the present study were to describe the clinical and CT signs associated with wooden foreign  
60 bodies in a series of dogs, to test the hypothesis that signs differ between acute and chronic cases, and  
61 to estimate the accuracy of CT for detection of wooden foreign bodies.

62

### 63 **Methods**

64 For this retrospective cross-sectional study, computerized patient records at The Queen Mother Hospital  
65 for Animals in the period 2005-2015 were searched (by EHWP) for dogs that had a history of suspected  
66 stick injury and CT of the affected body part, or possible wooden foreign object reported on CT, and  
67 surgical exploration during the same period of hospitalization. Data extracted from records included  
68 signalment, history, duration and nature of clinical signs, results of CT, results of surgery, and type of  
69 wood retrieved (where known).

70 For the purposes of this study, CT images were reviewed by a board-certified radiologist (CRL) with  
71 knowledge of the surgical findings. Images were reviewed using soft tissue (level 50HU, width 350HU)  
72 and lung (level -500HU, width 2000HU) windows. Oblique multiplanar reconstructions were used as  
73 required on an individual case basis. When suspected foreign material was identified, its maximal short  
74 and long axis dimensions were determined using the measuring tool in a proprietary DICOM viewer  
75 (Image Viewer Version 4.1, Visbion Ltd, Chertsey, UK) and its attenuation in Hounsfield units (HU) was  
76 measured using a hand-drawn region of interest fitted to the cross-section of the object. Within a

77 suspected foreign body, zones of variable density were included in the region of interest, but if a cavity  
78 appeared gas-filled because its HU values were comparable to air surrounding the patient, it was not  
79 included. The average of three HU measurements was recorded.

80 Clinical data were summarized using median and range. Cases with duration of clinical signs prior to  
81 referral <7 days were classified as acute and those with duration of 7 or more days were classified as  
82 chronic.<sup>4</sup> Statistical testing was done by CRL. Differences in prevalence of clinical signs in acute versus  
83 chronic cases were tested using Fisher's exact test. The difference in median attenuation values of wood  
84 found in acute versus chronic cases was tested using the Mann-Whitney test. (SPSS Statistics V22, IBM  
85 Corporation, New York, USA). Differences with  $p < 0.05$  were considered significant. Binomial 95%  
86 confidence intervals (CI) for estimates of sensitivity, specificity, positive likelihood ratio (PLR), and  
87 negative likelihood ratio (NLR) were determined using the online statistical calculator provided by the  
88 Center for Evidence Based Medicine (<http://ktclearinghouse.ca/cebmlpractise/ca/calculators/statscalc>).

89

## 90 **Results**

91 Records were found of 72 dogs that satisfied the inclusion criteria. No dogs were excluded. There were  
92 51 male dogs (35 neutered) and 21 females (17 neutered). The median age of the dogs was 3.9 years  
93 (range 9 months-11.9 years). Their median body weight was 25.0kg (range 6.8-50kg). The most  
94 prevalent breeds were 20 (28%) Labrador retrievers, 12 (17%) mixed breed, 11 (15%) English Springer  
95 Spaniels 7 (10%) Border Collies, and 5 (7%) Staffordshire bull terriers. In addition, there were 1 or 2 dogs  
96 of 14 other canine breeds.

97 A single slice helical scanner (PQ5000, Universal Medical Systems, Solon, OH) was used for CT before  
98 August 2009 and a 16-slice helical scanner (Mx8000 IDT, Philips, Best, The Netherlands) was used from  
99 August 2009. CT settings were optimized for examination of soft tissues. Settings included helical

100 acquisition, slice thickness 2-5mm, 120-250mAs, 120kVp, field of view 240-400mm, matrix 512x512, and  
101 medium frequency ('soft tissue') reconstruction algorithm. Pre- and post-contrast CT series were  
102 obtained in all patients. For post-contrast series, 2ml/kg of iohexol 300mg/ml (Omnipaque 300, GE  
103 Healthcare, Oslo, Norway) was injected intravenously either by hand (for small dogs) or at 2ml/s using a  
104 power injector (Stellant, Medrad Inc., Pennsylvania, USA), and the acquisition initiated at 60 seconds  
105 after the start of intravenous injection.

106 In 35 (48%) instances the owners reported observing trauma that caused a penetration injury, and in 21  
107 (29%) instances a piece of wood had been removed from a wound prior to referral. Duration of clinical  
108 signs prior to referral was <7 days in 48 (67%) dogs and 7 or more days in 24 (33%) dogs. In addition to  
109 the wound caused by stick entry, which was evident in 35 (49%) dogs, the most prevalent clinical signs  
110 were soft tissue swelling in 25 (35%) dogs, signs of pain on palpation in 20 (28%), pyrexia (rectal  
111 temperature >39.1C) in 15 (21%) dogs, and lethargy in 8 (11%). Prevalence of clinical signs observed in at  
112 least 2 dogs is summarized in table 1. A wound was significantly associated with acute cases ( $p=0.001$ ),  
113 whereas a draining sinus ( $p=0.01$ ) and vomiting ( $p=0.03$ ) were significantly associated with chronic  
114 cases.

115 Of the 72 dogs included in this study, 50 (69%) had history of trauma and/or clinical signs of a  
116 penetrating wound. In these dogs, lesions were found in the cervical region in 20 (40%) dogs, the oral  
117 cavity in 12 (24%), pharynx in 12 (24%), thorax in 5 (10%), and inguinal region in 1 (2%). In contrast, in  
118 the 22 (31%) dogs with no history of trauma nor clinical signs of a penetrating wound, lesions were  
119 found in the abdomen in 10 (46%) dogs, cervical region in 5 (23%), thorax in 4 (18%), pharynx in 2 (9%),  
120 and nasal cavity in 1 (5%).

121 Over the course of their veterinary care, wood was removed from 55 dogs, including 21 in which  
122 removal was reported to have occurred prior to referral. Wood was identified as a piece of a branch of a

123 tree or shrub in 33 (60%) instances, kebab stick in 8 (15%), piece of bamboo garden cane in 2 (4%),  
124 cocktail stick in 2 (4%), thorn in 1 (2%) and was unidentified in the remaining 9 (16%) instances.

125 Evidence of a wooden foreign body was identified in CT images in 36 dogs; however, after CT, fragments  
126 of wood or sticks were removed surgically from 43 dogs (including 9 dogs from which another piece of  
127 wood had already been removed from a wound prior to referral). Wood was removed surgically from  
128 the cervical region in 18 (42%) dogs, pharynx in 7 (16%), peritoneal cavity in 7 (16%), lung in 4 (9%),  
129 thoracic wall in 3 (7%), nasal cavity in 1 (2%), stomach in 1 (2%), duodenum in 1 (2%), and inguinal  
130 region in 1 (2%). All of the kebab sticks and cocktail sticks were retrieved from the thoracic wall or  
131 abdomen.

132 In 2 dogs, a suspected wooden foreign body was identified in CT images but no foreign material was  
133 found at exploratory surgery, hence these were classified as false positive results. In one of these dogs,  
134 clinical follow-up over a period exceeding 2 years found no persistent or recurrent signs, supporting the  
135 conclusion that there was no retained foreign material. The other dog, which had chronic immune-  
136 mediated anemia and signs of a wooden foreign body detected unexpectedly at abdominal CT, collapsed  
137 and died 4 weeks after discharge, and no necropsy was performed.

138 Classification of CT results with respect to presence/absence of wooden foreign bodies was true positive  
139 34, false positive 2, true negative 27, and false negative 9. Sensitivity of CT for wooden foreign bodies  
140 was 79% (95% CI 65-89%), specificity was 93% (78-98%), PLR was 11.5 (2.9- 44.1), and NLR was 0.23  
141 (0.13-0.41).

142 In CT images, the 34 wooden foreign bodies confirmed as true positives were predominantly rectangular  
143 or linear, with their length greatly exceeding their width. The median length of wooden foreign bodies  
144 was 48mm (range 2-270mm) and the median thickness was 3mm (range 1-22mm). The median  
145 attenuation of wooden foreign bodies was 111HU (range -344-+640HU). In 6/34 (18%) instances, the



146 foreign material contained linear variations in density compatible with wood structure (Figure 1). A  
147 median attenuation value  $<0\text{HU}$  was observed in CT images of 6 wooden foreign bodies found in acute  
148 cases, but in none of the chronic cases. The difference in median attenuation of wooden foreign bodies  
149 between dogs with acute versus chronic histories was of borderline statistical significance ( $p=0.05$ ).  
150 CT signs observed in dogs in this series are summarized in table 2. A CT finding of gas in soft tissues was  
151 significantly associated with acute cases ( $p<0.001$ ), whereas suspected foreign material ( $p<0.001$ ),  
152 cavitory lesions ( $p=0.002$ ), fat stranding ( $p=0.01$ ), and periosteal reaction on adjacent bones ( $p=0.01$ )  
153 were significantly associated with chronic cases (Figure 2).

154

## 155 **Discussion**

156 The predominance of lesions affecting the oral cavity, pharynx and cervical region in dogs with history  
157 and/or clinical signs of a traumatic injury in this study corresponds with the belief that stick injuries  
158 frequently occur when dogs run onto a stick or collide with a tree or shrub.<sup>1-4</sup> Conversely, the current  
159 findings that none of the dogs with sticks in the abdomen had a history of trauma, and that the kebab  
160 sticks and cocktail sticks were all retrieved from the thoracic wall or abdomen, suggest that these sticks  
161 were ingested, probably because at the time of ingestion they had food attached.<sup>9,11</sup>

162 Just as the etiology of wooden foreign bodies can be divided into penetrating trauma and ingestion, the  
163 clinical and imaging findings can be divided into those associated with acute and chronic presentations.<sup>4</sup>

164 Clinical signs of a wound and CT signs of gas in soft tissues were significantly associated with acute  
165 cases, whereas a draining sinus and CT findings of cavitory lesions, fat stranding, and periosteal reaction  
166 on adjacent bones were significantly associated with chronic cases. Visualization of the wooden foreign  
167 body by CT was possible in a larger proportion of chronic cases than acute cases in the present study.

168 This observation may reflect multiple factors, such as removal of obvious penetrating foreign bodies

169 prior to referral for imaging, and the difficulty distinguishing by CT small low-density wood fragments  
170 from gas introduced by a penetrating injury. Although not specifically tested in the present study,  
171 authors propose that post-contrast CT images would likely be most useful as an aid to detection of  
172 cavitory lesions, such as abscess, rather than as a means of detecting foreign material directly, as  
173 suggested previously.<sup>22</sup> This suggests that, careful scrutiny of initial, non-contrast CT images is advisable  
174 because of the possibility that subsequent contrast accumulation in tissues could obscure a small  
175 hyperdense foreign object.

176 Confirmed wooden foreign bodies were found to have a wide range of attenuation values in CT images (  
177 344-+640HU). This variation likely reflects normal differences in the native density of wood (depending  
178 on the plant species, its maturity, and condition prior to entry into the body) and changes that may  
179 occur as a result of absorption of body fluids by porous wood.<sup>34,36</sup> Wood with median attenuation value  
180 <0HU was observed in CT images of dogs with acute histories, but not in dogs with chronic histories.  
181 Although of borderline statistical significance, this difference is compatible with the hypothesis that  
182 there is an increase over time in attenuation of wood lodged in tissues.<sup>34,36</sup> When a piece of wood is  
183 markedly hyperdense compared to surrounding tissues, it is liable to be recognized easily; however, low-  
184 density pieces of wood resemble gas in CT images obtained using a typical soft tissue window, and  
185 hence may be missed, particularly in acute cases when a penetrating wound has allowed gas into the  
186 tissues.<sup>36</sup> When examining a wound containing gas, use of a wide window is recommended to help  
187 distinguish gas from low-density wood fragments.<sup>33, 36, 37</sup> Similarly, use of multiplanar reconstructions  
188 aids CT identification of foreign bodies.<sup>26</sup> Even large pieces of wood may be difficult to recognize when  
189 orientated obliquely to the image plane (Figure 3).

190 In the present study, the sensitivity of CT for wooden foreign bodies was 79% and specificity was 93%.  
191 These results are similar to those of the previously reported canine cadaver study that employed non-  
192 contrast CT.<sup>35</sup> PLR was 11.5 and NLR was 0.23 indicating that in a dog with a suspected wooden foreign

193 body, both positive and negative results for CT significantly affect the probability of diagnosis; however,  
194 these estimates cannot be considered definitive because of limitations associated with the study design.  
195 For example, dogs that did not undergo surgical exploration because foreign body was not considered in  
196 the differential diagnosis or not recognized by CT would not be included. This omission would contribute  
197 to over-estimation of the sensitivity of CT. Similarly, it is possible that some dogs with wooden foreign  
198 bodies were misclassified as negatives because the object was not retrieved surgically. This error would  
199 lead to over-estimation of the sensitivity and/or under-estimation of the specificity of CT. Clinical  
200 suspicion of foreign material and careful review of the CT images in collaboration with the surgeon  
201 should help minimize this error, but will not eliminate it. Images were reviewed with knowledge of the  
202 surgical findings in order to ensure optimal search for foreign material, but under this circumstance  
203 sensitivity is likely to be higher than when images are reviewed during case management. Based on  
204 previous publications, it is clear that CT has a much higher sensitivity than radiography, which fails to  
205 demonstrate wooden foreign bodies in many instances.<sup>10,20, 29, 35</sup> In the present study, direct comparison  
206 with sensitivity of other modalities was not possible because few dogs had other imaging studies.  
207 On the basis of these findings, CT may be recommended as a technique with moderate sensitivity and  
208 high specificity for detection of wooden foreign bodies in dogs following penetrating injury or ingestion.

209

## 210 **Acknowledgement**

211 We thank our nursing and veterinary colleagues who contributed to the management of the cases  
212 described in this study.

213

## 214 **List of Author Contributions**

215 Category 1

- 216 (a) Conception and Design
- 217 Author name (s) C.R. Lamb, E.H.W. Pope, K.C.L. Lee
- 218 (b) Acquisition of Data
- 219 Author name (s) C.R. Lamb, E.H.W. Pope, K.C.L. Lee
- 220 (c) Analysis and Interpretation of Data
- 221 Author name (s) C.R. Lamb, E.H.W. Pope, K.C.L. Lee
- 222
- 223 Category 2
- 224 (a) Drafting the Article
- 225 Author name (s) C.R. Lamb, E.H.W. Pope, K.C.L. Lee
- 226 (b) Revising Article for Intellectual Content
- 227 Author name (s) C.R. Lamb, E.H.W. Pope, K.C.L. Lee
- 228
- 229 Category 3
- 230 (a) Final Approval of the Completed Article
- 231 Author name (s) C.R. Lamb, K.C.L. Lee

232 **References**

- 233 1. Devriendt N, Van Goethem B, Kitshoff A, Furcas A, Van der Vekens E, de Rooster H. Oropharyngeal  
234 stick injury in the dog. *Vlaams Diergeneeskundig Tijdschrift* 2015;84: 162-168.
- 235 2. Doran IP, Wright CA, Moore AH. Acute oropharyngeal and esophageal stick injury in forty-one  
236 dogs. *Veterinary Surgery* 2008;37: 781-785.
- 237 3. Griffiths LG, Tiruneh R, Sullivan M, Reid SWJ. Oropharyngeal penetrating injuries in 50 dogs: A  
238 retrospective study. *Veterinary Surgery* 2000;29: 383-388.
- 239 4. White RAS, Lane JG. Pharyngeal stick penetration injuries in the dog. *Journal of Small Animal  
240 Practice* 1988;29: 13-35.
- 241 5. Baglietto M, Cloquell A, Monteagudo S, Munoz A, Mateo I. Spinal epidural empyema associated  
242 with paravertebral and sublumbar abscesses in two dogs. Diagnosis with computed tomography  
243 and myelography. *Clinica Veterinaria De Pequeños Animales* 2011;31: 85-90.
- 244 6. Frenidin J, Funkquist B, Hansson K, Lonnemark M, Carlsten J. Diagnostic imaging of foreign body  
245 reactions in dogs with diffuse back pain. *Journal of Small Animal Practice* 1999;40: 278-285.
- 246 7. Lamb CR, White RN, McEvoy FJ. Sinography in the investigation of draining tracts in small animals  
247 - retrospective review of 25 cases. *Veterinary Surgery* 1994;23: 129-134.
- 248 8. Moon SJ, Lee JH, Jeong SW, Kim JW, Park HM. Chronic bronchocutaneous fistula caused by  
249 toothpick foreign body in a Maltese dog. *Journal of Veterinary Medical Science* 2012;74: 651-655.
- 250 9. Stander N, Kirberger RM. Diagnostic imaging of migrating kebab (sosatie) sticks - a review of 8  
251 cases. *Journal of the South African Veterinary Association* 2011;82: 160-165.

- 252 10. Thiel C, Frese H, Tacke S, Herde K, Kramer M. Injuries through wooden foreign bodies in dogs. A  
253 retrospective study of frequently underestimated wounds. *Tieraerztliche Praxis Ausgabe Kleintiere*  
254 *Heimtiere* 2006;34: 157-167.
- 255 11. Hunt GB, Worth A, Marchevsky A. Migration of wooden skewer foreign bodies from the  
256 gastrointestinal tract in eight dogs. *Journal of Small Animal Practice* 2004;45: 362-367.
- 257 12. Sereda NC, Towl S, Maisenbacher HW, 3rd, Bleweis MS, Levy JK, Byrne BJ, et al. Intracardiac  
258 foreign body in a dog. *Journal of Veterinary Cardiology* 2009;11: 53-58.
- 259 13. O'Reilly A, Beck C, Mouatt JG, Stenner VJ. Exophthalmos due to a wooden foreign body in a dog.  
260 *Australian Veterinary Journal* 2002;80: 268-271.
- 261 14. Mueller M, Thiel C, Mildner-Mueller K, Hartmann A, Hassdenteufel E, Kramer M. Penetrating  
262 laryngeal injury caused by a wooden foreign body in a crossbreed bitch. *Kleintierpraxis* 2013;58:  
263 512-522.
- 264 15. Pelosi A, Hauptman JG, Eyster GE, Beal MW, Anderson LK, Olivier NB. Myocardial perforation by a  
265 stick foreign body in a dog. *Journal of Veterinary Emergency and Critical Care* 2008;18: 184-187.
- 266 16. Clarke SP, Ferguson JF. Bacterial infective arthritis following a penetrating stick injury of the stifle  
267 joint in a dog. *Journal of Small Animal Practice* 2012;53: 483-486.
- 268 17. Menard J, Schoeffler GL. Colonic, ureteral, and vascular injuries secondary to stick impalement in  
269 a dog. *Journal of Veterinary Emergency and Critical Care* 2011;21: 387-394.
- 270 18. Stuehrenberg W, Heinrich M. Oropharyngeal penetrating injury in a mixed-breed male dog -  
271 fistulography as a simple alternative in imaging wound sinuses and foreign bodies. *Kleintierpraxis*  
272 2010;55: 423-428.

- 273 19. Penninck D, Mitchell SL. Ultrasonographic detection of ingested and perforating wooden foreign  
274 bodies in four dogs. *Journal of the American Veterinary Medical Association* 2003;223: 206-209.
- 275 20. Armbrust LJ, Biller DS, Radlinsky MG, Hoskinson JJ. Ultrasonographic diagnosis of foreign bodies  
276 associated with chronic draining tracts and abscesses in dogs. *Veterinary Radiology & Ultrasound*  
277 2003;44: 66-70.
- 278 21. Hinken K, Kaiser S, Hinken M. Computed tomographic imaging of plant foreign bodies associated  
279 with oropharyngeal injuries. *Kleintierpraxis* 2010;55: 5-13.
- 280 22. Nicholson I, Halfacree Z, Whatmough C, Mantis P, Baines S. Computed tomography as an aid to  
281 management of chronic oropharyngeal stick injury in the dog. *Journal of Small Animal Practice*  
282 2008;49: 451-457.
- 283 23. Woolfson JM, Wesley RE. Magnetic-resonance-imaging and computed tomographic scanning of  
284 fresh (green) wood foreign-bodies in dog orbits. *Ophthalmic Plastic and Reconstructive Surgery*  
285 1990;6: 237-240.
- 286 24. Appleby R, zur Linden A, Singh A, Finck C, Crawford E. Computed tomography diagnosis of a  
287 thoracic and abdominal penetrating foreign body in a dog. *Canadian Veterinary Journal* 2015;56:  
288 1149-1152.
- 289 25. Bouabdallah R, Moissonnier P, Delisle F, De Fornel P, Manassero M, Maaoui M, et al. Use of  
290 preoperative computed tomography for surgical treatment of recurrent draining tracts. *Journal of*  
291 *Small Animal Practice* 2014;55: 89-94.
- 292 26. Jones JC, Ober CP. Computed tomographic diagnosis of non-gastrointestinal foreign bodies in  
293 dogs. *Journal of the American Animal Hospital Association* 2007;43: 99-111.

- 294 27. Dobromylskyj MJ, Dennis R, Ladlow JF, Adams VJ. The use of magnetic resonance imaging in the  
295 management of pharyngeal penetration injuries in dogs. *Journal of Small Animal Practice* 2008;49:  
296 74-79.
- 297 28. Young B, Klopp L, Albrecht M, Kraft S. Imaging diagnosis: Magnetic resonance imaging of a cervical  
298 wooden foreign body in a dog. *Veterinary Radiology & Ultrasound* 2004;45: 538-541.
- 299 29. Peterson JJ, Bancroft LW, Kransdorf MJ. Wooden foreign bodies: imaging appearance. *American*  
300 *Journal of Roentgenology* 2002;178: 557-562.
- 301 30. Jacobsen JA, Powell A, Craig JG, Bouffard JA, van Holsbeeck MT. Wooden foreign bodies in soft  
302 tissue: detection at US. *Radiology* 1998;206: 45-48.
- 303 31. Boncoeur-Martel MP, Adenis JP, Rulfi JY, Robert PY, Dupuy JP, Maubon A. CT appearances of  
304 chronically retained wooden intraorbital foreign bodies. *Neuroradiology* 2001;43: 165-168.
- 305 32. Uchino A, Kato A, Takase Y, Kudo S. Intraorbital wooden and bamboo foreign bodies: CT.  
306 *Neuroradiology* 1997;39: 213-215.
- 307 33. Yoon JH, Kim SH, Lee Y, Kim OH, Ryu JH, Lee KH, et al. Detection of an accidentally implanted  
308 wooden foreign body using CT: case report and literature review. *Clinical Imaging* 2015;39: 158-  
309 160.
- 310 34. Imokawa H, Tazawa T, Sugiura N, Oyake D, Yosino K. Penetrating neck injuries involving wooden  
311 foreign bodies: the role of MRI and the misinterpretation of CT images. *Auris Nasus Larynx*  
312 2003;30: S145-S147.
- 313 35. Ober CP, Jones JC, Larson MM, Lanz OI, Werre SR. Comparison of ultrasound, computed  
314 tomography, and magnetic resonance imaging in detection of acute wooden foreign bodies in the  
315 canine manus. *Veterinary Radiology & Ultrasound* 2008;49: 411-418.



- 316 36. Krimmel M, Cornelius CP, Stojadinovic S, Hoffmann J, Reinert S. Wooden foreign bodies in facial  
317 injury: a radiological pitfall. *International Journal of Oral and Maxillofacial Surgery* 2001;30: 445-  
318 447.
- 319 37. Dalley RW. Intraorbital wood foreign-bodies on CT - use of wide bone window settings to  
320 distinguish wood from air. *American Journal of Roentgenology* 1995;164: 434-435.
- 321

322 Table 1. Prevalence of clinical signs in dogs with suspected wooden stick foreign bodies<sup>†</sup>

323	Clinical Sign	Number of affected dogs		
324		Acute cases	Chronic cases	Total
325		(n=48)	(n=24)	(n=72)
326				
327	Wound caused by stick entry	31 (65%) <sup>a</sup>	4 (17%) <sup>a</sup>	35 (49%)
328	Soft tissue swelling	17 (35%)	11 (46%)	28 (39%)
329	Pain on palpation	17 (35%)	3 (13%)	20 (28%)
330	Pyrexia	8 (17%)	7 (29%)	15 (21%)
331	Lethargy	4 (8%)	4 (17%)	8 (11%)
332	Dyspnea	4 (8%)	1 (4%)	5 (7%)
333	Tachypnoea	3 (6%)	0	3 (4%)
334	Epistaxis	2 (4%)	0	2 (3%)
335	Inappetence	2 (4%)	2 (8%)	4 (6%)
336	Ptyalism	2 (4%)	1 (4%)	3 (4%)
337	Stiff neck	2 (4%)	0	2 (3%)
338	Retching	2 (4%)	0	2 (3%)
339	Cyanosis	2 (4%)	0	2 (3%)
340	Coughing	1 (2%)	2 (8%)	3 (4%)
341	Draining sinus	0 <sup>b</sup>	4 (17%) <sup>b</sup>	4 (6%)
342	Vomiting	0 <sup>c</sup>	3 (13%) <sup>c</sup>	3 (4%)
343	Diarrhea	0	2 (8%)	2 (3%)
344				

---

<sup>†</sup> Values with same superscript are significantly different, p<0.05



346 Table 2. Prevalence of CT signs in dogs with suspected wooden stick foreign bodies<sup>‡</sup>

347 Sign	348 Number of affected dogs		
	349 Acute cases (n=48)	Chronic cases (n=24)	Total (n=72)
351 Gas in soft tissues	46 (96%) <sup>a</sup>	3 (13%) <sup>a</sup>	49 (68%)
352 Soft tissue swelling	21 (44%)	17 (71%)	38 (53%)
353 Enlarged regional nodes	20 (42%)	12 (50%)	32 (44%)
354 Suspected foreign material	16 (33%) <sup>b</sup>	20 (83%) <sup>b</sup>	36 (50%)
355 Pneumothorax	9 (19%)	0	9 (13%)
356 Cavitory lesion	7 (15%) <sup>c</sup>	12 (50%) <sup>c</sup>	19 (26%)
357 Fat stranding	5 (10%) <sup>d</sup>	9 (38%) <sup>d</sup>	14 (19%)
358 Intra-cavity fluid	3 (15%)	3 (50%)	6 (26%)
359 Periosteal reaction	0 <sup>e</sup>	4 (17%) <sup>e</sup>	4 (6%)


360

361

---


<sup>‡</sup> Values with same superscript are significantly different,  $p < 0.05$

## 362 Legends




363  
 364 Figure 1. Examples of the CT appearance of wooden foreign bodies. A) Transverse, non-contrast CT  
 365 image (level 50HU, width 350HU) showing two relatively dense (110HU) angular wood splinters  
 366 (arrowheads) lodged in the base of the tongue of a dog with chronic signs of swelling (\*), dysphagia, and  
 367 pyrexia. Foci of gas are visible dorsal to the wood fragments. B) Oblique, non-contrast CT image (level -  
 368 500HU, width 2000HU) showing a tree branch (between arrowheads) (-342HU) lodged in the axilla  
 369 following an acute penetrating injury. Multiple small gas foci are present in the adjacent tissues. C)  
 370 Transverse, post-contrast CT image (level 50HU, width 350HU) of the caudal abdomen showing a well-  
 371 defined, linear lucent tract through the left gluteal muscles of a dog that had an acute penetrating  
 372 wound. D) Same image as C displayed using a lung window (level -500HU, width 2000HU). The foreign  
 373 body (-320HU) is now visible as wood with a low density core. A piece of garden cane was removed  
 374 surgically.

375



376  
 377 Figure 2. Examples of CT signs associated with chronic wooden foreign bodies in dogs. A) Sagittal, post-  
 378 contrast CT image (level 50HU, width 350HU) showing an abscess (between arrowheads) surrounding a  
 379 linear wood fragment (150HU) in the cervical region of a dog with focal painful swelling following  
 380 suspected oral stick injury 3 weeks previously. B) Transverse, non-contrast CT image (level 50HU, width  
 381 350HU) showing marked, diffuse fat stranding (\*) affecting the right inguinal region of a dog with acute  
 382 onset of swelling and right pelvic lameness. A narrow linear foreign body (70HU) and associated gas  
 383 tract are visible (arrowheads). A kebab stick was removed surgically. C) Surface-rendered, 3-dimensional  
 384 CT image of the caudal thorax showing a solid, irregular periosteal reaction affecting the right 12<sup>th</sup> rib in  
 385 a dog with a chronic draining sinus on the right thoracic wall. A kebab stick was removed surgically from  
 386 this site. L, left; R, right.

387



388

389 Figure 3. Example of use of multiplanar reconstruction to aid recognition of a large wooden foreign  
 390 body. A) Transverse image of the cranial thorax of a dog with deep wound in the left axilla shows  
 391 extensive sub-cutaneous emphysema, a small volume pneumothorax (black arrowhead), pleural fluid  
 392 (\*), and an oblong heterogeneous structure (between white arrowheads) on the lateral aspect of the left  
 393 lung. The nature of this structure is not evident from this image alone. B) Oblique, approximately  
 394 sagittal, image of the same dog shows clearly an 88mm long stick (between arrowheads) lodged in the  
 395 thoracic wall and lung. Level -500HU, width 2000HU for both images.