

This is the peer-reviewed, manuscript version of the following article:

Clark, E. L., Tomley, F. M. and Blake, D. P. 'Are *Eimeria* Genetically Diverse, and Does It Matter?', *Trends in Parasitology*.

The final version is available online via <http://dx.doi.org/10.1016/j.pt.2016.08.007>.

© 2016. This manuscript version is made available under the CC-BY-NC-ND 4.0 license <http://creativecommons.org/licenses/by-nc-nd/4.0/>.

The full details of the published version of the article are as follows:

TITLE: Are *Eimeria* Genetically Diverse, and Does It Matter?

AUTHORS: Emily L. Clark, Fiona M. Tomley, Damer P. Blake

JOURNAL TITLE: Trends in Parasitology

PUBLISHER: Elsevier (Cell Press)

PUBLICATION DATE: August 2016 (online)

DOI: 10.1016/j.pt.2016.08.007

1 **Are *Eimeria* genetically diverse, and does it matter?**

2

3 **Emily L. Clark [§], Fiona M. Tomley and Damer P. Blake**

4

5 Department of Pathology and Pathogen Biology, Royal Veterinary College, North
6 Mymms, Hertfordshire, UK.

7 [§]Current address: The Roslin Institute, The University of Edinburgh, Midlothian, UK.

8

9 **Correspondence:** dblake@rvc.ac.uk (D.P. Blake)

10 **Keywords:** *Eimeria tenella*, genetic diversity, chickens, vaccines

11

12

13 **Abstract**

14 *Eimeria* pose a risk to all livestock species as a cause of coccidiosis, reducing
15 productivity and compromising animal welfare. Pressure to reduce drug use in the
16 food chain makes development of cost-effective vaccines against *Eimeria* essential.
17 For novel vaccines to be successful, understanding genetic and antigenic diversity in
18 field populations is key. *Eimeria* species that infect chickens are most significant, with
19 *Eimeria tenella* among the best studied and most economically important. Genome-
20 wide single nucleotide polymorphism-based haplotyping has been used to determine
21 population structure, genotype distribution, and potential for cross-fertilization
22 between *E. tenella* strains. Here, we discuss recent developments in our
23 understanding of diversity for *Eimeria* in relation to its specialized lifecycle,
24 distribution across the globe, and the challenges posed to vaccine development.

25

26

27

Genome sequences pave new roads for anticoccidial vaccine development

Eimeria species, protozoan parasites that can cause the damaging intestinal disease coccidiosis, pose a significant risk to global poultry production [1, 2]. Members of the phylum Apicomplexa, the genus *Eimeria* encompasses at least 1 200 species, almost all of which are restricted to a single host [3]. Seven *Eimeria* species are recognized to infect the chicken, causing a considerable disease burden across the globe [2, 4, 5]. Similarly, several species are considered to be highly pathogenic in turkeys (reviewed in [6]). Whilst the pathology associated with each *Eimeria* species infecting chickens has long been understood [7, 8], parasite population structures and the extent of genetic diversity in field populations are only now emerging. Interest in parasite occurrence, diversity and epidemiology is driven by a global need for cheap and effective vaccines as alternatives to anticoccidial drugs. Details of regional variation in *Eimeria* species prevalence, distribution of genetically and antigenically distinct strains, and the frequency at which polymorphic strains cross-fertilize, all provide valuable knowledge that can underpin rational vaccine design and development. In particular the recent availability of genome sequence resources for all seven *Eimeria* species of the chicken [9] provides opportunities to define many of the variables outlined above [10].

Here we review and discuss recent findings relating to the genetic and antigenic diversity of *Eimeria* species which infect chickens in the context of vaccine development and the potential for future successes based on new sequencing technologies and the search for novel vaccine candidate antigens.

Current control strategies for *Eimeria* which infect chickens

More than 60 billion chickens are produced in the world every year, yielding 1.1 trillion eggs and more than 90 million tonnes of meat [11]. The poultry industry in the United States of America (USA) alone is worth in excess of \$38.1 billion, which includes the combined production value of chickens and turkeys [12]. Consequently, effective means of controlling pathogens which infect chickens are essential and of increasing importance as trends for expansion and intensification of global poultry production are maintained (Grace *et al.*, 2012; https://cgspace.cgiar.org/bitstream/handle/10568/21161/ZooMap_July2012_final.pdf).

Control of coccidiosis in poultry relies predominantly on chemoprophylaxis, although resistance to anticoccidial drugs is common in *Eimeria* field populations [13-15]. Prior to the year 2000, anticoccidial drugs were used in ~95% of flocks where anticoccidial control was employed, including ~99% of commercial broiler flocks [16]. More recently, a study from the USA has reported that this percentage has fallen to between 60 and 99%, depending upon the time of year [16]. While anticoccidial drugs remain essential to chicken production and these trends are not yet reflected in much of the world, reductions in drug application throughout the food chain driven by legislative and consumer pressure is encouraging alternatives for coccidiosis control [13, 17]. The use of live oocyst vaccines comprising mixes of species of non-attenuated (formerly wild-type) or attenuated parasites [2] are well established. Oral exposure to controlled numbers of vaccine oocysts is designed to result in low grade coccidial infection, inducing a protective immune response that is boosted by re-infection as the live vaccine re-circulates through the chicken house. However, vaccine production costs and the requirement for multiple parasite lines in each vaccine have been

76 significant barriers to the widespread use of live vaccines in the majority broiler
77 production sector [2]. Nonetheless, non-attenuated vaccines are now included in
78 anticoccidial rotation programs by 35-40% of commercial broiler companies in the
79 USA [16].

80 Recombinant subunit vaccines have been considered as potential alternatives
81 for coccidiosis control for many years, and the concept has returned to the fore in the
82 past decade with the discovery and testing of many partially immunoprotective
83 antigens and expansion of the number of vaccine delivery systems available for use in
84 chickens. Low genetic variability in the target antigen(s) is a key requirement for
85 success precisely because recombinant vaccines rely on the expression of a single, or
86 a small number of antigens [18, 19]. Vaccination using such a small subset of antigens
87 from a complex parasite such as *Eimeria* may provide a significant driving force for
88 immune selection, which could lead to the rapid appearance and dissemination of
89 alleles which confer vaccine-escape (resistance) [10]. The phylum Apicomplexa
90 encompasses a number of parasites important for human and/or animal health
91 including *Plasmodium falciparum* and *Toxoplasma gondii*. The well-characterized
92 population structures and genetic diversity of these parasites have shown that there
93 are numerous barriers to the success of subunit vaccines, but have inspired relevant
94 vaccine development (e.g. [20-22]). In contrast, rather little is known of the genetic
95 diversity and structure of field populations of *Eimeria* parasites, the potential for
96 mixing between genotypes or the selective pressures imposed on loci which encode
97 immunoprotective antigens, highlighting the numerous challenges posed to the
98 development of novel subunit vaccines (reviewed in [11]).

Defining genetic diversity within *Eimeria* species

Eimeria parasites have been recognized for more than a century [7]. Early approaches to understanding parasite diversity focused on parasite (mainly oocyst) morphology, lifecycle (location and timing of development in the gut) and pathogenicity [23]. Differences in the mobility of specific metabolic enzymes during starch gel electrophoresis by isoelectric focusing permitted discrimination between *Eimeria* species and some strains [24], but it was only with the application of techniques that visualize DNA such as pulsed field gel electrophoresis to examine chromosomes, and amplified fragment length polymorphism to examine polymorphisms, that genetic variation began to be explored (reviewed in [25]). Now, advances in molecular biology permit the detailed definition of genetic diversity at specified loci of interest and across whole genomes (Box 1).

Assessing genetic diversity of *Eimeria* using defined locus sequencing

Sequencing short genomic regions, such as internal transcribed spacer (ITS) or mitochondrial cytochrome oxidase subunit 1 (COX1) loci, has been used widely to infer the relatedness of *Eimeria* isolates, particularly those collected from the field. The technique is relatively inexpensive, can be carried out with limited laboratory resources, and is supported by a published sequence archive with ~1 000 and ~100 sequences currently available for ITS and COX1 respectively (GenBank; accessed 7th June, 2016 <http://www.ncbi.nlm.nih.gov/pubmed> using the 'nucleotide' menu).

ITS sequencing has been the molecular technology used most widely for assessing *Eimeria* occurrence in field populations. Initially, studies focused largely on separate countries or continents with examples including Australia, India, Africa and

the USA [5, 26-28]. The most comprehensive survey of *Eimeria* field isolates was published recently in which 512 pooled faecal samples were surveyed from poultry farms situated in 20 countries across five continents [29]. Here, ITS1-5.8S-ITS2 sequence analysis revealed some interesting aspects of population structure. The genetic signatures of *Eimeria acervulina* and *Eimeria mitis* indicated that regular interbreeding occurs between genotypes, while *Eimeria tenella* exhibited a more restricted population structure [10, 29]. The inclusion of sequences derived from laboratory reference strains that are progenitors to many vaccine parasites in the comparison suggested that the samples collected were representative of wild-type field strains, not re-sampling of vaccinal lines [29]. It was suggested that the faster generation time and greater fecundity of *E. acervulina* and *E. mitis* compared to *E. tenella* (~33% shorter prepatent period and 2.5-4 times more oocysts produced per oocyst ingested [30, 31]) could account for the observed differences in population structure. As a consequence, *E. acervulina* and *E. mitis* parasites have greater opportunity for co-infection and hybridization and their genomes may evolve more rapidly.

Analysis of ITS sequence datasets has also led to the discovery of three new *Eimeria* 'operational taxonomic units' (OTUs) [5, 26, 29, 32]. Initially, ITS2 sequencing of isolates from Australia provided the first definition of the three *Eimeria* OTU genotypes termed OTUx, OTUy and OTUz [26], which have been supported by subsequent ITS1 and ITS2 sequencing of isolates covering a greater geographical range [29, 32, 33]. These divergent parasites appear to be restricted at present to southern regions of the world below 30°N latitude [29, 34], although future human and trade movements risk the expansion of their range. The spread of parasites with these novel

genotypes may have significant consequences for vaccine development and application. At present it is unclear whether these variants can evade the immune protection offered by live vaccines, although Morris and colleagues have provided one example of escape from the field [35]. Sequence comparison currently suggests that OTUx is most closely related to *Eimeria maxima*, with *Eimeria brunetti* the closest link to OTUy [32]. Comparison of ITS1-5.8S rDNA-ITS2 sequences has revealed the greatest divergence for OTUz with distinct long and short forms, as described previously for *E. maxima* and *E. mitis* [27, 29].

The development of next generation sequencing technologies has moved analysis of genomic diversity from the single gene to genome wide levels, vastly increasing available genetic and genomic resources for *Eimeria*. For example, fully resolving the phylogenetic relationships between *Eimeria* species which infect chickens and turkeys has proven difficult based on COX1 and 18S rDNA sequences alone [36-38]. The robustness of separation of the seven *Eimeria* species recognized to infect chickens was greatly improved using whole-genome phylogenies [9] and may prove beneficial in future analyses of field isolates. Mitochondrial genome sequencing has also been used effectively to separate *Eimeria* species which infect domestic turkeys [39] (Table 1). The addition of genome sequences resources for cloned OTU x, y and z lines are a high priority and should resolve the cryptic status of these genotypes.

A Dynamic and Adaptable Genome?

Beyond the resolution of phylogenetic debate, whole genome sequencing has revealed interesting aspects of genome structure for *Eimeria* (refer to Table 1 for a

summary of resources). Initial analysis of *E. tenella* chromosome 1, sequenced following purification from pulse field gel electrophoresis-resolved karyotypes, revealed alternating regions of repeat-poor (P) and repeat-rich (R) sequences [40]. More recently, Illumina-based genome sequencing and assembly demonstrated that the P and R structure was not limited to chromosome 1, but was conserved in all chromosomes of *E. tenella*, and across the genomes of *E. acervulina*, *E. brunetti*, *E. maxima*, *E. mitis*, *E. necatrix*, and *E. praecox*, as well as *Eimeria falciformis* (a parasite specific to the mouse) [9, 41]. Interestingly, differences were observed in repeat content between species. *Eimeria tenella*, for example, has fewer R regions than the other six species which infect chickens, while *E. necatrix* was more repeat rich across its genome, most notably in regions syntenic with *E. tenella* [9]. This P/R structure does not appear in the genomes of other coccidia such as *Neospora caninum* or *T. gondii* [9, 42], although it has been detected in the more closely related *Cyclospora cayetanensis* genome [43]. Ling and colleagues have suggested that the unusual genome organization might pose an evolutionary advantage to the parasite by facilitating rapid evolution and diversification. Variation in restriction fragment length polymorphism (RFLP) fragment size between different strains of *E. tenella* associated with R-, but not P-regions, lending some support for genome plasticity [9, 40, 44]. A disproportionately high repeat content in protein coding sequences could confer some evolutionary advantage, although their effects on protein structure appear to be neutral and genes known to be integral to host–parasite interaction were relatively free of repeats [9]. The seven *Eimeria* species which infect poultry do not appear to possess the sub-telomeric regions which in *P. falciparum* contain a set of plastic genes involved in host immune system evasion [9, 40, 45]. Telomere-like repeats are,

however, dispersed throughout the R-segments, suggesting there are complexities in the structure of the genome that we do not fully yet understand. Telomere-like repeats have previously been described in the *Plasmodium knowlesi* genome where they associate with variant antigen families, although a similar linkage has not been described for *Eimeria* [46]. The impact of the segmented *Eimeria* genome structure on the appearance and extent of genetic diversity is yet to be determined, although it may well associate with hotspots of genetic recombination. The implications of such hotspots on vaccine development are similarly unclear.

The importance of population structure

Genetic mapping has been useful in establishing the population structure of some apicomplexans (reviewed in [47]), as have other molecular tools (reviewed in [25]). Population structure varies across the Apicomplexa. *Plasmodium falciparum* has been shown to exhibit signatures of panmictic or clonal population structures, influenced in part by regional transmission rates [48, 49]. *Toxoplasma gondii* is commonly clonal in much of the world, with a small number of dominant genotypes described, although a higher level of genetic diversity has been detected in regions such as South America where population mixing appears to occur at a greater frequency [21, 50]. For *E. tenella*, comparison of haplotype occurrence and diversity defined following multiplex single nucleotide polymorphism (SNP) genotyping revealed that north Indian and north African field populations were characterized by a limited number of distinct haplotypes and significant linkage disequilibrium (Figure 1), resembling the region specific population structure of *T. gondii* [10]. This population structure suggests that limited opportunities exist for cross-fertilization

and genetic recombination, and that the expansion of a small number of haplotypes might be common although not necessarily clonal. In contrast, a greater haplotype diversity was reported in southern India and Nigeria with multiple haplotypes appearing, all at a very low frequency, indicating that co-infection with heterologous isolates and cross-fertilization is common during sexual reproduction, and that genetic diversity is likely to be greater than estimated by current sampling [10]. These findings suggest there are numerous opportunities for recombination in the field in these regions.

The regional differences in population structure observed for *E. tenella* may have several underlying causes. In southern India there is a greater poultry density than found in the north (Grace *et al.*, 2012; https://cgspace.cgiar.org/bitstream/handle/10568/21161/ZooMap_July2012_final.pdf), and therefore more opportunity for parasite co-infection and cross-fertilization. Further, the climate in south India is commonly more humid than in north India [51], likely favoring higher levels of oocyst sporulation and increased parasite survival in the poultry house environment as reported in comparisons of rainy versus dry seasons [52]. Higher rates of transmission commonly associate with elevated levels of outcrossing and increased genotype abundance for other apicomplexans such as *P. falciparum* [49]. Importantly, co-infection of a single host with two or more genetically distinct *Eimeria* isolates does not guarantee cross-fertilization. The *Eimeria* life cycle includes a single, transiently diploid phase during sexual reproduction and oocyst maturation [53], so timing of the co-infection has to be essentially simultaneous for genetic recombination to occur. Additionally, the *in vivo* phase of the *Eimeria* life cycle is predominantly self-limiting, with features such as prepatent period and the number

of rounds of schizogony stable, unless subjected to deliberate selection for developmental rate [10, 54, 55]. Studies using major histocompatibility complex (MHC) class I or II knockout mice suggest little or no role for the host immune response in the conclusion of parasite replication [56]. Thus, gametes of each genotype must mature in parallel for cross-fertilization to take place. *In vivo* experiments using laboratory strains of *E. tenella* have shown that, given the opportunity, cross-fertilization is common, highlighting the potential that *E. tenella* has to hybridize in field populations and indicating the ease with which vaccine or drug-resistant alleles could propagate in field parasite populations [10]. Combined, these factors emphasize the importance of considering region specific environmental and social variables in implementation of novel control strategies for *Eimeria* species. Fornace and colleagues demonstrated that the diversity of species present in small-scale production systems in Africa was directly linked to profitability [5]. However, there have been few similar studies and the potential is there to link population structure and the burden of coccidiosis to profitability in particular regions of the globe.

The relevance of antigenic diversity

Selection of candidate antigens for vaccine development has proved to be a significant barrier to progress in other Apicomplexa such as *T. gondii* and the *Plasmodium* species [57-59]. Differentiating immunogenicity from 'true' immune protection can be difficult, making selection of protective antigens problematic [19]. In one example, homologs of apical membrane antigen 1 (AMA-1) have been shown to be protective in a range of apicomplexan parasites including *E. maxima* [19], *E. tenella* [60] and *P. falciparum* [61-63], and it has been widely proposed as a candidate

for subunit vaccine development. However, extensive allelic diversity has limited development of *P. falciparum* AMA-1, with more than 60 polymorphic amino acid residues detected and more than 200 haplotypes within even a single population [61, 62, 64, 65]. Despite such discouraging reports from *P. falciparum*, AMA-1 has shown promise as a vaccine candidate for *E. tenella*, with a potent inhibitory effect on parasite invasion [60]. More recently, genotyping *E. tenella* field isolates collected from Africa and India suggested that polymorphisms in the EtAMA-1 locus are lower than expected in field populations with largely neutral signatures of selection. The functionality of AMA-1 may outweigh the potential benefit to the parasite of immune evasion, which may be of limited value in the self-limiting eimerian life cycle [10]. Similarly, just four nucleotide polymorphisms exist between EmAMA-1 coding sequences from the *E. maxima* Houghton and Weybridge laboratory strains, two causing non-synonymous changes, one situated in the putative pro-domain and one located in domain 1 [9, 66]. Nonetheless, despite such limited diversity within the coding region of at least one vaccine candidate, strain specific immune escape has been reported *in vivo* for *E. acervulina* [67, 68], *E. mitis* [69], *E. maxima* [70] and *E. tenella* [71-73]. Comparison of *E. tenella* isolates collected from chickens reared in British and Indian poultry houses revealed incomplete immune protection between isolates, most notably following low-level primary exposure [10]. Despite these reports, there is no evidence that vaccine resistance has evolved in response to whole live parasite vaccination [2, 11]. One possible explanation for this is that throughout its lifecycle each *Eimeria* species expresses between 6 000 and 9 000 proteins [9], exposing the host to a complex portfolio of antigens. Selection targeting multiple immunoprotective antigens in parallel during replication in the chicken is likely to limit

the capacity for any individual parasite to evade the host immune response as a consequence of diversifying selection. Thus, the complexity of the antigenic repertoire might explain why resistance to live parasite vaccination has not yet developed [10] [11]. Incorporating multiple antigens, in addition to AMA-1, in novel subunit vaccines would therefore be likely to extend their potential for long term success by buffering the effects of diversifying selection on a single target antigen.

Life cycle stage-specific antigen expression and immune selection

Each *Eimeria* life cycle features a series of extra- and intracellular stages within the definitive host as the parasite undergoes successive rounds of asexual, and then sexual replication [9, 53]. Throughout this process *Eimeria* expresses many of its genes in a stage-specific manner which can impact on the development of novel vaccines. In *T. gondii*, for example, vaccination with life cycle stage-specific antigens leads to stage-limited protection [57, 58]. In *Eimeria*, the early life cycle stages are important to the induction of protective immunity during natural infection [9, 11, 60]. Importantly, vaccine candidates such as AMA-1 are primarily expressed by a single life cycle stage and are unlikely to be subjected to a protein-specific adaptive immune response during primary infection given the absence of protracted colonization [10, 54, 60]. Thus, the large oocyst output resulting from even low dose primary infections results in considerable environmental contamination with parasites which have never been exposed to immune selection.

Future directions

315 A clear direction for future work is to expand our understanding of population
316 structure to other *Eimeria* species in the field as has been reported recently for *E.*
317 *tenella* [10]. Elucidating the population structure and potential for mixing is key in the
318 development of novel control strategies for *Eimeria*. Understanding the possible
319 biological, environmental, industrial and social drivers which underpin the observed
320 diversity may be even more important, demanding detailed epidemiological
321 interrogation. Opportunities to develop medium/high throughput tools such as
322 Sequenom-based genotyping, and new high-throughput sequencing technologies
323 such as restriction site associated DNA (RAD) sequencing, will facilitate the move away
324 from ITS sequencing to genome wide analysis of genetic diversity with particular
325 relevance to field samples. *Eimeria* genomic resources have increased greatly in recent
326 years (reviewed in [74]). Additionally, since the cost of sequencing a genome the size
327 of *E. tenella* is now relatively modest (51.8 Mb DNA in the current genome assembly
328 [9]), the opportunity exists to build on the available genomic resources with whole
329 genome sequencing of other *Eimeria* strains and species. Parasites of the three OTU
330 genotypes are obvious candidates, with species which infect other livestock species
331 further priorities. The genomes of non-target species can yield clues as to the
332 structure and function of other closely related species. Comparative analysis of the *E.*
333 *falciformis* genome with *T. gondii* revealed a shared emergence and diversification
334 across the Coccidia of gene families associated with motility and invasion [41]. Building
335 on information from whole genome sequencing, another relatively new technology,
336 RNA sequencing (RNASeq) can be used for transcriptomic profiling of other key
337 antigens of interest and is likely to offer clues as to their function and suitability as
338 vaccine targets. RNASeq has already been used successfully to define transcriptomes

from several *Eimeria* life cycle stages [9, 53]. Indeed in the near future Isoform sequencing (IsoSeq), which at present generates transcripts >3Kb [75], could be utilized to sequence the entire transcriptome of a single parasite in full length fragments. Genome editing techniques such as the CRISPR/Cas system have huge potential and could be used, for example, to switch allelic type for a small number of target antigen coding genes. The CRISPR/Cas system has been used successfully in *P. falciparum* [76, 77] and *T. gondii* [78, 79], but is not yet available for *Eimeria*. These tools should improve the molecular definition of diversity, expand our understanding of parasite evolution and host evasion, and highlight regions of the genome that show promise in the development of novel sub-unit vaccines.

Concluding Remarks

There are several key challenges posed by population, genetic and antigenic diversity of *Eimeria* parasites to the development of novel vaccines (see Outstanding Questions). How genetic, particularly antigenic, diversity influences pathogenicity, vaccine specificity and epidemiology, and the implications of this for effective intervention and control, are important questions that need to be answered for all apicomplexan parasites. Recent studies have revealed a polarized global occurrence for genetically divergent *Eimeria* strains, and possibly even new species, that may be capable of replicating within chickens vaccinated using current generation vaccines. These parasites pose a significant risk to vaccine efficacy, and thus food security and animal welfare, in production systems which rely on anticoccidial vaccination. Considering social and environmental variables in novel control strategies is of great importance, with factors including choice of production system, geographic

separation of farms and climatic conditions likely to influence parasite population dynamics. The recent expansion in genetic and genomic resources available for *Eimeria* has dramatically improved our ability to genotype parasites recovered from field populations and begin to assess how many of these variables will affect genetic diversity, and whether that diversity will impact on vaccine efficacy and longevity.

Acknowledgements

Much of the work reviewed here was funded by DFID and the BBSRC through the Combating Infectious Diseases of Livestock for International Development (CIDLID) initiative under project reference BB/H009337 and BB/L00478X. This manuscript has been assigned the reference PPB_01354 by the RVC.

References

- 1 Dalloul, R.A. and Lillehoj, H.S. (2006) Poultry coccidiosis: recent advancements in control measures and vaccine development. *Expert Rev Vaccines* 5, 143-163
- 2 Shirley, M.W., *et al.* (2005) The biology of avian *Eimeria* with an emphasis on their control by vaccination. *Adv Parasitol* 60, 285-330
- 3 Chapman, H.D., *et al.* (2013) A selective review of advances in coccidiosis research. *Advances In Parasitology* 83, 93-171
- 4 Williams, R.B. (1998) Epidemiological aspects of the use of live anticoccidial vaccines for chickens. *International Journal of Parasitology* 28, 1089-1098
- 5 Fornace, K.M., *et al.* (2013) Occurrence of *Eimeria* species parasites on small-scale commercial chicken farms in Africa and indication of economic profitability. *PLoS ONE* 8, e84254
- 6 Chapman, H. (2008) Coccidiosis in the turkey. *Avian Pathology* 37, 205-223
- 7 Chapman, H.D. (2014) Milestones in avian coccidiosis research: a review. *Poult Sci* 93, 501-511
- 8 Long, P., *et al.* (1976) A guide to laboratory techniques used in the study and diagnosis of avian coccidiosis. *Folia Veterinaria Latina* 6, 201-217
- 9 Reid, A., *et al.* (2014) Genomic analysis of the causative agents of coccidiosis in domestic chickens. *Genome Research* 24, 1676-1685
- 10 Blake, D.P., *et al.* (2015) Population, genetic, and antigenic diversity of the apicomplexan *Eimeria tenella* and their relevance to vaccine development. *Proc Natl Acad Sci U S A* 112, E5343-5350
- 11 Blake, D.P. and Tomley, F.M. (2014) Securing poultry production from the ever-present *Eimeria* challenge. *Trends Parasitol* 30, 12-19

399 12 NASS (2012) Poultry - Production and Value. (Agriculture, U.S.D.o., ed), USDA
 400 13 Shirley, M.W., *et al.* (2007) Challenges in the successful control of the avian
 401 coccidia. *Vaccine* 25, 5540-5547
 402 14 Joyner, L. and Norton, C. (1975) Transferred drug-resistance in *Eimeria*
 403 *maxima*. *Parasitology* 71, 385-392
 404 15 Chapman, H.D. (1976) *Eimeria tenella* in chickens: studies on resistance to the
 405 anticoccidial drugs monensin and lasalocid. *Veterinary Parasitology* 2, 187-196
 406 16 Chapman, H.D. and Jeffers, T.K. (2014) Vaccination of chickens against
 407 coccidiosis ameliorates drug resistance in commercial poultry production.
 408 *International journal for parasitology. Drugs and drug resistance* 4, 214-217
 409 17 Godfray, H.C., *et al.* (2010) Food security: the challenge of feeding 9 billion
 410 people. *Science* 327, 812-818
 411 18 Blake, D.P., *et al.* (2004) Parasite genetics and the immune host:
 412 recombination between antigenic types of *Eimeria maxima* as an entree to the
 413 identification of protective antigens. *Mol Biochem Parasitol* 138, 143-152
 414 19 Blake, D.P., *et al.* (2011) Genetic mapping identifies novel highly protective
 415 antigens for an apicomplexan parasite. *PLoS Pathog* 7, e1001279
 416 20 Amambua-Ngwa, A., *et al.* (2012) Population genomic scan for candidate
 417 signatures of balancing selection to guide antigen characterization in malaria
 418 parasites. *PLoS Genet* 8, e1002992
 419 21 Minot, S., *et al.* (2012) Admixture and recombination among *Toxoplasma*
 420 *gondii* lineages explain global genome diversity. *Proc Natl Acad Sci U S A* 109,
 421 13458-13463
 422 22 Manske, M., *et al.* (2012) Analysis of *Plasmodium falciparum* diversity in
 423 natural infections by deep sequencing. *Nature* 487, 375-379
 424 23 Tyzzer, E. (1929) Coccidiosis in gallinaceous birds. *American Journal of*
 425 *Hygiene* 10, 269-383
 426 24 Shirley, M.W., *et al.* (1989) Enzyme variation and pathogenicity of recent field
 427 isolates of *Eimeria tenella*. *Res Vet Sci* 46, 79-83
 428 25 Beck, H.P., *et al.* (2009) Molecular approaches to diversity of populations of
 429 apicomplexan parasites. *Int J Parasitol* 39, 175-189
 430 26 Cantacessi, C., *et al.* (2008) Genetic characterization of three unique
 431 operational taxonomic units of *Eimeria* from chickens in Australia based on
 432 nuclear spacer ribosomal DNA. *Vet Parasitol* 152, 226-234
 433 27 Schwarz, R.S., *et al.* (2009) Genomic analysis of *Eimeria* spp. populations in
 434 relation to performance levels of broiler chicken farms in Arkansas and North
 435 Carolina. *J Parasitol* 95, 871-880
 436 28 Kundu, K., *et al.* (2015) Cloning and sequencing of beta-tubulin and internal
 437 transcribed spacer-2 (ITS-2) of *Eimeria tenella* isolate from India. *J Parasit Dis* 39,
 438 539-544
 439 29 Clark, E.L., *et al.* (2016) Cryptic *Eimeria* genotypes are common across the
 440 southern but not northern hemisphere. *Int J Parasitol* 46, 537-544
 441 30 Bumstead, N. and Millard, B. (1992) Variation in susceptibility of inbred lines
 442 of chickens to seven species of *Eimeria*. *Parasitology* 104, 407-413
 443 31 Eckert, J., *et al.* (1995) Guidelines on Techniques in Coccidiosis Research.
 444 (Comission, E., ed)
 445 32 Godwin, R.M. and Morgan, J.A. (2015) A molecular survey of *Eimeria* in
 446 chickens across Australia. *Vet Parasitol* 214, 16-21

447 33 Jatau, I.D., *et al.* (2016) Three operational taxonomic units of *Eimeria* are
 448 common in Nigerian chickens and may undermine effective molecular diagnosis
 449 of coccidiosis. *BMC Vet Res* 12, 86
 450 34 Clark, E.L. and Blake, D.P. (2012) Genetic mapping and coccidial parasites:
 451 past achievements and future prospects. *J Biosci* 37, 879-886
 452 35 Morris, G.M., *et al.* (2007) Investigating a persistent coccidiosis problem on a
 453 commercial broiler-breeder farm utilising PCR-coupled capillary electrophoresis.
 454 *Parasitol Res* 101, 583-589
 455 36 Ogedengbea, J.D., *et al.* (2011) DNA barcoding identifies *Eimeria* species and
 456 contributes to the phylogenetics of coccidian parasites (Eimeriorina,
 457 Apicomplexa, Alveolata). *International Journal of Parasitology* 41, 843-850
 458 37 El-Sherry, S., *et al.* (2013) Divergent nuclear 18S rDNA paralogs in a turkey
 459 coccidium, *Eimeria meleagridis*, complicate molecular systematics and
 460 identification. *International Journal of Parasitology* 43, 679-685
 461 38 Miska, K.B., *et al.* (2010) Molecular characterization and phylogenetic analysis
 462 of *Eimeria* from turkeys and gamebirds: implications for evolutionary
 463 relationships in Galliform birds. *J Parasitol* 96, 982-986
 464 39 Ogedengbe, M., *et al.* (2014) Complete mitochondrial genome sequences from
 465 five *Eimeria* species (Apicomplexa; Coccidia; Eimeriidae) infecting domestic
 466 turkeys. *Parasite Vectors* 17, 335
 467 40 Ling, K., *et al.* (2007) Sequencing and analysis of chromosome 1 of *Eimeria*
 468 *tenella* reveals a unique segmental organization. *Genome Research* 17, 311-319
 469 41 Heitlinger, E., *et al.* (2014) The genome of *Eimeria falciformis*--reduction and
 470 specialization in a single host apicomplexan parasite. *BMC Genomics* 15, 696
 471 42 Reid, A.J., *et al.* (2012) Comparative genomics of the apicomplexan parasites
 472 *Toxoplasma gondii* and *Neospora caninum*: Coccidia differing in host range and
 473 transmission strategy. *PLoS Pathog* 8, e1002567
 474 43 Liu, S., *et al.* (2016) Comparative genomics reveals *Cyclospora cayetanensis*
 475 possesses coccidia-like metabolism and invasion components but unique surface
 476 antigens. *BMC Genomics* 17, 316
 477 44 Shirley, M.W. (1994) Coccidial parasites from the chicken: discrimination of
 478 different populations of *Eimeria tenella* by DNA hybridisation. *Research In*
 479 *Veterinary Science* 57, 10-14
 480 45 Gardner, M., *et al.* (2002) Genome sequence of the human malaria parasite
 481 *Plasmodium falciparum*. *Nature* 419, 498-511
 482 46 Pain, A., *et al.* (2008) The genome of the simian and human malaria parasite
 483 *Plasmodium knowlesi*. *Nature* 455, 799-803
 484 47 Clark, E.L. and Blake, D.P. (2012) Genetic mapping and coccidial parasites:
 485 past achievements and future prospects. *Journal of Biosciences* 37, 879-886
 486 48 Annan, Z., *et al.* (2007) Population genetic structure of *Plasmodium falciparum*
 487 in the two main African vectors, *Anopheles gambiae* and *Anopheles funestus*. *Proc*
 488 *Natl Acad Sci U S A* 104, 7987-7992
 489 49 Larranaga, N., *et al.* (2013) Genetic structure of *Plasmodium falciparum*
 490 populations across the Honduras-Nicaragua border. *Malar J* 12, 354
 491 50 Su, C., *et al.* (2012) Globally diverse *Toxoplasma gondii* isolates comprise six
 492 major clades originating from a small number of distinct ancestral lineages. *Proc*
 493 *Natl Acad Sci U S A* 109, 5844-5849
 494 51 Deichmann, U. and Eklundh, L. (1991) *Global digital data sets for land*
 495 *degradation studies: a GIS approach*. UNEP/GEMS and GRID; Nairobi, Kenya

496 52 Awais, M.M., *et al.* (2012) Seasonal prevalence of coccidiosis in industrial
 497 broiler chickens in Faisalabad, Punjab, Pakistan. *Trop Anim Health Prod* 44, 323-
 498 328
 499 53 Walker, R., *et al.* (2015) RNA Seq analysis of the *Eimeria tenella* gametocyte
 500 transcriptome reveals clues about the molecular basis for sexual reproduction
 501 and oocyst biogenesis. *BMC Genomics* 16, 94
 502 54 Lal, K., *et al.* (2009) Proteomic comparison of four *Eimeria tenella* life-cycle
 503 stages: unsporulated oocyst, sporulated oocyst, sporozoite and second-
 504 generation merozoite. *Proteomics* 9, 4566-4576
 505 55 Shirley, M.W. and Harvey, D.A. (2000) A genetic linkage map of the
 506 apicomplexan protozoan parasite *Eimeria tenella*. *Genome Res* 10, 1587-1593
 507 56 Smith, A.L. and Hayday, A.C. (2000) Genetic dissection of primary and
 508 secondary responses to a widespread natural pathogen of the gut, *Eimeria*
 509 *vermiformis*. *Infect Immun* 68, 6273-6280
 510 57 Alexander, J., *et al.* (1996) Immunological control of *Toxoplasma gondii* and
 511 appropriate vaccine design. *Current Topics in Microbiology and Immunology* 219,
 512 183-195
 513 58 Liu, Q., *et al.* (2012) Vaccines against *Toxoplasma gondii*: status, challenges
 514 and future directions. *Human Vaccines and Immunotherapeutics* 8, 1305-1308
 515 59 Stanisic, D.I., *et al.* (2013) Escaping the immune system: How the malaria
 516 parasite makes vaccine development a challenge. *Trends Parasitol* 29, 612-622
 517 60 Jiang, L., *et al.* (2012) Identification and characterization of *Eimeria tenella*
 518 apical membrane antigen-1 (AMA1). *PLoS ONE* 7, e41115
 519 61 Healer, J., *et al.* (2004) Allelic polymorphisms in apical membrane antigen-1
 520 are responsible for evasion of antibody-mediated inhibition in *Plasmodium*
 521 *falciparum*. *Molecular Microbiology* 52, 159-168
 522 62 Drew, D., *et al.* (2012) Defining the antigenic diversity of *Plasmodium*
 523 *falciparum* apical membrane antigen 1 and the requirements for a multi-allele
 524 vaccine against malaria. *PloS One* 7, e51023
 525 63 Eisen, D., *et al.* (2002) Alterations in *Plasmodium falciparum* genotypes during
 526 sequential infections suggest the presence of strain specific immunity. *American*
 527 *Journal of Tropical Medical Hygiene* 67, 8-16
 528 64 Hodder, A.N., *et al.* (2001) Specificity of the protective antibody response to
 529 apical membrane antigen 1. *Infect Immun* 69, 3286-3294
 530 65 Terheggen, U., *et al.* (2014) Limited antigenic diversity of *Plasmodium*
 531 *falciparum* apical membrane antigen 1 supports the development of effective
 532 multi-allele vaccines. *BMC Medicine* 12, 183
 533 66 Blake, D.P., *et al.* (2012) EmaxDB: Availability of a first draft genome sequence
 534 for the apicomplexan *Eimeria maxima*. *Mol Biochem Parasitol* 184, 48-51
 535 67 Joyner, L.P. (1969) Immunological variation between two strains of *Eimeria*
 536 *acervulina*. *Parasitology* 59, 725-732
 537 68 Wu, L., *et al.* (2014) Biological characteristics of Chinese precocious strain of
 538 *Eimeria acervulina* and its immune efficacy against different field strains. *Avian*
 539 *Diseases* 58, 367-372
 540 69 McDonald, V., *et al.* (1985) Attenuation of *Eimeria* species: further
 541 characterisation of two lines of *Eimeria mitis*. *Research in Veterinary Science* 39,
 542 328-332

543 70 Smith, A., *et al.* (2002) Antigenic diversity in *Eimeria maxima* and the
 544 influence of host genetics and immunization schedule on cross-protective
 545 immunity. *Infection and Immunity* 70, 2472-2479
 546 71 Awad, A.M., *et al.* (2013) Evaluation of the protective efficacy of the
 547 anticoccidial vaccine Coccivac-B in broilers, when challenged with Egyptian field
 548 isolates of *E. tenella*. *Parasitol Res* 112, 113-121
 549 72 Abu-Akkada, S. and Awad, A. (2012) Isolation, propagation, identification and
 550 comparative pathogenicity of five Egyptian field strains of *Eimeria tenella* from
 551 broiler chickens in five different provinces in Egypt. *Research In Veterinary*
 552 *Science* 92, 92-95
 553 73 Fitz-Coy, S. (1992) Antigenic variation among strains of *Eimeria maxima* and
 554 *E. tenella* of the chicken. *Avian Diseases* 36, 40-43
 555 74 Blake, D. (2015) *Eimeria* genomics: Where are we now and where are we
 556 going? *Veterinary Parasitology* 212, 68-74
 557 75 Tilgner, H., *et al.* (2014) Defining a personal, allele-specific, and single-
 558 molecule long-read transcriptome. *Proc Natl Acad Sci U S A* 111, 9869-9874
 559 76 Wagner, J.C., *et al.* (2014) Efficient CRISPR-Cas9-mediated genome editing in
 560 *Plasmodium falciparum*. *Nat Methods* 11, 915-918
 561 77 Ghorbal, M., *et al.* (2014) Genome editing in the human malaria parasite
 562 *Plasmodium falciparum* using the CRISPR-Cas9 system. *Nat Biotechnol* 32, 819-
 563 821
 564 78 Sidik, S.M., *et al.* (2014) Efficient genome engineering of *Toxoplasma gondii*
 565 using CRISPR/Cas9. *PLoS One* 9, e100450
 566 79 Shen, B., *et al.* (2014) Efficient gene disruption in diverse strains of
 567 *Toxoplasma gondii* using CRISPR/CAS9. *MBio* 5, e01114-01114
 568 80 Hikosaka, K., *et al.* (2011) Concatenated mitochondrial DNA of the coccidian
 569 parasite *Eimeria tenella*. *Mitochondrion* 11, 273-278
 570 81 Lin, R.Q., *et al.* (2011) Characterization of the complete mitochondrial
 571 genomes of five *Eimeria* species from domestic chickens. *Gene* 480, 28-33
 572 82 Ogedengbe, M.E., *et al.* (2013) Sequencing the complete mitochondrial
 573 genome of *Eimeria mitis* strain USDA 50 (Apicomplexa: Eimeriidae) suggests
 574 conserved start positions for mtCOI- and mtCOIII-coding regions. *Parasitol Res*
 575 112, 4129-4136
 576 83 Ogedengbe, M.E., *et al.* (2014) Complete mitochondrial genome sequences
 577 from five *Eimeria* species (Apicomplexa; Coccidia; Eimeriidae) infecting domestic
 578 turkeys. *Parasit Vectors* 7, 335
 579 84 Liu, G.H., *et al.* (2012) The complete mitochondrial genome sequence of
 580 *Eimeria mitis* (Apicomplexa: Coccidia). *Mitochondrial DNA* 23, 341-343
 581 85 Hafeez, M.A., *et al.* (2016) The complete mitochondrial genome sequence of
 582 *Eimeria innocua* (Eimeriidae, Coccidia, Apicomplexa). *Mitochondrial DNA A DNA*
 583 *MappSeq Anal* 27, 2805-2806
 584 86 Vrba, V. and Pakandl, M. (2014) Coccidia of turkey: from isolation,
 585 characterisation and comparison to molecular phylogeny and molecular
 586 diagnostics. *Int J Parasitol* 44, 985-1000
 587 87 Hnida, J.A. and Duszynski, D.W. (1999) Taxonomy and phylogeny of some
 588 *Eimeria* (Apicomplexa:Eimeriidae) species of rodents as determined by
 589 polymerase chain reaction/restriction-fragment-length polymorphism analysis
 590 of 18S rDNA. *Parasitol Res* 85, 887-894
 591

592

593

Figure legends




Figure 1. Median-joining phylogenetic NETWORKs illustrating genome-wide and antigen specific diversity reported for *Eimeria tenella*. A. The influence of geographic origin on *E. tenella* single nucleotide polymorphism (SNP) haplotype occurrence and complexity. Parasite populations from Nigeria and south India presented high haplotype diversity and apparent panmixia, compared to more restricted variation in north African and north Indian populations. Node size indicates the frequency of haplotype occurrence. Figure reproduced from [10]. B. Coding sequence polymorphism within the apical membrane antigen 1 (AMA-1) locus. Eight allelic types were detected with less geographic specificity than described for genome haplotypes. Figure derived from data presented in [10].

606 **Table 1. Genome size and available genetic resources for *Eimeria* species in comparison with the apicomplexan species**
607 ***Toxoplasma gondii* and *Plasmodium falciparum*^a.**
608

| Species | Host | Genome Size (Mb) | Reference Genome ² | Mitochondrial Genome | RNASeq | Defined Locus Sequencing | SNP Arrays | Proteomics |
|------------------------------|-----------------|------------------|-------------------------------|----------------------|--------|--------------------------|------------|------------|
| <i>Eimeria falciformis</i> | Mouse | 43.67 | • | | | • | | |
| <i>Eimeria acervulina</i> | Chicken | 45.83 | • | • | | • | | |
| <i>Eimeria brunetti</i> | Chicken | 66.89 | • | • | | • | | |
| <i>Eimeria maxima</i> | Chicken | 45.98 | • | • | | • | | |
| <i>Eimeria mitis</i> | Chicken | 72.24 | • | • | | • | | |
| <i>Eimeria necatrix</i> | Chicken | 55.01 | • | • | | • | | |
| <i>Eimeria praecox</i> | Chicken | 60.08 | • | • | | • | | |
| <i>Eimeria tenella</i> | Chicken | 51.86 | • | • | • | • | • | • |
| <i>Eimeria adenoeides</i> | Turkey | - | | • | | • | | |
| <i>Eimeria dispersa</i> | Turkey | - | | • | | • | | |
| <i>Eimeria gallopavonis</i> | Turkey | - | | • | | • | | |
| <i>Eimeria innocua</i> | Turkey | - | | • | | • | | |
| <i>Eimeria meleagridis</i> | Turkey | - | | • | | • | | |
| <i>Eimeria meleagritidis</i> | Turkey | - | | • | | • | | |
| <i>Toxoplasma gondii</i> GT1 | Cat, others | 63.95 | • | • | • | • | • | • |
| <i>Plasmodium falciparum</i> | Mosquito, human | 23.3 | • | • | • | • | • | • |

609 ^aTable adapted from the *Toxoplasma* Genetics Resource *ToxoDB* (<http://www.toxodb.org/toxo/showApplication.do>, accessed 15th June,
610 2016), supplemented by [80-87].

611 ^b• = sequence resource available.

612

613

Box 1. The utility of SNP genotyping assays in determining parasite population dynamics in the field.

Defined locus sequencing has been widely used to genotype *Eimeria* field isolates since it is cost effective and relatively quick to accomplish. However, with the advent of new high throughput sequencing technologies analysis of genetic diversity across whole genomes is now possible. When whole genome sequencing first became available the associated costs were prohibitive, but this is changing rapidly as the technology becomes cheaper. Reference genome sequence assemblies are now available for the seven *Eimeria* species that infect chickens [9]. Large-scale genome re-sequencing of field isolates will soon be possible but is not yet affordable. In the interim period sequencing a small number of additional strains for comparison with the relevant reference genome provides a resource to design genotyping tools based on specific single nucleotide polymorphisms (SNPs). Custom SNP-based assays are a cost effective method of genotyping parasites which can be applied effectively to large-scale collections of field isolates [10]. SNP genotyping technologies provide useful tools to assess the level of cross-fertilization and genetic recombination in field populations. Such knowledge can be employed to improve the prospects of future subunit vaccines being effective in the field.