

This is the peer-reviewed, manuscript version of the following article:

C.P. Rubio, S. Martínez-Subiela, J. Hernández-Ruiz, A. Tvarijonaviciute, J.J. Cerón, K. Allenspach, Serum biomarkers of oxidative stress in dogs with idiopathic inflammatory bowel disease, *The Veterinary Journal* (2017), <http://dx.doi.org/10.1016/j.tvjl.2017.02.003>.

© 2016. This manuscript version is made available under the CC-BY-NC-ND 4.0 license <http://creativecommons.org/licenses/by-nc-nd/4.0/>.

The full details of the published version of the article are as follows:

TITLE: Serum biomarkers of oxidative stress in dogs with idiopathic inflammatory bowel disease

AUTHORS: C.P. Rubio, S. Martínez-Subiela, J. Hernández-Ruiz, A. Tvarijonaviciute, J.J. Cerón, K. Allenspach

JOURNAL: The Veterinary Journal

PUBLISHER: Elsevier

PUBLICATION DATE: 9 February 2017 (online)

DOI: 10.1016/j.tvjl.2017.02.003

1 **Original Article**

2
3
4 **Serum biomarkers of oxidative stress in dogs with idiopathic inflammatory bowel**
5 **disease**

6
7
8 C.P. Rubio ^a, S. Martínez-Subiela ^a, J. Hernández-Ruiz ^b, A. Tvarijonaviciute ^{a,c}, J.J.
9 Cerón ^{a,*}, K. Allenspach ^{d,1}

10
11 ^a *Department of Animal Medicine and Surgery, Faculty of Veterinary Medicine,*
12 *Regional Campus of International Excellence 'Campus Mare Nostrum', University of*
13 *Murcia, Espinardo, Murcia, 30100, Spain*

14 ^b *Department of Plant Biology (Plant Physiology), Faculty of Biology, University of*
15 *Murcia, Espinardo, Murcia, 30100, Spain*

16 ^c *Department of Animal Medicine and Surgery, Veterinary School, University Autònoma*
17 *of Barcelona, Barcelona, 08193, Spain*

18 ^d *Department of Veterinary Clinical Sciences and Services, Royal Veterinary College,*
19 *University of London, North Mymms, AL97PT, UK*

20
21
22 * Corresponding author. Tel.: +34 86884722.
23 *E-mail address: jjceron@um.es (J.J. Cerón)*

24
25
26 ¹ *Dr Allenspach's current address is Veterinary Clinical Sciences, College of Veterinary*
27 *Medicine, Iowa State University, Ames, IA 50011, USA*

29 **Highlights**

- 30 • Decreased total antioxidant capacity, measured by CUPRAC and TEAC assays,
31 was demonstrated in sera from dogs with IBD.
- 32 • Serum thiol and PON1 activity was significantly decreased in sera from dogs
33 with IBD compared with healthy dogs.
- 34 • Oxidative damage in dogs with IBD was demonstrated by increased serum FOX,
35 ROS and TBARS.

36

37 **Abstract**

38 The objective of this study was to evaluate and compare a panel of various
39 serum biomarkers evaluating both the antioxidant response and oxidative damage in
40 dogs with idiopathic inflammatory bowel disease (IBD). Eighteen dogs with IBD and
41 20 healthy dogs were enrolled in the study. Trolox equivalent antioxidant capacity
42 (TEAC), cupric reducing antioxidant capacity (CUPRAC), ferric reducing ability of the
43 plasma (FRAP), total thiol concentrations, and paraoxonase 1 (PON1) activity were
44 evaluated in serum to determine antioxidant response. To evaluate oxidative status,
45 ferrous oxidation-xylenol orange (FOX), thiobarbituric acid reactive substances
46 (TBARS) and reactive oxygen species production (ROS) concentrations in serum were
47 determined.

48

49 Mean concentrations of all antioxidant biomarkers analysed, with exception of
50 FRAP, were significantly lower ($P < 0.0001$) in the sera of dogs with IBD than in
51 healthy dogs. The oxidant markers studied were significantly higher ($P < 0.0001$) in sera
52 of dogs with IBD than in healthy dogs. These findings support the hypothesis that
53 oxidative stress could play an important role in the pathogenesis of canine IBD.

54

55 *Keywords:* Antioxidant; Cupric; PON1; Reactive oxygen species; Thiol

Accepted Manuscript

56 **Introduction**

57 Idiopathic inflammatory bowel disease (IBD) is characterised by persistent or
58 recurrent activation of the mucosal immune system accompanied by infiltrations of
59 inflammatory cells in the intestinal mucosa (Allenspach et al., 2007; Simpson and
60 Jergens, 2011). It is the most common cause of chronic intestinal disease in dogs, and
61 results in diverse and often debilitating clinical signs (Jergens et al., 2003; Allenspach et
62 al., 2016).

63
64 The pathogenesis of IBD in dogs is not completely understood; however it is
65 believed that intestinal inflammation results from a dysregulated immune response to
66 intestinal antigens (Allenspach et al., 2010). There is evidence that oxidative stress
67 plays an important role in the pathogenesis of IBD in human patients particularly in the
68 initiation and perpetuation of inflammation and in subsequent tissue damage. Oxidative
69 stress occurs when there is a marked imbalance between the production of reactive
70 oxygen species (ROS) and their removal by antioxidants (Rezaie et al., 2007). Recent
71 studies suggest that oxidative stress could also represent a significant factor in the
72 pathogenesis of IBD in dogs. One study that evaluated the metabolomics profile in dogs
73 with IBD using an untargeted metabolomics approach suggested the presence of
74 oxidative stress and a functional alteration of the GI microbiota in dogs with IBD,
75 which persisted even in the face of a clinical response to medical therapy (Minamoto et
76 al., 2014). Other recent studies have reported that various serum antioxidant biomarkers,
77 such as Trolox equivalent antioxidant capacity (TEAC), cupric reducing antioxidant
78 capacity (CUPRAC) and paraoxonase 1 (PON1), were decreased in the sera of dogs
79 with IBD (Rubio et al., 2016a,b; Segarra et al., 2016), again suggesting that oxidative
80 stress could play an important role in the pathogenesis of canine IBD.

81

82 The purpose of this study was to evaluate and compare a panel of various serum
83 biomarkers evaluating both the antioxidant response and oxidative damage response in
84 sera from dogs with IBD. These included previously the described antioxidant serum
85 biomarkers TEAC, CUPRAC and PON1, and additional antioxidants that have not
86 previously been studied in canine IBD, such as ferric reducing ability of plasma (FRAP)
87 and total serum thiol concentrations. To investigate oxidative damage, we measured
88 ferrous oxidation-xylene orange (FOX) and thiobarbituric acid reactive substances
89 (TBARS), which measures products of lipid peroxidation in serum. Finally, we
90 measured serum concentrations of reactive oxygen species.

91

92 **Material and methods**

93 *Animals*

94 In this retrospective study, a group of 18 dogs diagnosed with IBD at the Royal
95 Veterinary College (RVC), London, were included. Dogs with a history typical for
96 chronic enteropathy (≥ 3 weeks of vomiting, diarrhea or both, with or without weight
97 loss) were included. The diagnosis of chronic enteropathy was confirmed based on
98 established criteria (no clinically relevant abnormalities on routine hematology, serum
99 biochemistry; trypsin-like immunoreactivity [TLI], canine pancreatic lipase [cPL] and
100 adrenocorticotrophic hormone [ACTH]-stimulation test results within the reference
101 ranges; no abnormalities on abdominal imaging [radiographs, abdominal ultrasound
102 examination or both]; Allenspach et al., 2007). The histopathological criteria for IBD
103 were based on the guidelines for evaluation of gastrointestinal inflammation in
104 companion animals, as established by Washabau et al. (2010). In all cases biopsies were
105 obtained and reviewed by a board-certified pathologist.

106

107 In addition, 20 clinically healthy dogs were included in this study as a control
108 group.

109

110 Archived sera from healthy dogs and those with IBD and biopsy samples were
111 originally obtained in 2007 for diagnostic purposes only and were residual samples,
112 stored and available in the RVC archive. They were originally obtained with informed
113 owner consent under the Veterinary Surgeons Act (residual clause) and approved by the
114 RVC Ethics and Welfare Committee and were frozen immediately after blood sampling
115 and stored at -80 °C until further analysis.

116

117 *Antioxidant capacity*

118 The TEAC assay, based on the inhibition of the radical ABTS by the sample,
119 was performed according to the assay described by Arnao et al. (1996) validated by
120 Rubio et al. (2016a). The CUPRAC assay, based on the capacity of the sample in
121 reducing Cu(II) to Cu(I), was performed as previously described for use in canine serum
122 (Rubio et al., 2016b). The FRAP assay was performed following the method by Benzie
123 and Strain (1996) which measures the ferric to ferrous ion reduction by the sample.

124

125 Serum thiol was determined according to the method by Jocelyn (1987), and
126 serum PON1 activity was analysed following a previously described method for use in
127 canine serum (TvariJonaviciute et al., 2012).

128

129 All analyses were performed at Murcia University using an automated
130 biochemistry analyser (Olympus AU600 Automatic Chemistry Analyser, Olympus). All

131 assays showed inter- and intra-assay imprecision of <15%. Lower detection limits of
132 TEAC, CUPRAC and FRAP were 0.090, 0.003, and 0.031 mmol/L, respectively. The
133 lower detection limit of serum thiol and PON1 were 4.0 $\mu\text{mol/L}$ and 0.6 U/mL,
134 respectively.

135

136 *Oxidant biomarkers*

137 The FOX assay was based on the automated method described by Arab and
138 Steghens (2004) and performed using the Olympus AU600 Automatic Chemistry
139 Analyser (Olympus). The TBARS assay was determined following the method by
140 Buege and Aust (1978) using a microplate reader (Powerwave XS, Biotek instruments).

141

142 Reactive oxygen species (ROS) were estimated by luminol-mediated
143 chemiluminescence assay (Vong et al., 2014) using a microplate reader (Victor 2 1420
144 Multilabel Counter; PerkinElmer, Finland) and expressed in counts per second (cps).
145 All oxidant assays showed inter- and intra-assay imprecision less than 15% when
146 evaluated in our laboratory. In addition, the lower detection limits of FOX and TBARS
147 were 55.91 and 0.81 $\mu\text{mol/L}$, respectively. The ROS assay showed a lower detection
148 limit of of 1,300 cps.

149

150 *Statistical analysis*

151 Data were analysed using Graphpad Prism software (version 5 for Windows).
152 Concentrations of antioxidants and oxidant biomarkers were compared between dogs
153 with IBD and healthy control dogs. The results for each parameter were evaluated for
154 normality using the Shapiro-Wilk test. Thiol, PON1, TBARS and ROS results were not
155 normally distributed, therefore they were presented as median and interquartile range

156 (IQR). Normally distributed data were presented as means \pm standard deviation. We
157 determined statistical differences between healthy dogs and dogs with IBD using
158 unpaired t test (normally distributed data) and Mann Whitney U test (not normally
159 distributed data). Correlations between variables were determined using Spearman
160 correlation analysis. A *P*-value (two-tailed) of <0.05 was taken as statistically
161 significant in all cases.

162

163 **Results**

164 *Animals*

165 The 18 dogs with IBD included Border collie ($n=1$), Boxer ($n=3$), Cocker
166 spaniel ($n=2$), East-European shepherd ($n=1$), Greyhound ($n=1$), Labrador ($n=2$), Mixed
167 breed ($n=1$), Old English sheepdog ($n=1$), Polish Lowland sheepdog ($n=1$), Rottweiler
168 ($n=1$), Schnauzer ($n=1$), Staffordshire terrier ($n=2$), West Highland white terrier ($n=1$).
169 Their ages ranged from 7 months to 11 years; eight were females ($n=6$ spayed; $n=2$
170 intact) and 10 were males ($n=4$ neutered; $n=6$ intact). The clinical status of each dog
171 was evaluated at the time of diagnosis using the canine chronic enteropathy clinical
172 activity index (CCECAI) scoring system established by Allenspach et al. (2007), which
173 is based on nine variables, including attitude and activity, appetite, vomiting, stool
174 consistency, stool frequency, weight loss, ascites, pruritus and serum albumin
175 concentration. Based on CCECAI, the disease was mild (score 4-5) in two dogs,
176 moderate (score 6-8) in 11 dogs, severe (score 9-11) in two dogs, and very severe (score
177 ≥ 12) in three dogs.

178

179 Histopathologic findings of intestinal mucosal biopsies showed in all cases a
180 lympho-plasmacytic inflammation, in some cases with eosinophils. There was no
181 neutrophilic or macrophagic inflammation in any case.

182

183 The 20 control dogs consisted of various breeds (Mixed breed, Staffordshire
184 terrier, Italian Spinone, Labrador, Cavalier King Charles spaniel, English Springer
185 spaniel, Akita Inu, Golden retriever, Border collie, Dogue de Bordeaux, Rottweiler,
186 Great Dane, Japanese Spitz, Leonberger). Their ages ranged from 2 to 13 years; seven
187 were female ($n=6$ spayed; $n=1$ intact) and 13 were male ($n=11$ neutered; $n=2$ intact).

188

189 *Antioxidant response*

190 Dogs with IBD had lower TEAC concentrations (0.35 ± 0.06 vs. 0.51 ± 0.04
191 mmol/L; $P < 0.0001$), lower CUPRAC concentrations (0.3 ± 0.05 vs. 0.44 ± 0.05 mmol/L;
192 $P < 0.0001$), lower serum thiol concentrations (median, 63; IQR, 34-94 vs. median, 245;
193 IQR, 216-277 $\mu\text{mol/L}$; $P < 0.0001$), and lower PON1 activity than control dogs (median,
194 2.2; IQR, 1.7-2.8 vs. median, 3.5; IQR, 3.1-3.8 IU/mL; $P < 0.0001$; Fig. 1).

195

196 However, serum FRAP did not differ between dogs with IBD and control dogs
197 (0.41 ± 0.09 vs. 0.42 ± 0.07 mmol/L; $P = 0.6459$).

198

199 *Oxidant biomarkers*

200 Dogs with IBD had higher serum ROS counts (median, 8,861; IQR, 5,900-
201 14,763 cps vs. median, 1,502; IQR, 1,384-1,648 cps; $P < 0.0001$), higher FOX
202 concentrations (148 ± 65 vs. 72 ± 13 $\mu\text{mol/L}$; $P < 0.0001$) and higher TBARS

203 concentrations than control dogs (median, 8.0; IQR, 5.9-10.5 vs. median, 2.4; IQR: 1.9-
204 2.8 $\mu\text{mol/L}$; $P < 0.0001$; Fig. 2).

205

206 *Correlation study*

207 We observed correlations between all antioxidant biomarkers except for FRAP
208 and TEAC, and FRAP and CUPRAC (Table 1). The highest correlations ($\rho > 0.90$)
209 were observed between TEAC, CUPRAC and serum thiol.

210

211 Similarly, all oxidant biomarkers were positively correlated, with the highest
212 coefficient of correlation being between TBARS and ROS ($\rho = 0.84$; $P < 0.001$). In
213 addition, TBARS and ROS correlated negatively with TEAC, CUPRAC, and PON1 (all
214 $\rho \leq 0.45$; $P < 0.01$). A negative correlation was also observed between all oxidant
215 biomarkers and serum thiol (all $\rho \leq 0.54$; $P < 0.001$).

216

217 The CCECAI was not correlated with any of the biomarkers studied (all P
218 ≥ 0.05).

219

220 **Discussion**

221 Our study, evaluating the oxidative stress in dogs with IBD compared to healthy
222 control dogs using a comprehensive panel of serum biomarkers, suggests increased
223 oxidative stress status in canine IBD.

224

225 Serum TEAC and CUPRAC were significantly reduced in dogs with IBD
226 compared with healthy dogs. These results corroborate recent studies in dogs (Rubio et
227 al., 2016a,b; Segarra et al., 2016). Furthermore, Rubio et al. (2016a) observed similar

228 decreases (about 30%) in the TEAC concentration in dogs with IBD, as we did in the
229 present study. In this cohort of dogs, dogs with IBD had mean CUPRAC serum
230 concentrations that were 33% lower than in healthy dogs, compared to a 17% decrease
231 observed by Rubio et al. (2016b). Decreased TEAC and individual antioxidant (biotin,
232 folate, β -carotene, and vitamins A, C, and B) concentrations have previously been
233 reported in human patients with ulcerative colitis (UC; Fernandez-Banares et al., 1989;
234 Geerling, 1999; Aslan et al., 2011). The decreased antioxidant response in dogs with
235 IBD could be due to severe, persistent oxidative stress that depletes antioxidant
236 resources and overtakes the ability of the body to produce more antioxidants (Rezaie et
237 al., 2007). We failed to detect a difference in serum FRAP between healthy dogs and
238 dogs with IBD. This might be explained by the different individual antioxidants that
239 contribute to serum FRAP compared with other total antioxidant capacity (TAC) assays.
240 Ascorbic acid, α -tocopherol and principally uric acid are the contributors to FRAP in
241 humans, whereas thiols and albumin are the main contributors to CUPRAC and TEAC.
242 Therefore, we recommend that several different methods be used to measure TAC in
243 biological samples, because individual assays have different biochemical bases,
244 resulting in different results and interpretations (Cao and Prior, 1998; Hettyey et al.,
245 2007; Jansen and Ruskovska, 2013).

246
247 Studies have shown that mucosal thiol proteins are targets of oxidative injury
248 (Grisham et al., 1990). In the present study, serum thiol and PON1 were also
249 determined as individual antioxidants and both were significantly diminished in dogs
250 with IBD. Paraoxonase 1 (PON1) is widely distributed among tissues, including the
251 intestine. This protein is considered to be an antioxidant enzyme as it hydrolyses lipid
252 peroxides, in addition to its anti-inflammatory role in disease (Ceron et al., 2014). One

253 of the explanations for decreased PON1 activity could be its inactivation due to an
254 exacerbated oxidative environment (Nguyen and Sok, 2003). Our results are in
255 agreement with other studies that reported diminished PON1 activity in dogs with IBD
256 and human patients with IBD (Baskol et al., 2006; Boehm et al., 2009; Segarra et al.,
257 2016). In addition, there is evidence that diminished PON1 activity could also be a
258 consequence of a reduced number of the free SH groups on PON1, because of
259 exposition to oxygen free radicals (Jaouad et al., 2006), which could explain the high
260 correlation between serum PON1 activity and total serum thiol found in this study.

261

262 We demonstrated increased ROS and products of lipid peroxidation in the sera
263 of dogs with IBD by TBARS and FOX assays. These results are in agreement with
264 reports of IBD in humans (Levy et al., 2000; Sampietro et al., 2002; Baskol et al.,
265 2006). It has been reported that phagocytic cells (neutrophils and macrophages) isolated
266 from inflamed intestinal tissue in human patients with IBD release large amounts of
267 ROS during stimulation that could act locally or be secreted into the circulation to
268 produce different systemic effects (Kitahora et al., 1988; Alzoghaibi, 2013). However,
269 in our study, inflamed intestinal mucosa in the dogs with IBD contained lymphocytes
270 and plasma cells, which could suggest that these cells might also be a source of
271 systemic ROS, as has been suggested in humans (Lantow et al., 2006). Nevertheless,
272 studies using larger group sizes are needed to evaluate the association between the type
273 of cells detected in the inflamed mucosa of the dogs with IBD and the systemic
274 concentrations of oxidants (ROS and lipid peroxides) during the disease.

275

276 The relatively low number of dogs with IBD in our study might have limited our
277 ability to detect correlations between the various serum biomarkers and clinical disease

278 activity. However, our data show that antioxidants, which act to control oxidative stress,
279 are decreased, and oxidants generated in association with oxidative stress are increased
280 in dogs with IBD. These findings could help explain the pathophysiology of the disease.
281 However, further studies with larger group sizes are necessary to evaluate the potential
282 of these markers as possible predictors of disease and biomarkers for treatment
283 monitoring.

284

285 Although we did not determine the stability of the oxidant and antioxidants we
286 measured in canine sera, previous reports demonstrated that antioxidants were stable in
287 human serum stored at -80 °C for at least 1 year (Jansen et al., 2013). In addition, the
288 values of the control group were inside the reference values of our laboratory,
289 determined with fresh samples. However, possible changes during the long storage of
290 the samples in any of the analytes measured cannot be disregarded and this should be
291 considered as a limitation of our study.

292

293 **Conclusions**

294 Our study demonstrated the presence of oxidative stress in dogs with IBD. Based
295 on our results, a profile including biomarkers of total antioxidant status such as TEAC
296 and CUPRAC, individual antioxidant biomarkers such as PON1 and thiol, and
297 biomarkers of oxidant status measuring lipid peroxidation, such as FOX, or TBARS, or
298 directly measuring ROS production, might be useful in the comprehensive evaluation of
299 the oxidative stress response in dogs with IBD.

300

301 **Conflict of interest statement**

302 The authors have no financial and personal relationships with people or
303 organisations that could have inappropriately influenced his work.

304

305 **Acknowledgment**

306 The authors would like to thank the staff of the Clinical Investigation Center at
307 the RVC for their help in sample acquisition and storage.

308

309 **Funding**

310 The National Council for Scientific and Technological Development (CNPq) of
311 Brazil funded the PhD scholarship of CPR. The Ministerio de Economía y
312 Competitividad of Spain funded AT through a postdoctoral grant (Juan de la Cierva).
313 This study was also funded by the Seneca Foundation of Murcia Region
314 (19894/GERM/15).

315

316 **References**

- 317 Allenspach, K., Culverwell, C., Chan, D., 2016. Long-term outcome in dogs with
318 chronic enteropathies: 203 cases. *Veterinary Record* 178, 368-368.
- 319
- 320 Allenspach, K., House, A., Smith, K., McNeill, F.M., Hendricks, A., Elson-Riggins, J.,
321 Riddle, A., Steiner, J.M., Werling, D., Garden, O.A., Catchpole, B., Suchodolski,
322 J.S., 2010. Evaluation of mucosal bacteria and histopathology, clinical disease
323 activity and expression of Toll-like receptors in German shepherd dogs with
324 chronic enteropathies. *Veterinary Microbiology* 146, 326–335.
- 325
- 326 Allenspach, K., Wieland, B., Gröne, A., Gaschen, F., 2007. Chronic enteropathies in
327 dogs: evaluation of risk factors for negative outcome. *Journal of Veterinary*
328 *Internal Medicine* 21, 700–708.
- 329
- 330 Alzogaibi, M.A., 2013. Concepts of oxidative stress and antioxidant defense in
331 Crohn's disease. *World Journal of Gastroenterology* 19, 6540-6547.
- 332
- 333 Arab, K., Steghens, J. P., 2004. Plasma lipid hydroperoxides measurement by an
334 automated xylenol orange method. *Analytical Biochemistry* 325, 158-163.
- 335
- 336 Arnao, M.B., Cano, A., Hernandez-Ruiz, J., Garcia-Cánovas, F., Acosta, M., 1996.
337 Inhibition by l-ascorbic acid and other antioxidants of the 2,2'-azino-bis(3-

- 338 ethylbenzthiazoline-6-sulfonic acid) oxidation catalyzed by peroxidase: a new
339 approach for determining total antioxidant status of foods. *Analytical Biochemistry*
340 236, 255–261.
- 341
- 342 Aslan, M., Nazligu, Y., Bolukbas, C., Bolukbas, F.F., Horoz, M., Dulger, A.C., Erdur,
343 F.M., Celik, H., Kocyigit, A., 2011. Peripheral lymphocyte DNA damage and
344 oxidative stress in patients with ulcerative colitis. *Polish Archives of Internal*
345 *Medicine* 121, 223–229.
- 346
- 347 Benzie, I.F., Strain, J.J., 1996. The ferric reducing ability of plasma (FRAP) as a
348 measure of “antioxidant power”: the FRAP assay. *Analytical Biochemistry* 239,
349 70–76.
- 350
- 351 Baskol, G., Baskol, M., Yurci, A., Ozbakir, O., Yucesoy, M., 2006. Serum paraoxonase
352 1 activity and malondialdehyde levels in patients with ulcerative colitis. *Cell*
353 *Biochemistry and Function* 24, 283–286.
- 354
- 355 Boehm, D., Krzystek-Korpacka, M., Neubauer, K., Matusiewicz, M., Berdowska, I.,
356 Zielinski, B., Paradowski, L., Gamian, A., 2009. Paraonase-1 status in Crohn’s
357 disease and ulcerative colitis. *Inflammatory Bowel Diseases* 15, 93–99.
- 358
- 359 Buege, J. A., Aust, S. D., 1978. [30] Microsomal lipid peroxidation. *Methods in*
360 *Enzymology* 52, 302-310.
- 361
- 362 Cao, G., Prior, R.L., 1998. Comparison of different analytical methods for assessing
363 total antioxidant capacity of human serum. *Clinical Chemistry* 44, 1309–1315.
- 364
- 365 Ceron, J.J., Tecles, F., Tvarijonaviciute, A., 2014. Serum paraoxonase 1 (PON1)
366 measurement: an update. *BMC Veterinary Research* 10, 74.
- 367
- 368 Fernandez-Banares, F., Abad-Lacruz, A., Xiol, X., Gine, J.J., Dolz, C., Cabre, E.,
369 Esteve, M., Gonzalez-Huix, F., Gassull, M.A., 1989. Vitamin status in patients
370 with inflammatory bowel disease. *American Journal of Gastroenterology* 84.
- 371
- 372 Geerling, B.J., 1999. The relation between antioxidant status and alterations in fatty acid
373 profile in patients with Crohn disease and controls. *Scandinavian Journal of*
374 *Gastroenterology* 34, 1108–1116.
- 375
- 376 Grisham, M.B., Gaginella, T.S., von Ritter, C., Tamai, H., Robert, M.B., Granger, D.N.,
377 1990. Effects of neutrophil-derived oxidants on intestinal permeability, electrolyte
378 transport, and epithelial cell viability. *Inflammation* 14, 531–542.
- 379
- 380 Heteyey, C.S., Manczur, F., Dudás-Györki, Z., Reiczigel, J., Ribiczey, P., Vajdovich, P.,
381 Vörös, K., 2007. Plasma antioxidant capacity in dogs with naturally occurring
382 heart diseases. *Journal of Veterinary Medicine series* 54, 36–39.
- 383
- 384 Jansen, E.H., Beekhof, P.K., Cremers, J.W.J.M., Viezeliene, D., Muzakova, V.,
385 Skalicky, J., 2013. Long-term stability of parameters of antioxidant status in
386 human serum. *Free Radical Research* 47, 535-540.
- 387

- 388 Jansen, E.H., Ruskovska, T., 2013. Comparative analysis of serum (anti)oxidative status
389 parameters in healthy persons. *International Journal of Molecular Sciences* 14,
390 6106–6115.
391
- 392 Jaouad, L., Guise, C. de, Berrougui, H., 2006. Age-related decrease in high-density
393 lipoproteins antioxidant activity is due to an alteration in the PON1's free
394 sulfhydryl groups. *Atherosclerosis* 185, 191-200.
395
- 396 Jergens, A.E., Schreiner, C.A., Frank, D.E., Niyo, Y., Ahrens, F.E., Eckersall, P.D.,
397 Benson, T.J., Evans, R., 2003. A scoring index for disease activity in canine
398 inflammatory bowel disease. *Journal of Veterinary Internal Medicine* 17, 291-297.
399
- 400 Jocelyn, P., 1987. Spectrophotometric assay of thiols. *Methods in Enzymology* 143, 44-
401 67.
402
- 403 Kitahora, T., Suzuki, K., Asakura, H., Yoshida, T., Suematsu, M., Watanabe, M., Aiso,
404 S., Tsuchiya, M., 1988. Active oxygen species generated by monocytes and
405 polymorphonuclear cells in Crohn's disease. *Digestive diseases and Sciences* 33,
406 951–955.
407
- 408 Lantow, M., Lupke, M., Frahm, J., Mattsson, M.O., Kuster, N., Simko, M., 2006. ROS
409 release and Hsp70 expression after exposure to 1,800 MHz radiofrequency
410 electromagnetic fields in primary human monocytes and lymphocytes. *Radiation
411 and Environmental Biophysics*, 45, 55-62.
412
- 413 Levy, E., Rizwan, Y., Thibault, L., Lepage, G., Brunet, S., Bouthillier, L., Seidman, E.,
414 2000. Altered lipid profile, lipoprotein composition, and oxidant and antioxidant
415 status in pediatric Crohn disease. *The American Journal of Clinical Nutrition* 71,
416 807–815.
417
- 418 Minamoto, Y., Otoni, C.C., Steelman, S.M., Büyükleblebici, O., Steiner, J.M., Jergens,
419 A.E., Suchodolski, J.S., 2014. Alteration of the fecal microbiota and serum
420 metabolite profiles in dogs with idiopathic inflammatory bowel disease. *Gut
421 Microbes* 6, 33-47.
422
- 423 Nguyen, S.D., Sok, D.E., 2003. Oxidative inactivation of paraoxonase1, an antioxidant
424 protein and its effect on antioxidant action. *Free Radical Research* 37, 1319–1330.
425
- 426 Rezaie, A., Parker, R.D., Abdollahi, M., 2007. Oxidative stress and pathogenesis of
427 inflammatory bowel disease: an epiphenomenon or the cause? *Digestive Diseases
428 and Sciences* 52, 2015–21.
429
- 430 Rubio, C.P., Hernández-Ruiz, J., Martínez-Subiela, S., Tvarijonaviciute, A., Arnao,
431 M.B., Ceron, J.J., 2016a. Validation of three automated assays for total antioxidant
432 capacity (TAC) determination in canine serum samples. *Journal of Veterinary
433 Diagnostic Investigation* 1040638716664939.
434
- 435 Rubio, C.P., Tvarijonaviciute, A., Martínez-Subiela, S., Hernández-Ruiz, J., Ceron, J.J.,
436 2016b. Validation of an automated assay for the measurement of cupric reducing
437 antioxidant capacity in serum of dogs. *BMC Veterinary Research* 12, 1.

- 438
439 Sampietro, G.M., Cristaldi, M., Cervato, G., Maconi, G., Danelli, P., Cervellione, R.,
440 Rovati, M., Bianchi Porro, G., Cestaro, B., Taschieri, A.M., 2002. Oxidative stress,
441 vitamin A and vitamin E behaviour in patients submitted to conservative surgery
442 for complicated Crohn's disease. *Digestive and Liver Disease* 34, 696–701.
443
- 444 Segarra, S., Martinez-Subiela, S., Cerda-Cuellar, M., Martinez-Puig, D., Munoz-Prieto,
445 A., Rodriguez-Franco, F., Rodriguez-Bertos, A., Allenspach, K., Velasco, A.,
446 Ceron, J., 2016. Oral chondroitin sulfate and prebiotics for the treatment of canine
447 Inflammatory Bowel Disease: a randomized, controlled clinical trial. *BMC*
448 *Veterinary Research* 12, 49.
449
- 450 Simpson, K.W., Jergens, A.E., 2011. Pitfalls and progress in the diagnosis and
451 management of canine inflammatory bowel disease. *Veterinary Clinics of North*
452 *America: Small Animal Practice* 41, 381–98.
453
- 454 Tvarijonaviciute, A., Tecles, F., Caldin, M., Tasca, S., Cerón, J., 2012. Validation of
455 spectrophotometric assays for serum paraoxonase type-1 measurement in dogs.
456 *American Journal of Veterinary Research* 73, 34–41.
457
- 458 Vong, L., Lorentz, R., Assa, A., 2014. Probiotic *Lactobacillus rhamnosus* inhibits the
459 formation of neutrophil extracellular traps. *The Journal of Immunology* 192,
460 1870-1877.
461
- 462 Washabau, R.J., Day, M.J., Willard, M.D., Hall, E.J., Jergens, A.E., Mansell, J.,
463 Minami, T., Bilzer, T.W., 2010. Endoscopic, biopsy, and histopathologic
464 guidelines for the evaluation of gastrointestinal inflammation in companion
465 animals. *Journal of Veterinary Internal Medicine* 24, 10-26.
466

467

468 Fig. 1. Comparisons of antioxidant biomarkers in healthy dogs and dogs with IBD. The
469 plots show median (line within box), 25th and 75th percentiles (box) and minimum and
470 maximum values (whiskers). TEAC, Trolox equivalent antioxidant capacity; CUPRAC,
471 cupric reducing antioxidant capacity; FRAP, ferric reducing ability of plasma; PON1,
472 paraoxonase 1.

473

474 Fig. 2. Comparisons of oxidant biomarkers in healthy dogs and dogs with IBD. The
475 plots show median (line within box), 25th and 75th percentiles (box) and minimum and
476 maximum values (whiskers). ROS, reactive oxygen species; FOX, ferrous oxidation-
477 xyleneol orange; TBARS, thiobarbituric acid reactive substances.

Accepted Manuscript

478 **Table 1.** Spearman correlation coefficients, levels of statistical significance (95%
 479 confidence intervals) between all biomarkers studied.

	TEAC	CUPRA C	FRAP	Thiol	PON1	FOX	TBARS
CUP	0.92***						
RAC	(0.84 to 0.95)						
FRAP	0.26 (- 0.07 to 0.55)	0.27 (- 0.07 to 0.55)					
Thiol	0.90*** (0.81 to 0.95)	0.90*** (0.81 to 0.95)	0.04 (- 0.30 to 0.37)				
PON1	0.68*** (0.45 to 0.82)	0.72*** (0.51 to 0.85)	0.34* (0.01 to 0.61)	0.65*** (0.41 to 0.81)			
FOX	-0.40* (-0.65 to -0.08)	-0.39* (-0.64 to -0.06)	0.46** (0.16 to 0.69)	-0.54*** (-0.74 to -0.26)	-0.20 (-0.50 to 0.13)		
TBA	-0.54*** (-0.74 to -0.26)	-0.59*** (-0.77 to -0.32)	0.12 (-0.22 to -0.44)	-0.65*** (-0.81 to -0.41)	-0.45** (-0.68 to -0.14)	0.74*** (0.54 to -0.86)	
ROS	-0.70*** (-0.84 to -0.49)	-0.77*** (-0.88 to -0.58)	-0.07 (-0.40 to -0.27)	-0.77*** (-0.88 to -0.60)	-0.65*** (-0.81 to -0.40)	0.61*** (0.35 to 0.78)	0.84*** (0.70 to 0.91)

480 * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$.

481 CUPRAC, cupric reducing antioxidant capacity; FRAP, ferric reducing ability of
 482 plasma; FOX, ferrous oxidation-xylenol orange; PON1, paraoxonase 1; ROS, reactive
 483 oxygen species; TBARS, thiobarbituric acid reactive substances; TEAC, Trolox
 484 equivalent antioxidant capacity.

485

Accepted Manuscript