

This is the peer-reviewed, manuscript version of the following article:

Volk, H. A. (2016) 'Canine epilepsy: separating the wood from the trees', Veterinary Record, 178(16), 394-396.

The final version is available online via <http://dx.doi.org/10.1136/vr.i1999>.

The full details of the published version of the article are as follows:

TITLE: Canine epilepsy: separating the wood from the trees

AUTHORS: H.A. Volk

JOURNAL TITLE: Veterinary Record

VOLUME/EDITION: 178/16

PUBLISHER: BMJ Publishing Group

PUBLICATION DATE: April 2016

DOI: 10.1136/vr.i1999

## **Canine epilepsy: separating the wood from the trees**

Volk HA

Department of Clinical Science and Services, Royal Veterinary College, Hatfield, AL9 7TA,  
Hertfordshire, UK (Holger Volk [hvolk@rvc.ac.uk](mailto:hvolk@rvc.ac.uk))

**Corresponding author:** Holger A Volk, Department of Clinical Science and Services, Royal  
Veterinary College, Hatfield, Hertfordshire, UK; [hvolk@rvc.ac.uk](mailto:hvolk@rvc.ac.uk)

**Key words:** Epilepsy, Seizures, Dog, Classification, Semiology

## 1 Editorial

2 Every 100th patient appearing on the doorstep of a busy first opinion practice will be  
3 presented with seizures and two thirds of these patients will have epilepsy (Heske and  
4 others 2014; Kearsley-Fleet and others 2013). Despite the frequency in occurrence and a  
5 plethora of studies been published in the field, the clinician remains puzzled of what  
6 terminology is the correct one to use, how to diagnose the different types of epilepsy, how  
7 and when to treat best and when change in treatment is necessary?

8 In 2015, a group of Veterinary Neurology Specialists and Non-specialists came together to  
9 form the International Veterinary Epilepsy Task Force (IVETF). The IVETF's main aim is "to  
10 provide the veterinary community, breeders and dog (and in part cat) owners with  
11 consensus statements on the key areas in the field of epilepsy"(Volk 2015). The IVETF wants  
12 to represent the "chain of care" involving a broad range of stakeholders (veterinary and  
13 human neurologists and neuroscientists, practitioners, neuropharmacologists and  
14 neuropathologists) to ensure that the produced consensus statements are pragmatic and  
15 'user-friendly' for daily use. More than 25 co-authors were involved in the process of  
16 developing seven consensus statements:

1. International Veterinary Epilepsy Task Force consensus report on epilepsy  
definition, classification and terminology in companion animals (Berendt and others  
2015)
- 17 2. International Veterinary Epilepsy Task Force Consensus Proposal: Diagnostic  
18 approach to epilepsy in dogs (De Risio and others 2015)
- 19 3. International Veterinary Epilepsy Task Force current understanding of idiopathic  
20 epilepsy of genetic or suspected genetic origin in purebred dogs (Huelsmeyer and  
21 others 2015)
- 22 4. International Veterinary Epilepsy Task Force consensus proposal: Medical treatment  
23 of canine epilepsy in Europe (Bhatti and others 2015)

5. International Veterinary Epilepsy Task Force Consensus Proposal: Outcome of therapeutic interventions in canine and feline epilepsy (Potschka and others 2015)
6. International Veterinary Epilepsy Task Force recommendations for a veterinary epilepsy-specific MRI protocol (Rusbridge and others 2015)
7. International Veterinary Epilepsy Task Force recommendations for systematic sampling and processing of brains from epileptic dogs and cats (Matiasek and others 2015)

In 2016, a complimentary consensus statement about seizure management was published under the umbrella of the American College of Veterinary Internal Medicine (ACVIM) (Podell and others 2016). The ACVIM's and IVETF's consensus statements are based on evidence-based medicine, but also consider collective expertise where such evidence is conflicting or lacking. All the consensus statements are freely available online for the interested reader<sup>1</sup>. A potted version of some of the highlights can be found below:

#### **What terminology is the correct one to use (Berendt and others 2015)?**

How best to talk the talk in Epilepsy - the IVETF proposed to differentiate between the term seizure and epileptic seizure. The term seizure can be used for any sudden occurring, brief and transient episode and does not necessarily imply that the event is epileptic. It is therefore better to use the term epileptic seizure when you are certain of the nature of the episode. Epileptic seizures are defined as "manifestations of excessive synchronous, usually self-limiting epileptic activity of neurons in the brain. This results in a transient occurrence of signs which may be characterised by short episodes with convulsions or focal motor, autonomic or behavioural features and due to abnormal excessive and/or synchronous epileptic neuronal activity in the brain" (Berendt and others 2015).

---

<sup>1</sup> [http://www.biomedcentral.com/bmcvetres/series/IVETF\\_consensus\\_reports](http://www.biomedcentral.com/bmcvetres/series/IVETF_consensus_reports);  
<http://onlinelibrary.wiley.com/doi/10.1111/jvim.13841/abstract>

47 Epileptic seizures can be differentiated into focal and generalised epileptic seizures, and  
48 focal epileptic seizures can evolve into generalised epileptic seizures. Focal epileptic  
49 seizures originate within a neural network of one cerebral hemisphere. The onset and the  
50 resulting clinical signs are consistent from one epileptic seizure to another. An example  
51 could be the animal presents with epileptic seizures which always start with a twitching of  
52 its right facial muscles. Focal epileptic seizures are often asymmetrical in presentation. On  
53 the other hand, a generalised epileptic seizure is characterised by major involvement of both  
54 cerebral hemispheres. Epileptic seizures can further be described depending on their  
55 expression(s) of autonomic, motor or behavioural features.

56 The IVETF defined epilepsy conceptually as a brain disease which is “characterised by an  
57 enduring predisposition to generate epileptic seizures”, however, realised that a more  
58 practical definition was needed. The definition for epilepsy which can be used in daily  
59 practice is that epilepsy is defined as “at least two unprovoked epileptic seizures >24 h  
60 apart”. The IVETF proposed not to use the term epilepsy for epileptic seizures which are  
61 secondary or reactive to a metabolic or toxic transient disturbance in function, which when  
62 rectified will result in a cessation of epileptic activity. The term reactive seizures should be  
63 used for these.

64

## 65 **How to diagnose the different types of epilepsy (Berendt and others 2015; De Risio and** 66 **others 2015)?**

67 After the clinician has determined that the animal has recurrent epileptic seizures, the cause  
68 of the epileptic seizures need to be determined. Broadly speaking, the IVETF proposed to  
69 differentiate idiopathic epilepsy from structural epilepsy. The former term cryptogenic  
70 (possible symptomatic) epilepsy has been heavily discussed in human medicine, as some of  
71 them were identified later to be genetic in origin and this resulted human medicine to move  
72 away from this term and use the term epilepsy of unknown cause for epilepsy in which an

73 underlying cause could not be identified. This term was also listed in the IVETF statement  
74 as a “bucket term” for all the epilepsies which cannot be classified as idiopathic or structural  
75 epilepsy.

76 Structural epilepsy is caused by any disease which alters the brain structure such as  
77 vascular, inflammatory/infectious, traumatic, anomalous/developmental, neoplastic and  
78 degenerative diseases. After long debate the IVETF continues to recommend to use the term  
79 idiopathic epilepsy as it is well established. However, idiopathic epilepsy should be seen as  
80 a framework term which can be further differentiated into

- 81 1. “Idiopathic epilepsy (genetic epilepsy)—a causative gene for epilepsy has been  
82 identified/confirmed genetic background” (Berendt and others 2015).
- 83 2. “Idiopathic epilepsy (suspected genetic epilepsy)” – this applies for those cases  
84 where you have family or breed history of epilepsy (see also Huelsmeyer and others  
85 2015).
- 86 3. “Idiopathic epilepsy (epilepsy of unknown cause)—epilepsy in which the nature of  
87 the underlying cause is as yet unknown and with no indication of structural  
88 epilepsy”.

89

90 The IVETF also provided guidelines of how to confirm idiopathic epilepsy diagnostically  
91 and introduced a three tier level of confidence for the diagnosis of idiopathic epilepsy (De  
92 Risio and others 2015). In brief, the first tier can be diagnosed by any first opinion  
93 practitioner if an animal has “≥ 2 unprovoked epileptic seizures occurring ≥ 24h apart, age  
94 at epileptic seizure onset 6 months to 6 years, unremarkable inter-ictal physical and  
95 neurological examination, no clinically significant abnormalities on minimum database  
96 blood tests and urinalysis”. A family history of epilepsy adds strength to the diagnosis. The  
97 next level of confidence can be reached (tier 2) by having also an unremarkable dynamic

98 bile acid test, MRI (epilepsy-specific MRI protocol (Rusbridge and others 2015)) and CSF  
99 analysis. The highest level of confidence (tier 3) can be reached when in addition to the  
100 requirements fulfilled in tier 1 and 2 characteristic EEG changes are present.

101

102 **How and when to treat best and when change in treatment is necessary** (Bhatti and others  
103 2015; Podell and others 2016)?

104 The ACVIM's and IVETF's consensus statements agree in most parts of when medical  
105 treatment should be initiated. Treatment should be started when structural epilepsy is  
106 present, the period between seizures is equal or less than 6 months, the frequency of seizures  
107 is increasing over the last 3 inter-ictal periods, the animal had a status epilepticus, cluster  
108 seizures, severe or debilitating postictal signs. Both also agree that the evidence  
109 (Charalambous and others 2014; Charalambous and others 2016) is strongest for starting an  
110 otherwise healthy dog with epilepsy on Imepitoin or Phenobarbital. The IVETF, however,  
111 pointed out that in Europe, Imepitoin should only be prescribed for dogs with idiopathic  
112 epilepsy which have single generalised epileptic seizures.

113 The IVETF's aim is to ideally achieve seizure freedom or at least an "extension of the inter-  
114 seizure interval to three times the longest pre-treatment inter-seizure interval and for a  
115 minimum of three months"(Bhatti and others 2015; Potschka and others 2015). The IVETF  
116 does recognizes partial treatment success which is characterized by an at least 50%  
117 reduction in seizure frequency and/or seizure severity. It is known that a risk factor for  
118 poorer seizure control is a high seizure density (Packer and others 2014) and patients with  
119 cluster seizures might need to be treated more aggressively sooner. If treatment is not  
120 successful and/or treatment is not well tolerated treatment might need to be altered. Other  
121 factors apart from treatment success which might need to be considered when selecting a  
122 second anti-epileptic drug is the selection of a drug with a different mechanism of action,

123 potential harmful drug-interactions and a risk-benefit analysis of polypharmacy versus  
124 quality of life (Podell and others 2016).

125 The IVETF mainly recommends potassium bromide as an add-on medication, considering  
126 also local legal frameworks (Bhatti and others 2015). The ACVIM provides only a moderate  
127 level of recommendation of which drug best to chose from for second line treatment; Drugs  
128 mentioned are Phenobarbital, potassium bromide, Levetiracetam and Zonisamide (Podell  
129 and others 2016). The ACVIM also provides recommendations about non-pharmacological  
130 treatment options and the potential of novel diets for the management of epilepsy (Podell  
131 and others 2016), which might improve seizure control and reduce behavioural  
132 comorbidities(Law and others 2015; Packer and others 2016). Both groups highlight the  
133 importance of owner education and the importance of considering quality of life in the  
134 successful management of epilepsy.

135 In conclusion, as a busy clinician one can easily not see the forest for the trees in epilepsy.  
136 Epilepsy is a rather complex disease and can be challenging to diagnose and treat. The recent  
137 published consensus statements are useful resources to help give an expert overview of  
138 what is relevant.

139



## 140 **References**

- 141 BERENDT, M., FARQUHAR, R. G., MANDIGERS, P. J. J., PAKOZDY, A., BHATTI, S. F. M., DE RISIO, L., FISCHER,  
142 A., LONG, S., MATIASEK, K., MUNANA, K., PATTERSON, E. E., PENDERIS, J., PLATT, S., PODELL, M.,  
143 POTSCHKA, H., PUMAROLA, M. B., RUSBRIDGE, C., STEIN, V. M., TIPOLD, A. & VOLK, H. A. (2015)  
144 International veterinary epilepsy task force consensus report on epilepsy definition, classification and  
145 terminology in companion animals. *Bmc Veterinary Research* 11
- 146 BHATTI, S. F. M., DE RISIO, L., MUNANA, K., PENDERIS, J., STEIN, V. M., TIPOLD, A., BERENDT, M.,  
147 FARQUHAR, R. G., FISCHER, A., LONG, S., LOESCHER, W., MANDIGERS, P. J. J., MATIASEK, K., PAKOZDY, A.,  
148 PATTERSON, E. E., PLATT, S., PODELL, M., POTSCHKA, H., RUSBRIDGE, C. & VOLK, H. A. (2015) International  
149 Veterinary Epilepsy Task Force consensus proposal: medical treatment of canine epilepsy in Europe. *Bmc*  
150 *Veterinary Research* 11
- 151 CHARALAMBOUS, M., BRODBELT, D. & VOLK, H. A. (2014) Treatment in canine epilepsy—a systematic  
152 review. *Bmc Veterinary Research* 10, 257
- 153 CHARALAMBOUS, M., BRODBELT, D. & VOLK, H. A. (2016) The Evidence Behind the Treatment of Canine  
154 Idiopathic Epilepsy. *Veterinary Evidence* 1
- 155 DE RISIO, L., BHATTI, S., MUNANA, K., PENDERIS, J., STEIN, V., TIPOLD, A., BERENDT, M., FARQUHAR, R.,  
156 FISCHER, A., LONG, S., MANDIGERS, P. J., MATIASEK, K., PACKER, R. M., PAKOZDY, A., PATTERSON, N.,  
157 PLATT, S., PODELL, M., POTSCHKA, H., BATLLE, M. P., RUSBRIDGE, C. & VOLK, H. A. (2015) International  
158 veterinary epilepsy task force consensus proposal: diagnostic approach to epilepsy in dogs. *BMC Vet Res*  
159 11, 148
- 160 HESKE, L., NODTVEDT, A., JADERLUND, K. H., BERENDT, M. & EGENVALL, A. (2014) A cohort study of  
161 epilepsy among 665,000 insured dogs: incidence, mortality and survival after diagnosis. *Vet J* 202, 471-476
- 162 HUELSMEYER, V.-I., FISCHER, A., MANDIGERS, P. J. J., DERISIO, L., BERENDT, M., RUSBRIDGE, C., BHATTI, S.  
163 F. M., PAKOZDY, A., PATTERSON, E. E., PLATT, S., PACKER, R. M. A. & VOLK, H. A. (2015) International  
164 Veterinary Epilepsy Task Force's current understanding of idiopathic epilepsy of genetic or suspected  
165 genetic origin in purebred dogs. *Bmc Veterinary Research* 11
- 166 KEARSLEY-FLEET, L., O'NEILL, D. G., VOLK, H. A., CHURCH, D. B. & BRODBELT, D. C. (2013) Prevalence and  
167 risk factors for canine epilepsy of unknown origin in the UK. *The Veterinary record* 172, 338
- 168 LAW, T. H., DAVIES, E. S. S., PAN, Y., ZANGHI, B., WANT, E. & VOLK, H. A. (2015) A randomised trial of a  
169 medium-chain TAG diet as treatment for dogs with idiopathic epilepsy. *The British journal of nutrition* 114,  
170 1438-1447
- 171 MATIASEK, K., PUMAROLA I BATLLE, M., ROSATI, M., FERNANDEZ-FLORES, F., FISCHER, A., WAGNER, E.,  
172 BERENDT, M., BHATTI, S. F. M., DE RISIO, L., FARQUHAR, R. G., LONG, S., MUNANA, K., PATTERSON, E. E.,  
173 PAKOZDY, A., PENDERIS, J., PLATT, S., PODELL, M., POTSCHKA, H., RUSBRIDGE, C., STEIN, V. M., TIPOLD, A. &  
174 VOLK, H. A. (2015) International veterinary epilepsy task force recommendations for systematic sampling  
175 and processing of brains from epileptic dogs and cats. *Bmc Veterinary Research* 11
- 176 PACKER, R. M., LAW, T. H., DAVIES, E., ZANGHI, B., PAN, Y. & VOLK, H. A. (2016) Effects of a ketogenic diet  
177 on ADHD-like behavior in dogs with idiopathic epilepsy. *Epilepsy Behav* 55, 62-68
- 178 PACKER, R. M. A., SHIHAB, N. K., TORRES, B. B. J. & VOLK, H. A. (2014) Clinical Risk Factors Associated with  
179 Anti-Epileptic Drug Responsiveness in Canine Epilepsy. *Plos One* 9
- 180 PODELL, M., VOLK, H. A., BERENDT, M., LOSCHER, W., MUNANA, K., PATTERSON, E. E. & PLATT, S. R. (2016)  
181 2015 ACVIM Small Animal Consensus Statement on Seizure Management in Dogs. *J Vet Intern Med*

182 POTSCHKA, H., FISCHER, A., LOESCHER, W., PATTERSON, N., BHATTI, S., BERENDT, M., DE RISIO, L.,  
 183 FARQUHAR, R., LONG, S., MANDIGERS, P., MATIASEK, K., MUNANA, K., PAKOZDY, A., PENDERIS, J., PLATT,  
 184 S., PODELL, M., RUSBRIDGE, C., STEIN, V., TIPOLD, A. & VOLK, H. A. (2015) International veterinary epilepsy  
 185 task force consensus proposal: outcome of therapeutic interventions in canine and feline epilepsy. *Bmc*  
 186 *Veterinary Research* 11

187 RUSBRIDGE, C., LONG, S., JOVANOVIK, J., MILNE, M., BERENDT, M., BHATTI, S. F. M., DE RISIO, L.,  
 188 FARQUHAR, R. G., FISCHER, A., MATIASEK, K., MUNANA, K., PATTERSON, E. E., PAKOZDY, A., PENDERIS, J.,  
 189 PLATT, S., PODELL, M., POTSCHKA, H., STEIN, V. M., TIPOLD, A. & VOLK, H. A. (2015) International  
 190 Veterinary Epilepsy Task Force recommendations for a veterinary epilepsy-specific MRI protocol. *Bmc*  
 191 *Veterinary Research* 11

192 VOLK, H. A. (2015) International Veterinary Epilepsy Task Force consensus reports on epilepsy definition,  
 193 classification and terminology, affected dog breeds, diagnosis, treatment, outcome measures of  
 194 therapeutic trials, neuroimaging and neuropathology in companion animals. *Bmc Veterinary Research* 11

195  
 196  
 197