

This research was originally published in the Journal of Biological Chemistry:

Clive Bate, William Nolan, and Alun Williams. Sialic Acid on the Glycosylphosphatidylinositol Anchor Regulates PrP-mediated Cell Signaling and Prion Formation. *J. Biol. Chem.* 2016; Vol:291, pp.17093-17101. © the American Society for Biochemistry and Molecular Biology. <http://dx.doi.org/10.1074/jbc.M116.731117>.

# Sialic Acid on the Glycosylphosphatidylinositol Anchor Regulates PrP-mediated Cell Signaling and Prion Formation\*

Received for publication, June 13, 2015, and in revised form, November 5, 2015. Published, JBC Papers in Press, November 9, 2015, DOI 10.1074/jbc.M115.672394

Clive Bate<sup>‡1</sup>, William Nolan<sup>‡</sup>, and Alun Williams<sup>§</sup>

From the <sup>‡</sup>Department of Pathology and Pathogen Biology, Royal Veterinary College, North Mymms, Hertfordshire AL9 7TA, United Kingdom and the <sup>§</sup>Department of Veterinary Medicine, University of Cambridge, Cambridge CB3 0ES, United Kingdom

The prion diseases occur following the conversion of the cellular prion protein (PrP<sup>C</sup>) into disease-related isoforms (PrP<sup>Sc</sup>). In this study, the role of the glycosylphosphatidylinositol (GPI) anchor attached to PrP<sup>C</sup> in prion formation was examined using a cell painting technique. PrP<sup>Sc</sup> formation in two prion-infected neuronal cell lines (ScGT1 and ScN2a cells) and in scrapie-infected primary cortical neurons was increased following the introduction of PrP<sup>C</sup>. In contrast, PrP<sup>C</sup> containing a GPI anchor from which the sialic acid had been removed (desialylated PrP<sup>C</sup>) was not converted to PrP<sup>Sc</sup>. Furthermore, the presence of desialylated PrP<sup>C</sup> inhibited the production of PrP<sup>Sc</sup> within prion-infected cortical neurons and ScGT1 and ScN2a cells. The membrane rafts surrounding desialylated PrP<sup>C</sup> contained greater amounts of sialylated gangliosides and cholesterol than membrane rafts surrounding PrP<sup>C</sup>. Desialylated PrP<sup>C</sup> was less sensitive to cholesterol depletion than PrP<sup>C</sup> and was not released from cells by treatment with glimepiride. The presence of desialylated PrP<sup>C</sup> in neurons caused the dissociation of cytoplasmic phospholipase A<sub>2</sub> from PrP-containing membrane rafts and reduced the activation of cytoplasmic phospholipase A<sub>2</sub>. These findings show that the sialic acid moiety of the GPI attached to PrP<sup>C</sup> modifies local membrane microenvironments that are important in PrP-mediated cell signaling and PrP<sup>Sc</sup> formation. These results suggest that pharmacological modification of GPI glycosylation might constitute a novel therapeutic approach to prion diseases.

The prion diseases are fatal neurodegenerative disorders that include scrapie in sheep and goats, bovine spongiform encephalopathy in cattle, and Creutzfeldt-Jakob disease in man. A key event in these diseases is the conversion of a normal host protein designated PrP<sup>C</sup> into disease-associated isoforms (PrP<sup>Sc</sup>). In this conversion process, a portion of PrP<sup>C</sup> that is mostly  $\alpha$ -helix and random coil becomes refolded into a  $\beta$ -pleated sheet in the PrP<sup>Sc</sup> molecule (1). Although the presence of PrP<sup>C</sup> is essential for prion formation (2), not all cells that express PrP<sup>C</sup> are permissive for PrP<sup>Sc</sup> replication. The reasons why some cells that express PrP<sup>C</sup> do not replicate PrP<sup>Sc</sup> are not fully understood. Reports that the targeting of PrP<sup>C</sup> to specific mem-

branes is required for efficient PrP<sup>Sc</sup> formation (3, 4) indicate that factors that affect the cellular targeting and intracellular trafficking of PrP<sup>C</sup> are critical in determining PrP<sup>Sc</sup> replication.

The majority of PrP<sup>C</sup> molecules are linked to membranes via a glycosylphosphatidylinositol (GPI)<sup>2</sup> anchor (5). GPI anchors play a complex role in the regulation of cell membrane composition, cell signaling and protein trafficking (6). For example, the presence of a GPI anchor targets PrP<sup>C</sup> to specific membrane microdomains called rafts that are required for efficient PrP<sup>Sc</sup> formation (3, 4). The role of the GPI anchor in prion diseases is controversial because transgenic mice that produce anchorless PrP<sup>C</sup> produce large amounts of soluble PrP<sup>Sc</sup> (7), suggesting that the GPI anchor was not essential for PrP<sup>Sc</sup> formation. In contrast, observations that neuronal cells producing anchorless PrP<sup>C</sup> were resistant to scrapie infection (8) and that monoacylated PrP<sup>C</sup> was not converted to PrP<sup>Sc</sup> within neuronal cells (9) indicate that the GPI is a critical factor in facilitating membrane-associated PrP<sup>Sc</sup> formation.

GPI-anchored molecules, including PrP<sup>C</sup>, are rapidly incorporated into living cells (10, 11), and introducing PrP<sup>C</sup> to prion infected cells by cell painting increased PrP<sup>Sc</sup> formation (9). This technique was used to examine the role of specific GPI structures on PrP<sup>C</sup> and PrP<sup>Sc</sup> formation. The GPI anchor attached to PrP<sup>C</sup> is unusual among mammalian GPIs in that it contains sialic acid (12). Neuraminidase digestion was used to produce a PrP<sup>C</sup> with a GPI anchor lacking sialic acid (desialylated PrP<sup>C</sup>), a modification that could not be achieved by genetic manipulation methods (13). We report three major observations: first, that desialylated PrP<sup>C</sup> behaves differently from PrP<sup>C</sup> with regards to its effects on membrane composition and cell signaling; second, that desialylated PrP<sup>C</sup> is not converted to PrP<sup>Sc</sup>; and third, that desialylated PrP<sup>C</sup> inhibits the conversion of endogenous PrP<sup>C</sup> to PrP<sup>Sc</sup>.

## Experimental Procedures

**Cell Lines**—Prion-infected ScN2a and ScGT1 cells were grown in Ham's F-12 medium containing 2 mM glutamine and 2% fetal calf serum. To determine the effect of test preparations on PrP<sup>Sc</sup> formation, ScN2a or ScGT1 cells were added to 6-well plates, allowed to adhere overnight, and then cultured in the presence or absence of test PrP<sup>C</sup> preparations. The cells were grown with daily changes of media ( $\pm$  PrP<sup>C</sup> preparations), and

\* This work was supported by a grant from the European Commission FP6 "Neuprion" Network of Excellence. The authors declare that they have no conflicts of interest with the contents of this article.

<sup>1</sup> To whom correspondence should be addressed: Department of Pathology and Pathogen Biology, Royal Veterinary College, Hawkshead Ln., North Mymms, Hertfordshire AL9 7TA, UK. Tel.: 707-666550; Fax: 707-661464; E-mail: cbate@rvc.ac.uk.

<sup>2</sup> The abbreviations used are: GPI, glycosylphosphatidylinositol; cPLA<sub>2</sub>, cytoplasmic phospholipase A<sub>2</sub>; PLAP, phospholipase A<sub>2</sub>-activating peptide; DRM, detergent-resistant membrane; HPTLC, high performance thin layer chromatography.

the amounts of PrP<sup>Sc</sup> were evaluated after 7 days. In other experiments, ScGT1 and ScN2a cells were pretreated for 1 h with control medium or desialylated PrP<sup>C</sup> and then incubated with phospholipase A<sub>2</sub>-activating peptide (PLAP) (Bachem) for a further 1 h.

**Primary Neurons**—Cortical neurons were prepared from the brains of mouse embryos (day 15.5) derived from Prnp wild type<sup>(+/+)</sup> and Prnp knock-out<sup>(0/0)</sup> mice after mechanical dissociation. Neurons (10<sup>6</sup>) were added to 6-well plates that had been coated with poly-L-lysine and incubated in Ham's F-12 medium containing 5% fetal calf serum for 2 h. Cultures were shaken (600 rpm for 5 min), and nonadherent cells were removed by two washes in PBS. Neurons were grown in neurobasal medium containing B27 components and 5 ng/ml nerve growth factor (Sigma) for 10 days. Immunostaining showed that the cells were greater than 95% neurofilament-positive. In some experiments, neurons were pretreated for 24 h with squalenstatin (GlaxoSmithKline). The fate of PrP<sup>C</sup> preparations was determined by incubating neurons with 10 ng of PrP<sup>C</sup> or desialylated PrP<sup>C</sup>. Cells were washed to remove unbound PrP<sup>C</sup> and incubated in fresh culture medium for different time periods. For infection studies, neurons from either Prnp<sup>(0/0)</sup> or Prnp<sup>(+/+)</sup> mice were pulsed with 1 ng of PrP<sup>Sc</sup> (derived from ScGT1 cells (14)) and then incubated with 10 ng of PrP<sup>C</sup> or 10 ng of desialylated PrP<sup>C</sup>. Media containing PrP<sup>C</sup> preparations were replaced daily for 10 days, and the amount of PrP<sup>Sc</sup> in neurons was measured by ELISA. All experiments were performed in accordance with European regulations (European Community Council Directive, 1986, 56/609/EEC) and approved by the local authority veterinary service/ethical committee.

**Cell Extracts**—Treated cells were washed three times with ice-cold PBS and homogenized in an extraction buffer containing 10 mM Tris-HCl, pH 7.4, 100 mM NaCl, 10 mM EDTA, 0.5% Nonidet P-40, 0.5% sodium deoxycholate, and 0.2% SDS at 10<sup>6</sup> cells/ml, and nuclei and large fragments were removed by centrifugation (1000 × *g* for 5 min). Mixed protease inhibitors (4-(2-aminoethyl) benzenesulfonyl fluoride hydrochloride, aprotinin, leupeptin, bestatin, pepstatin A, and E-46) (Sigma) and a phosphatase inhibitor mixture including PP1, PP2A, microcystin LR, cantharidin, and *p*-bromotetramisole (Sigma) were added to some membrane extracts. To determine the amount of PrP<sup>Sc</sup> in cells extracts were digested with 1 μg/ml proteinase K for 1 h at 37 °C, heated to 95 °C for 5 min, and tested in a PrP ELISA as described (9).

**Isolation of Detergent-resistant Membranes (DRMs)/Rafts**—The cells were homogenized in an ice-cold buffer containing 1% Triton X-100, 10 mM Tris-HCl, pH 7.2, 100 mM NaCl, 10 mM EDTA, and mixed protease inhibitors (10<sup>6</sup> cells/ml), and nuclei and large fragments were removed by centrifugation (1000 × *g* for 5 min). The supernatant was incubated on ice for 60 min prior to further centrifugation (16,000 × *g* for 30 min at 4 °C). The detergent-soluble material was reserved as the normal cell membrane. The insoluble DRMs were homogenized in extraction buffer (10 mM Tris-HCl, pH 7.4, 10 mM NaCl, 10 mM EDTA, 0.5% Nonidet P-40, 0.5% sodium deoxycholate, and 0.2% SDS and mixed protease inhibitors. After further centrif-

ugation (10 min at 16,000 × *g*), the supernatant was collected as the raft fraction.

**Sucrose Density Gradients**—Cells were harvested with a Teflon scraper and homogenized in a buffer containing 250 mM sucrose, 10 mM Tris-HCl, pH 7.2, and 1 mM EDTA, mixed protease inhibitors, 1 mM dithiothreitol. Nuclei and cell fragments were removed by centrifugation (1000 × *g* for 5 min). Membranes were washed by centrifugation at 16,000 × *g* for 20 min at 4 °C and solubilized in an ice-cold buffer containing 1% Triton X-100, 10 mM Tris-HCl, pH 7.2, 150 mM NaCl and 10 mM EDTA. 5, 10, 15, 20, 30, and 40% sucrose solutions were prepared and layered to produce a gradient at 4 °C. Solubilized membranes were layered on top and centrifuged at 50,000 × *g* for 18 h at 4 °C. Serial 0.8- or 0.25-ml volumes were collected from the bottom of gradients.

**Isolation of PrP<sup>C</sup>**—PrP<sup>C</sup> molecules were isolated from murine GT1 neuronal cell membranes using a combination of immunoaffinity columns, size exclusion chromatography (Superdex), and reverse phase chromatography on C18 columns (Waters) as described (9). PrP<sup>C</sup> was digested with 2 units/ml endoglycosidase F (Sigma) and/or 0.2 units/ml neuraminidase (*Clostridium perfringens*; Sigma) for 2 h at 37 °C. TFA was added (final concentration, 0.1%), and digested PrP<sup>C</sup> was loaded onto C18 columns. The samples were eluted under a gradient of acetonitrile and water containing 0.1% TFA. 1-ml samples were collected, split into two, and lyophilized. One sample was solubilized in an extraction buffer (as above) and tested in a PrP ELISA. PrP-containing fractions were solubilized in culture medium by sonication prior to use.

**Test PrP<sup>Sc</sup> Preparations**—PrP<sup>Sc</sup> preparations consisted of supernatants collected from ScGT1 cells that were centrifuged (500 × *g* for 5 min) to remove cell debris and concentrated using a 10-kDa filter (Sartorius Vivaspinn). Preparations were digested with 1 μg/ml proteinase K for 1 h at 37 °C to remove PrP<sup>C</sup>, and the remaining protease-resistant PrP was measured by ELISA (see below). The samples were frozen at −20 °C for storage. On the day of use, PrP<sup>Sc</sup> preparations were diluted to 1 ng/ml in culture medium and sonicated before addition to cells.

**PrP ELISA**—Maxisorb immunoplates (Nunc) were coated with mAb ICSM18 and blocked with 5% milk powder. Samples were applied and detected with biotinylated mAb ICSM35, followed by extravidin-alkaline phosphatase and 1 mg/ml 4-nitrophenyl phosphate (Sigma). Absorbance was measured on a microplate reader at 405 nm, and the amount of PrP in samples was calculated by reference to serial dilutions of recombinant murine PrP (Prionics).

**cPLA<sub>2</sub> ELISA**—Maxisorb immunoplates were coated with 0.5 μg/ml of mouse mAb anti-cPLA<sub>2</sub>, clone CH-7 (Upstate), and blocked. Samples were incubated for 1 h, and the amount of cPLA<sub>2</sub> was detected using a goat polyclonal anti-cPLA<sub>2</sub> (Santa Cruz Biotech) followed by biotinylated anti-goat IgG, extravidin-alkaline phosphatase, and 1 mg/ml 4-nitrophenyl phosphate. Absorbance was measured at 405 nm, and the amount of cPLA<sub>2</sub> protein is expressed in units: 100 units = amount of cPLA<sub>2</sub> in extracts or immunoprecipitates from 10<sup>6</sup> untreated cells.

**Activated cPLA<sub>2</sub> ELISA**—The activation of cPLA<sub>2</sub> is accompanied by phosphorylation of the 505 serine residue, which cre-



## Composition of GPI Affects Prion Formation

ates a unique epitope that was measured in an ELISA using an anti-cPLA<sub>2</sub> mAb (clone CH-7) combined with rabbit polyclonal anti-phospho-cPLA<sub>2</sub> (Cell Signaling Technology) followed by biotinylated anti-rabbit IgG (Dako), extravidin-alkaline phosphatase, and 1 mg/ml 4-nitrophenyl phosphate). Absorbance was measured on a microplate reader at 405 nm. The results were expressed as “units activated cPLA<sub>2</sub>” with 100 units defined as the amount of activated cPLA<sub>2</sub> in extracts derived from 10<sup>6</sup> untreated cells.

**Isolation and Analysis of GPIs**—GPIs were isolated and analyzed as described (13). Briefly, PrP<sup>C</sup> preparations were digested with proteinase K (100 µg/ml) overnight at 37 °C. Digested products were mixed with water-saturated butanol. The butanol phase was collected and washed with water a further three times before being loaded onto C18 columns. GPIs were eluted from C18 columns under a gradient of propanol and water. The presence of phosphatidylinositol in GPIs was determined using mAb (5AB3-11), and specific glycans were detected with biotinylated lectins. Isolated GPIs bound to nitrocellulose membranes by dot blot and were blocked with 5% milk powder. Samples were probed with mAb 5AB3-11, biotinylated *Sambucus nigra* lectin (detects terminal sialic acid residues bound α-2,6 or α-2,3 to galactose), biotinylated concanavalin A (detects mannose), or biotinylated *Ricinus communis* agglutinin I (detects terminal galactose) (Vector Laboratories). Bound lectins were visualized using extravidin peroxidase and enhanced chemiluminescence. The mAb was visualized by incubation with a horseradish peroxidase-conjugated anti-murine IgM and chemiluminescence. In other studies, GPIs were separated on silica gel 60 high performance thin layer chromatography (HPTLC) plates in a mixture of chloroform/methanol/water (10/10/3 v/v/v) and detected by mAb 5AB3-11. The concentrations of GPIs were measured by ELISA. Maxisorb immunoplates were coated with 0.5 µg/ml concanavalin A (binds mannose) and blocked with 5% milk powder in PBS-Tween. Samples were added, and any bound GPI was detected by the addition of mAb 5AB3-11 (15), followed by a biotinylated anti-mouse IgM (Sigma), extravidin-alkaline phosphatase and 1 mg/ml 4-nitrophenyl phosphate. Absorbance was measured on a microplate reader at 405 nm.

**Immunoprecipitations of PrP-specific Rafts**—Treated cells were homogenized in ice cold 1% Triton X-100, 10 mM Tris-HCl, pH 7.2, 100 mM NaCl, 10 mM EDTA at 10<sup>6</sup> cells/ml. Cell debris was removed from by centrifugation (1000 × g for 5 min), and the supernatant was incubated with a mAb to PrP (4F2) for 30 min at 4 °C on rollers. Magnetic microbeads containing protein G (Miltenyi Biotec) were added (10 µl/ml) for 30 min, and protein G-bound antibody complexes were isolated using a µMACS magnetic system (Miltenyi Biotec) at 4 °C. The amounts of sialylated gangliosides in precipitates were determined by diluting precipitates from 10<sup>6</sup> cells in 1 ml of carbonate buffer and plating in Maxisorb immunoplates overnight. The plates were blocked with 5% milk powder in PBS-Tween, and sialylated gangliosides were detected by the addition of biotinylated *S. nigra* lectin followed by extravidin-alkaline phosphatase and 1 mg/ml 4-nitrophenyl phosphate. Absorbance was measured on a microplate reader at 405 nm. Gangliosides within immunoprecipitates were separated by

HPTLC in a solvent containing chloroform/methanol/water (6/4/1 v/v/v). Plates were blocked, and sialic acid containing gangliosides were detected by incubation with biotinylated *S. nigra* lectin, extravidin-peroxidase, and chemiluminescence.

**Western Blot**—Samples were dissolved in 50 µl of Laemmli buffer, boiled, and subjected to electrophoresis on a 15% polyacrylamide gel. Proteins were transferred onto a Hybond-P PVDF membrane (Amersham Biosciences) by semidry blotting. Membranes were blocked using 10% milk powder in PBS containing 0.2% Tween 20. PrP was detected by incubation with mAb ICSM18, β-actin by mAb AC-74 (Sigma), and cPLA<sub>2</sub> with mAb CH-7, followed by a secondary anti-mouse IgG conjugated to peroxidase. Bound antibody was visualized by chemiluminescence.


**Cholesterol and Protein Content**—Cholesterol and protein content were determined in cell membrane extracts. Protein concentrations were measured using a micro-BCA protein assay kit (Pierce). The concentrations of cholesterol were measured using the Amplex Red cholesterol assay kit (Invitrogen). Cholesterol is oxidized by cholesterol oxidase to yield hydrogen peroxide, which reacts with 10-acetyl-3,7-dihydroxyphenoxazine (Amplex Red reagent) to produce highly fluorescent resorufin. Fluorescence was measured by excitation at 550 nm and emission detection at 590 nm. The concentration of cholesterol was calculated by reference to cholesterol standards.

**Statistical Analysis**—Comparison of treatment effects was carried out using Student's paired t-tests, one-way and two-way analysis of variance with Bonferroni's post hoc tests (IBM SPSS statistics 20). Error values are S.D., and significance was determined, where  $p < 0.01$ . Correlations between data sets were analyzed using Pearson's bivariate coefficient (IBM SPSS statistics 20).

## Results

**Deglycosylated PrP<sup>C</sup> Is Converted to PrP<sup>Sc</sup>**—Initial studies were performed to determine whether the N-linked glycans that are attached to PrP<sup>C</sup> affect PrP<sup>Sc</sup> formation in ScGT1 cells. N-Linked glycans were removed from PrP<sup>C</sup> by digestion with endoglycosidase F (16) (Fig. 1A), and deglycosylated PrP<sup>C</sup> was isolated on C18 columns (Fig. 1B). The addition of PrP<sup>C</sup> or endoglycosidase F-digested PrP<sup>C</sup> (deglycosylated PrP<sup>C</sup>) increased the PrP<sup>Sc</sup> content of ScGT1 cells in a dose-dependent manner (Fig. 1C); there were no significant differences in the amounts of PrP<sup>Sc</sup> in cells incubated with PrP<sup>C</sup> or with deglycosylated PrP<sup>C</sup>. Similarly, the addition of 10 ng deglycosylated PrP<sup>C</sup> increased the amount of PrP<sup>Sc</sup> in ScN2a cells (4.7 ng PrP<sup>Sc</sup>/10<sup>6</sup> cells ± 1.1 compared with 1.5 ng PrP<sup>Sc</sup> ± 0.7,  $n = 15$ ,  $p < 0.05$ ). Because the absence of N-linked glycans did not significantly affect PrP<sup>Sc</sup> formation in these cells, all the following experiments were performed with deglycosylated PrP<sup>C</sup> preparations.


**Sialic Acid on the GPI Anchor of PrP<sup>C</sup> Is Necessary for PrP<sup>Sc</sup> Formation**—The GPI anchor attached to PrP<sup>C</sup> is unusual for mammalian GPIs in that it contains sialic acid as first described by Stahl *et al.* (12) and illustrated in Fig. 2A. To confirm the presence of sialic acid in our PrP<sup>C</sup> preparations, GPIs were isolated from PrP<sup>C</sup> and desialylated PrP<sup>C</sup> (neuraminidase-digested) by reverse phase chromatography (Fig. 2B). GPIs



**FIGURE 1. N-Linked glycans did not affect the conversion of PrP<sup>C</sup> to PrP<sup>Sc</sup>.** A, immunoblot showing PrP<sup>C</sup> (lane 1) and deglycosylated PrP<sup>C</sup> (lane 2) separated by PAGE. B, the amounts of deglycosylated PrP<sup>C</sup> (●) in fractions from C18 columns eluted under a gradient of propanol and water. C, the amounts of PrP<sup>Sc</sup> in ScGT1 cells treated daily with concentrations of PrP<sup>C</sup> (□) or deglycosylated PrP<sup>C</sup> (■) as shown for 7 days. The values are means  $\pm$  S.D. from triplicate experiments performed four times ( $n = 12$ ).

derived from PrP<sup>C</sup> and desialylated PrP<sup>C</sup> eluted at different concentrations of propanol. HPTLC analysis confirmed that neuraminidase digestion altered the migration of the GPI isolated from PrP<sup>C</sup> consistent with the loss of sialic acid (Fig. 2C). Isolated GPIs were blotted onto nitrocellulose membranes and probed with mAb 5A3-11 and biotinylated lectins. The digestion of PrP<sup>C</sup> with neuraminidase did not affect the binding of mAb 5A3-11 (binds to phosphatidylinositol) or concanavalin A (binds to mannose) to the GPIs, showing that similar amounts of GPIs were added to blots. *S. nigra* lectin (which reacts with terminal sialic acid bound either  $\alpha$ -2,6 or  $\alpha$ -2,3 to galactose) bound to GPIs isolated from PrP<sup>C</sup>, but not to GPIs derived from neuraminidase-digested PrP<sup>C</sup>, indicating that sialic acid had been removed (Fig. 2D). The lectin *R. communis* agglutinin I bound to neuraminidase-digested GPIs, indicating that terminal galactose had been exposed.


The observation that PrP<sup>Sc</sup> does not replicate in Prnp<sup>(0/0)</sup> neurons (14) allowed us to examine whether desialylated PrP<sup>C</sup> was converted to PrP<sup>Sc</sup>. Both PrP<sup>C</sup> and desialylated PrP<sup>C</sup> bound readily to recipient Prnp<sup>(0/0)</sup> neurons; following the addition of 10-ng PrP<sup>C</sup> preparations for 2 h, there was no significant difference in the amounts of PrP<sup>C</sup> and desialylated PrP<sup>C</sup> bound (9.4 ng of PrP<sup>C</sup>  $\pm$  0.9 compared with 9.7 ng  $\pm$  0.9,  $n = 9$ ,  $p = 0.7$ ). Neurons from Prnp<sup>(0/0)</sup> mice were pulsed with 1 ng of PrP<sup>Sc</sup> for 2 h and incubated with culture media containing control medium, 10 ng of PrP<sup>C</sup>, or 10 ng of desialylated PrP<sup>C</sup>. Media containing the PrP<sup>C</sup>/desialylated PrP<sup>C</sup> was replaced daily for 10 days when the amounts of PrP<sup>Sc</sup> in neurons were measured. We did not detect any PrP<sup>Sc</sup> in neurons that had been pulsed with 1 ng of PrP<sup>Sc</sup> and incubated with control medium, suggesting that original inoculum was degraded by these neurons. However,



**FIGURE 2. Neuraminidase removed the sialic acid from the GPI anchor attached to PrP<sup>C</sup>.** A, cartoon showing the putative structures of the GPI anchor attached to PrP<sup>C</sup> and the desialylated PrP<sup>C</sup>. Glycan residues shown include mannose (Man), sialic acid (SA), galactose (Gal), inositol (Inos), N-acetyl galactosamine (GalNAc), and glucosamine (GlcN). B, GPIs isolated from PrP<sup>C</sup> (○) and desialylated PrP<sup>C</sup> (●) were separated by reverse phase chromatography using a C18 column. GPIs were detected with mAb 5A3-11 reactive with phosphatidylinositol. C, HPTLC showing the migration of GPIs derived from PrP<sup>C</sup> (lane 1) or desialylated PrP<sup>C</sup> (lane 2) separated on silica 60 plates. D, dot blots showing the binding of mAb 5A3-11 (phosphatidylinositol), biotinylated concanavalin A (mannose), biotinylated *S. nigra* lectin (sialic acid), and biotinylated *R. communis* agglutinin I (galactose) to GPI anchors isolated from PrP<sup>C</sup> or desialylated PrP<sup>C</sup>.

PrP<sup>Sc</sup> pulsed neurons incubated daily with 10 ng PrP<sup>C</sup> contained 25.7 ng of PrP<sup>Sc</sup>  $\pm$  4.2,  $n = 12$ , indicating that some of the added PrP<sup>C</sup> was converted to PrP<sup>Sc</sup>. In contrast, no PrP<sup>Sc</sup> was detected in neurons pulsed with 1 ng of PrP<sup>Sc</sup> and incubated daily with 10 ng of desialylated PrP<sup>C</sup>, indicating that it was not converted to PrP<sup>Sc</sup> (Fig. 3A). Immunoblots confirmed the ELISA data and showed that PrP<sup>Sc</sup> was formed in Prnp<sup>(0/0)</sup> neurons pulsed with PrP<sup>Sc</sup> and incubated daily with PrP<sup>C</sup> but not in neurons pulsed with PrP<sup>Sc</sup> and incubated daily with control medium or with desialylated PrP<sup>C</sup> (Fig. 3B).


**Desialylated PrP<sup>C</sup> Reduced PrP<sup>Sc</sup> Formation**—Next, the effects of desialylated PrP<sup>C</sup> on the conversion of endogenous PrP<sup>C</sup> to PrP<sup>Sc</sup> were determined. For this study, neurons derived from Prnp<sup>(+/+)</sup> mice were pulsed with 1 ng of PrP<sup>Sc</sup> for 2 h and incubated for a further 10 days with 10 ng of PrP<sup>C</sup> or 10 ng of desialylated PrP<sup>C</sup>. Neurons pulsed with PrP<sup>Sc</sup> and incubated in control medium contained 40.2 ng of PrP<sup>Sc</sup>  $\pm$  5.2,  $n = 12$ . The addition of PrP<sup>C</sup> increased the amount of PrP<sup>Sc</sup> in neurons to 68.5 ng of PrP<sup>Sc</sup>  $\pm$  7.1,  $n = 12$ , whereas the addition of desialylated PrP<sup>C</sup> reduced the amount of PrP<sup>Sc</sup> to 5.2 ng/ml (Fig. 4, A



**FIGURE 3. Desialylated PrP<sup>C</sup> is not converted to PrP<sup>Sc</sup>.** *A*, the amounts of PrP<sup>Sc</sup> in  $Prnp^{(0/0)}$  neurons pulsed with 1 ng of PrP<sup>Sc</sup> and incubated daily with control medium, 10 ng of PrP<sup>C</sup>, or 10 ng of desialylated PrP<sup>C</sup> for 10 days. The values are means  $\pm$  S.D. from triplicate experiments performed four times ( $n = 12$ ). *B*, immunoblots showing the amounts of  $\beta$ -actin and PrP<sup>Sc</sup> in extracts of  $Prnp^{(0/0)}$  neurons pulsed with 1 ng of PrP<sup>Sc</sup> and incubated daily with control medium (lane i), 10 ng of PrP<sup>C</sup> (lane ii), or 10 ng of desialylated PrP<sup>C</sup> (lane iii) for 10 days.

and *B*), indicating that desialylated PrP<sup>C</sup> suppressed the conversion of PrP<sup>C</sup> to PrP<sup>Sc</sup>. Furthermore, when neurons from  $Prnp^{(0/0)}$  mice were pulsed with 1 ng of PrP<sup>Sc</sup> and incubated with 10 ng of PrP<sup>C</sup> or a mixture of 10 ng of PrP<sup>C</sup> and 10 ng of desialylated PrP<sup>C</sup> for 10 days the presence of desialylated PrP<sup>C</sup> reduced PrP<sup>Sc</sup> production (Fig. 4C). Similarly, the daily addition of desialylated PrP<sup>C</sup> reduced the PrP<sup>Sc</sup> content of recipient ScN2a cells (Fig. 4D) and ScGT1 cells (Fig. 4E).

**Sialic Acid on the GPI Anchor Affects the Fate of PrP<sup>C</sup>**—To determine the mechanism(s) involved in desialylated PrP<sup>C</sup>-induced suppression of PrP<sup>Sc</sup> formation, its properties in noninfected cells were examined. Both PrP<sup>C</sup> and desialylated PrP<sup>C</sup> bound readily to recipient  $Prnp^{(0/0)}$  neurons. The targeting of PrP<sup>C</sup> to DRMs is necessary for efficient prion formation (4), and following the addition of 10-ng PrP<sup>C</sup> preparations, similar amounts of PrP<sup>C</sup> and desialylated PrP<sup>C</sup> were found within DRMs ( $8.2 \text{ ng of PrP}^C \pm 0.9$  compared with  $8.1 \text{ ng of desialylated PrP}^C \pm 1.2$ ,  $n = 9$ ,  $p = 0.7$ ). PrP<sup>C</sup> and desialylated PrP<sup>C</sup> were found within different membrane rafts; after  $Prnp^{(0/0)}$  neurons pulsed with 10 ng of PrP<sup>C</sup> or 10 ng of desialylated PrP<sup>C</sup> were homogenized in ice-cold Triton X-100 (which maintains the integrity of membrane rafts) and subjected to sucrose density gradient analysis; the desialylated PrP<sup>C</sup> was found in lower density rafts than those containing PrP<sup>C</sup> (Fig. 5A). These rafts were examined in detail after immunoprecipitation. Similar amounts of PrP<sup>C</sup> and desialylated PrP<sup>C</sup> were obtained in each precipitate (Fig. 5B). However, the rafts isolated with PrP<sup>C</sup> contained significantly less cholesterol than rafts surrounding desialylated PrP<sup>C</sup> (Fig. 5C). HPTLC analysis of immunoprecipitates demonstrated that rafts containing desialylated PrP<sup>C</sup> con-




**FIGURE 4. Desialylated PrP<sup>C</sup> inhibited PrP<sup>Sc</sup> formation.** *A*, the amounts of PrP<sup>Sc</sup> in  $Prnp^{(+/+)}$  neurons pulsed with 1 ng of PrP<sup>Sc</sup> and incubated daily with control medium ( $\square$ ), 10 ng of PrP<sup>C</sup> or 10 ng of desialylated PrP<sup>C</sup> ( $\blacksquare$ ) for 10 days. The values are means  $\pm$  S.D. from triplicate experiments performed four times ( $n = 12$ ). *B*, immunoblots showing amounts of  $\beta$ -actin and protease-resistant PrP in neurons pulsed with 1 ng of PrP<sup>Sc</sup> and incubated daily with control medium (lane 1), 10 ng of PrP<sup>C</sup> (lane 2), or 10 ng of desialylated PrP<sup>C</sup> (lane 3) for 10 days. *C*, the amounts of PrP<sup>Sc</sup> in  $Prnp^{(0/0)}$  neurons pulsed with 1 ng of PrP<sup>Sc</sup> and incubated daily with control medium, 10 ng of PrP<sup>C</sup>, or a combination of 10 ng of PrP<sup>C</sup> and 10 ng of desialylated PrP<sup>C</sup> for 10 days. The values are means  $\pm$  S.D. from triplicate experiments performed four times ( $n = 12$ ). *D*, the amounts of PrP<sup>Sc</sup> in ScN2a cells treated daily with control medium ( $\square$ ) or desialylated PrP<sup>C</sup> ( $\blacksquare$ ) as shown for 7 days. The values are means  $\pm$  S.D. from triplicate experiments performed four times ( $n = 12$ ). *E*, the amounts of PrP<sup>Sc</sup> in ScGT1 cells treated daily with control medium ( $\square$ ) or desialylated PrP<sup>C</sup> ( $\blacksquare$ ) as shown for 7 days. The values are means  $\pm$  S.D. from triplicate experiments performed four times ( $n = 12$ ).

tained more sialylated gangliosides than did membranes surrounding PrP<sup>C</sup> (Fig. 5D). ELISA studies confirmed that rafts co-precipitated with desialylated PrP<sup>C</sup> contained significantly more sialylated gangliosides than rafts co-precipitated with PrP<sup>C</sup> (Fig. 5E).


Reducing cellular cholesterol concentrations destabilized some rafts, resulting in the relocation of PrP<sup>C</sup> from rafts to detergent-soluble materials (4). Here we examined whether the targeting of desialylated PrP<sup>C</sup> to DRMs was also sensitive to cholesterol depletion.  $Prnp^{(0/0)}$  neurons pretreated with the cholesterol synthesis inhibitor squalenstatin for 24 h were pulsed with 10 ng of PrP<sup>C</sup> or 10 ng of desialylated PrP<sup>C</sup> and DRMs





**FIGURE 5. Sialic acid on the GPI affects the composition of membrane rafts.** Prnp<sup>0/0</sup> neurons were pulsed with 25 ng of PrP<sup>C</sup> or 25 ng of desialylated PrP<sup>C</sup> and homogenized in ice-cold Triton X-100. **A**, the amounts of PrP<sup>C</sup> (□) and desialylated PrP<sup>C</sup> (■) in membranes separated on a sucrose density gradient. The values are means  $\pm$  S.D. from an experiment run in triplicate. Membrane rafts containing PrP<sup>C</sup> were isolated by precipitation with the PrP<sup>C</sup>-reactive mAb 4F2. **B**, blots showing the amounts of PrP<sup>C</sup> and desialylated PrP<sup>C</sup> in immunoprecipitates. **C**, the concentrations of cholesterol in precipitates from PrP<sup>C</sup> (□) or desialylated PrP<sup>C</sup> (■). The values are means  $\pm$  S.D. from four precipitates, each measured in triplicate ( $n = 12$ ). \*, cholesterol significantly higher than in PrP<sup>C</sup> precipitates ( $p < 0.05$ ). **D**, gangliosides in immunoprecipitates from cells incubated with PrP<sup>C</sup> (lane 1) or desialylated PrP<sup>C</sup> (lane 2) separated by HPTLC on silica 60 plates. **E**, the concentrations of gangliosides in precipitates of PrP<sup>C</sup> (□) or desialylated PrP<sup>C</sup> (■). The values are means  $\pm$  S.D. from four precipitates, each measured in triplicate ( $n = 12$ ). \*, gangliosides significantly higher than in PrP<sup>C</sup> precipitates ( $p < 0.05$ ).

(rafts) isolated. Whereas squalestatin reduced the amounts of PrP<sup>C</sup> in rafts, desialylated PrP<sup>C</sup> remained within rafts (Fig. 6A). The concentrations of cholesterol within cell membranes were reduced by squalestatin in a dose-dependent manner. There was a significant correlation between amounts of raft PrP<sup>C</sup>, but not desialylated PrP<sup>C</sup>, and concentrations of cholesterol in squalestatin-treated cells (Fig. 6B). Neurons incubated with PrP<sup>C</sup>/desialylated PrP<sup>C</sup> were also treated with PI-PLC for 1 h. Although digestion with PI-PLC reduced the amounts of cell-associated desialylated PrP<sup>C</sup> to less than 0.5 ng (indicating that all desialylated PrP<sup>C</sup> was expressed at the cell surface), 3.6 ng of



**FIGURE 6. Sialic acid on the GPI anchor affects the stability of PrP<sup>C</sup>.** **A**, the amounts of PrP<sup>C</sup> in rafts (DRMs) from Prnp<sup>0/0</sup> neurons pretreated with squalestatin as shown and pulsed with 10 ng of PrP<sup>C</sup> (□) or desialylated PrP<sup>C</sup> (■) for 2 h. The values are means  $\pm$  S.D. from triplicate experiments performed four times ( $n = 12$ ). \*, significantly less PrP<sup>C</sup> in rafts. **B**, there was a significant correlation between concentrations of cholesterol and concentrations of PrP<sup>C</sup> (○), but not desialylated PrP<sup>C</sup> (●), in rafts derived from Prnp<sup>0/0</sup> neurons pretreated with squalestatin and pulsed with 10 ng of PrP<sup>C</sup>/desialylated PrP<sup>C</sup> for 2 h, Pearson's coefficient = 0.41,  $p < 0.01$ . **C**, the amounts of PrP<sup>C</sup> in Prnp<sup>0/0</sup> neurons pulsed with 10 ng of PrP<sup>C</sup> (□) or desialylated PrP<sup>C</sup> (■) for 2 h and incubated with control medium or PI-PLC for 1 h. The values are means  $\pm$  S.D. from triplicate experiments performed three times ( $n = 9$ ). \*, significantly less desialylated PrP<sup>C</sup> after digestion with PI-PLC. **D**, the amounts of PrP<sup>C</sup> remaining Prnp<sup>0/0</sup> neurons pulsed with 10 ng of PrP<sup>C</sup> (□) or desialylated PrP<sup>C</sup> (■) measured at time points thereafter. The values are means  $\pm$  S.D. from triplicate experiments performed three times ( $n = 9$ ). \*, significantly more desialylated PrP<sup>C</sup> remaining in neurons. **E**, the amounts of PrP<sup>C</sup> in Prnp<sup>0/0</sup> neurons incubated for 2 h with 10 ng of PrP<sup>C</sup> (□) or 10 ng of desialylated PrP<sup>C</sup> (■) and treated with control medium or 5  $\mu$ M glimepiride for 1 h. The values are means  $\pm$  S.D. from triplicate experiments performed three times ( $n = 9$ ). \*, significantly less PrP<sup>C</sup> in neurons after treatment with glimepiride.

PrP<sup>C</sup>  $\pm$  0.4 remained cell-associated after PI-PLC digestion (Fig. 6C). Increasing the concentration of PI-PLC, or incubation period, did not affect the amounts of cell-associated PrP<sup>C</sup>, suggesting that there is a pool of PrP<sup>C</sup> that is not expressed at the surface of neurons. Because proteins with different GPI anchors traffic via different pathways (17, 18), the effect of the GPI anchor on the fate of PrP<sup>C</sup> within cells was investigated.

## Composition of GPI Affects Prion Formation

**TABLE 1**

**Sialic acid in the GPI anchor affected prion-mediated activation of cPLA<sub>2</sub>**

The amounts of activated cPLA<sub>2</sub> in cell extracts from ScN2a or ScGT1 cells pretreated with control medium or with 25 ng of desialylated PrP<sup>C</sup> and then incubated with 1 μM PLAP for 1 h. The values are the means ± S.D. from triplicate experiments performed five times (*n* = 15). The amount of activated cPLA<sub>2</sub> in untreated cells was standardized as 1 for each cell line.


Treatment	Activated cPLA <sub>2</sub>	
	ScN2a cells	ScGT1 cells
	units/10 <sup>6</sup> cells	
None	1.0 ± 0.14	1.0 ± 0.11
Desialylated PrP <sup>C</sup>	0.48 ± 0.12	0.6 ± 0.18
PLAP	3.87 ± 0.58	2.42 ± 0.28
Desialylated PrP <sup>C</sup> + PLAP	3.52 ± 0.49	2.22 ± 0.29

Prnp<sup>(0/0)</sup> neurons were pulsed with 10 ng of PrP<sup>C</sup> preparations, and the amounts of PrP<sup>C</sup> were measured at time points thereafter. Whereas PrP<sup>C</sup> was rapidly cleared from neurons and was absent after 48 h, desialylated PrP<sup>C</sup> remained in neurons for up to 96 h (Fig. 6D). Some GPI-anchored proteins are released from cells following the glimepiride-induced activation of an endogenous GPI-PLC (19, 20). Treatment with 5 μM glimepiride caused the release of PrP<sup>C</sup> but not desialylated PrP<sup>C</sup> (Fig. 6E).

**Desialylated PrP<sup>C</sup> Reduced the Activation of cPLA<sub>2</sub> in Prion-infected Cells**—Because the concentration of cholesterol within cell membranes affects PrP<sup>Sc</sup> formation (4), the effects of PrP<sup>C</sup> and desialylated PrP<sup>C</sup> on cholesterol in prion-infected cells were also examined. The amount of cholesterol in ScGT1 cell membranes was significantly increased after incubation with 10 ng of PrP<sup>C</sup> (744 ng of cholesterol/10<sup>6</sup> cells ± 84 compared with 534 ± 48, *n* = 9, *p* < 0.05) or with 10 ng of desialylated PrP<sup>C</sup> (771 ng of cholesterol/10<sup>6</sup> cells ± 76 compared with 534 ± 48, *n* = 9, *p* < 0.05). Critically there was no significant difference in the amounts of cholesterol between ScGT1 cells treated with PrP<sup>C</sup> or with desialylated PrP<sup>C</sup> (744 ng of cholesterol/10<sup>6</sup> cells ± 84 compared with 771 ± 76, *n* = 9, *p* < 0.36).

Because PrP<sup>Sc</sup> formation is dependent upon the activation of cPLA<sub>2</sub> (21), the effects of desialylated GPI on this enzyme were studied. The addition of 25 ng of desialylated PrP<sup>C</sup> significantly reduced the amounts of activated cPLA<sub>2</sub> in both ScN2a and ScGT1 cells (Table 1). However, desialylated PrP<sup>C</sup> did not affect PLAP-induced activation of cPLA<sub>2</sub>, showing that it did not have a direct effect upon cPLA<sub>2</sub>. The possibility that the addition of PLAP could restore PrP<sup>Sc</sup> formation in cells treated with desialylated PrP<sup>C</sup> was investigated. PrP<sup>Sc</sup> was not detected in ScGT-1 cells treated with a combination of 25 ng of desialylated PrP<sup>C</sup> and 1 μM PLAP (PrP<sup>Sc</sup> < 0.05 ng/10<sup>6</sup> cells), showing that nonselective activation of cPLA<sub>2</sub> was not sufficient to reverse desialylated PrP<sup>C</sup>-induced suppression of PrP<sup>Sc</sup> formation.

The clustering of GPIs in the membrane that arises as a consequence of the self-aggregation of PrP<sup>Sc</sup> molecules can be mimicked by mAb-mediated cross-linkage of PrP<sup>C</sup>, which also activates cPLA<sub>2</sub> (22). When Prnp<sup>(0/0)</sup> neurons were pulsed with PrP<sup>C</sup> or desialylated PrP<sup>C</sup> and incubated with mAb 4F2 cross-linkage of PrP<sup>C</sup>, but not desialylated PrP<sup>C</sup>, it caused a dose-dependent increase in activated cPLA<sub>2</sub> (Fig. 7A). Furthermore, when Prnp<sup>(+/+)</sup> neurons were pulsed with desialylated PrP<sup>C</sup> and incubated with mAb 4F2 to cross-link PrP<sup>C</sup>, the presence of desialylated PrP<sup>C</sup> reduced activated cPLA<sub>2</sub> in a dose-dependent manner (Fig. 7B).



**FIGURE 7. Desialylated PrP<sup>C</sup> reduces activation of cPLA<sub>2</sub> by cross-linkage of PrP<sup>C</sup>.** A, the amounts of activated cPLA<sub>2</sub> in Prnp<sup>(0/0)</sup> neurons pretreated with PrP<sup>C</sup> (□) or desialylated PrP<sup>C</sup> (■) as shown and incubated with mAb 4F2. The values are means ± S.D. from triplicate experiments performed three times (*n* = 9). B, the amounts of activated cPLA<sub>2</sub> in Prnp<sup>(+/+)</sup> neurons pretreated with desialylated PrP<sup>C</sup> as shown and incubated with mAb 4F2. The values are means ± S.D. from triplicate experiments performed three times (*n* = 9).


**Sialic Acid on the GPI Anchor Stabilized cPLA<sub>2</sub> within PrP-containing Rafts**—Upon activation, cPLA<sub>2</sub> migrates from the cytoplasm to rafts (9, 23). Sucrose density gradients showed that in untreated ScGT1 cells, most of the cPLA<sub>2</sub> was in low density membranes (fractions 9–12), whereas in cells that had been incubated with 25 ng of desialylated PrP<sup>C</sup>, a proportion of cPLA<sub>2</sub> had relocated to fractions 4–7 (Fig. 8A). Immunoprecipitation studies showed that cPLA<sub>2</sub> was found within PrP<sup>Sc</sup>-containing rafts (9), raising the possibility that the sialic acid moiety on the GPI attached to PrP<sup>C</sup> or PrP<sup>Sc</sup> is required to capture cPLA<sub>2</sub> within lipid rafts. This hypothesis was tested by treating Prnp<sup>(+/+)</sup> neurons with 10 ng of PrP<sup>C</sup> or 10 ng of desialylated PrP<sup>C</sup> followed by the cross-linkage of PrP<sup>C</sup> by mAb 4F2. Although precipitates contained similar amounts of PrP<sup>C</sup>, the precipitates from neurons incubated with PrP<sup>C</sup> contained more cPLA<sub>2</sub> than precipitates from neurons containing desialylated PrP<sup>C</sup> (Fig. 8B).

## Discussion

Recent studies suggest that differential glycosylation of the GPI anchor can affect the properties of some proteins, including protein structure (24), membrane localization, and intracellular trafficking (25). Here we show that sialic acid in the GPI affected the composition of the membranes surrounding PrP<sup>C</sup>, cell signaling, and the conversion of PrP<sup>C</sup> to PrP<sup>Sc</sup>.

Desialylated PrP<sup>C</sup> was created by the sequential digestion of PrP<sup>C</sup> with an endoglycosidase F (to remove N-linked glycans)






**FIGURE 8. Desialylated PrP<sup>C</sup> reduced the targeting of cPLA<sub>2</sub> to rafts.** *A*, the amounts of cPLA<sub>2</sub> in membrane extracts separated on a sucrose density gradient from ScGT1 cells treated with control medium (□) or 25 ng of desialylated PrP<sup>C</sup> (■). The values are means ± S.D. from an experiment run in triplicate. *B*, immunoblots showing amounts of PrP<sup>C</sup> and cPLA<sub>2</sub> precipitated by mAb 4F2 (reactive with PrP<sup>S</sup>) from wild type Prnp<sup>(+/+)</sup> neurons treated for 3 h with control medium or 25 ng of desialylated PrP<sup>C</sup>.

followed by neuraminidase to remove sialic acid. The removal of *N*-linked glycans was necessary to show that the effects of the subsequent neuraminidase digestion was upon sialic acid contained within the GPI anchor and not due to the removal of sialic acid contained within the *N*-linked glycans. Prnp<sup>(0/0)</sup> neurons exposed to PrP<sup>S</sup> demonstrated that although PrP<sup>C</sup> was converted to PrP<sup>S</sup>, desialylated PrP<sup>C</sup> was not. Consequently, the factors that prevented the conversion of desialylated PrP<sup>C</sup> to PrP<sup>S</sup> were examined. Both PrP<sup>C</sup> and desialylated PrP<sup>C</sup> bound to recipient cells, increased the concentrations of cholesterol in cell membranes, and were targeted to rafts. However, membrane rafts are heterogeneous (26), and the composition of the GPI anchor is thought to target proteins to specific rafts. For example, PrP<sup>C</sup> and Thy-1 have different GPI anchors and are targeted to different rafts (27), and replacing the GPI of PrP<sup>C</sup> with that of Thy-1 altered the trafficking of PrP<sup>C</sup> to apical membranes (28). Therefore, it is possible that the differences in their GPIs direct PrP<sup>C</sup> and desialylated PrP<sup>C</sup> to different rafts and consequently their trafficking within cells.

GPI-anchored proteins are surrounded by specific phospholipids, glycolipids, and cholesterol that constitute a raft, the composition of which is dependent upon multiple interactions between the protein, glycans, and membrane lipids (29). Thus, a change in the composition of the GPI anchor attached to a protein could affect the composition of the surrounding raft. This hypothesis is supported by observations that the composition of GPIs attached to Thy-1 differs from those attached to PrP<sup>C</sup> (12, 30), and the membranes surrounding these molecules have different lipid compositions (31). We hypothesize that the presence of sialic acid in the GPI has a direct effect upon the



**FIGURE 9. PrP<sup>C</sup> with sialic acid may alter the underlying cell membrane.** *A* and *B*, schematic showing the proposed membranes surrounding PrP<sup>C</sup> (*A*) and desialylated PrP<sup>C</sup> (*B*) that include cholesterol, saturated phospholipids, unsaturated phospholipids, and gangliosides. *C*, proposed interactions between PrP<sup>C</sup> and PrP<sup>S</sup> and its effects on the surrounding membrane including the incorporation of cPLA<sub>2</sub> into membranes. *D*, proposed interactions between desialylated PrP<sup>C</sup> and PrP<sup>S</sup> and its effects on the surrounding membrane including gangliosides.

composition of the surrounding membrane rafts as illustrated in Fig. 9 (*A* and *B*). Our observation that the rafts surrounding desialylated PrP<sup>C</sup> contained more cholesterol and sialylated gangliosides than those surrounding PrP<sup>C</sup> supported this hypothesis. In addition, treatment with glimepiride caused the release of PrP<sup>C</sup>, but not desialylated PrP<sup>C</sup>, suggesting that only PrP<sup>C</sup> is associated with a raft containing endogenous GPI-PLC. How sialic acid affects the composition of rafts is unknown. One possibility is that the sialic acid contained within the GPI competes with gangliosides for sialic acid-binding proteins within rafts. Thus, the removal of sialic acid from the GPI would allow the incorporation of gangliosides, which are associated with the suppression of PrP<sup>S</sup> formation (32), into rafts. Gangliosides help sequester cholesterol and stabilize rafts (33, 34) especially in brain tissue (35). The increased amounts of gangliosides that surround desialylated PrP<sup>C</sup> would be expected to form a raft containing more cholesterol, as was demonstrated in these experiments. The concentrations of cholesterol in cell membranes affects membrane rigidity and helps stabilize membrane rafts (36). We noted that desialylated PrP<sup>C</sup> remained within rafts in cholesterol-depleted cells, whereas PrP<sup>C</sup> was

relocated to detergent-soluble materials, a finding indicating that the rafts surrounding desialylated PrP<sup>C</sup> are more cholesterol-dense and hence more resistant to cholesterol depletion than those surrounding PrP<sup>C</sup>. Such observations are compatible with reports that the amount of gangliosides in rafts affects the expression and function of some GPI-anchored proteins (37, 38) including PrP<sup>C</sup> (39).

Our observation that desialylated PrP<sup>C</sup> remained in cells longer than PrP<sup>C</sup> suggests that sialic acid in the GPI anchor affects the trafficking of PrP<sup>C</sup>. The endocytosis of PrP<sup>C</sup> involves a step in which it moves out of rafts and enters clathrin-coated pits (17, 40). We speculate that desialylated PrP<sup>C</sup> stabilized within cholesterol-dense rafts does not move out of rafts nor undergo the same endocytic process as PrP<sup>C</sup>. It is possible that the cellular location of desialylated PrP<sup>C</sup> results in reduced interactions with PrP<sup>Sc</sup>, which would explain why desialylated PrP<sup>C</sup> was not converted to PrP<sup>Sc</sup>.

Desialylated PrP<sup>C</sup> also inhibited the conversion of endogenous PrP<sup>C</sup> to PrP<sup>Sc</sup> in primary cortical neurons and ScN2a and ScGT1 cells. Previously we had shown that monoacylated PrP<sup>C</sup> reduced prion formation (9), and others have demonstrated that co-expression of mutant prion proteins alters the cellular localization of wild type PrP<sup>C</sup> and partitioning into DRMs/rafts (41). One possibility is that desialylated PrP<sup>C</sup> competes with endogenous PrP<sup>C</sup> for specific partner proteins involved in endocytosis and that the depletion of these partner proteins alters the trafficking of endogenous PrP<sup>C</sup> and hence PrP<sup>Sc</sup> formation. Here we explored the concept that the binding of desialylated PrP<sup>C</sup> to PrP<sup>Sc</sup> modifies the rafts that are involved in PrP<sup>Sc</sup> formation (3). The composition and hence the function of rafts is dynamic and controlled by an "induced fit" model (26). Because the composition of membranes is affected by the glycan structure of GPIs (29, 31), the membrane surrounding a complex between PrP<sup>Sc</sup> and PrP<sup>C</sup> (Fig. 9C) would be expected to differ from membranes surrounding PrP<sup>Sc</sup> and desialylated PrP<sup>C</sup> (Fig. 9D). Studies in T and B cell receptor signaling show that the coalescence of raft outer membrane proteins affects the composition of the cytoplasmic leaflet and its association with signaling molecules (42–45). We propose that the binding of desialylated PrP<sup>C</sup> to PrP<sup>Sc</sup> changes the composition of the local membrane so that it is unfavorable for the conversion of PrP<sup>C</sup> to PrP<sup>Sc</sup>.

The hypothesis that the clustering of sialic acid-containing GPIs attached to PrP leads to the activation of cPLA<sub>2</sub>, a factor that promotes PrP<sup>Sc</sup> formation (21), was explored. This occurs naturally as a consequence of PrP<sup>Sc</sup> self-aggregation and experimentally following cross-linkage of PrP<sup>C</sup> by mAbs. Whereas cross-linkage of PrP<sup>C</sup> activated cPLA<sub>2</sub>, cross-linkage of desialylated PrP<sup>C</sup> had no effect, demonstrating that sialic acid was required (13). Cross-linkage of a mixture of PrP<sup>C</sup> and desialylated PrP<sup>C</sup> reduced activation of cPLA<sub>2</sub> when compared with cross-linkage of PrP<sup>C</sup> alone (13), indicating that homogeneity of the GPI anchors was critical. Notably the presence of desialylated PrP<sup>C</sup> did not affect the activation of cPLA<sub>2</sub> by PLAP, indicating that it did not have a direct effect upon this enzyme.

To understand how desialylated PrP<sup>C</sup> inhibited activation of cPLA<sub>2</sub>, the cellular location of cPLA<sub>2</sub> was examined. The activation of cPLA<sub>2</sub> causes it to migrate from the cytoplasm to

membrane rafts (23). In control ScGT1 cells, activated cPLA<sub>2</sub> was found predominantly in rafts, and more specifically cPLA<sub>2</sub> co-localized to PrP<sup>Sc</sup>-containing rafts (9). The addition of desialylated PrP<sup>C</sup> caused the dissociation of cPLA<sub>2</sub> from rafts. The targeting of cPLA<sub>2</sub> to membranes containing their substrates can regulate the formation of second messengers such as platelet-activating factor that facilitate PrP<sup>Sc</sup> formation (21). We propose that the density of sialic acid attached to GPIs is critical to the stabilization and activation of cPLA<sub>2</sub> in membrane rafts and that the binding of desialylated PrP<sup>C</sup> to PrP<sup>Sc</sup> changed the composition of the underlying membrane so that it no longer captured and activated cPLA<sub>2</sub>. This reduced the activation of cPLA<sub>2</sub> by existing PrP<sup>Sc</sup> and hindered the conversion of PrP<sup>C</sup> to PrP<sup>Sc</sup>. It is noteworthy that desialylated PrP<sup>C</sup> is surrounded by more gangliosides than PrP<sup>C</sup>, which is consistent with reports that gangliosides inhibit the activation of cPLA<sub>2</sub> (46–48).

Some of the PrP<sup>C</sup> extracted from hamster brains have GPIs that did not contain sialic acid (12). The chemical composition of the GPI anchor is cell type-specific (49), and the GPI anchor attached to PrP<sup>C</sup> derived from a glial cell line (N9 cells) does not contain sialic acid (13). More recently we found that some neuronal cell lines produce predominantly desialylated PrP<sup>C</sup>, suggesting that these cells may be resistant to prion infection, an observation that may explain the different sensitivities of neurons to prion infection.

In conclusion, we show that sialic acid attached to the GPI affects the properties of PrP<sup>C</sup>, altering the surrounding cell membrane, trafficking of PrP<sup>C</sup>, and PrP<sup>C</sup>-induced cell signaling. Critically the presence of desialylated PrP<sup>C</sup> reduced the activation of cPLA<sub>2</sub> and PrP<sup>Sc</sup> formation in prion-infected neuronal cells. We propose that sialic acid on the GPI anchor attached to PrP<sup>C</sup> affects the membrane targeting and cell signaling that is conducive to its conversion to PrP<sup>Sc</sup>. Consequently, therapeutics that block the incorporation of sialic acid into GPI anchors could prove useful in the treatment of prion diseases.

**Author Contributions**—C. B. was responsible for conception and design, data collection and analysis, and manuscript writing and revision; W. N. was responsible for experimental design, and data collection and analysis; and A. W. was responsible for conception and design and manuscript writing. All authors read and approved the final manuscript.

**Acknowledgment**—We thank Dr. Mourad Tayebi for supplying mAbs ICSM18 and ICSM35.

## References

1. Pan, K. M., Baldwin, M., Nguyen, J., Gasset, M., Serban, A., Groth, D., Mehlhorn, I., Huang, Z., Fletterick, R. J., and Cohen, F. E. (1993) Conversion of  $\alpha$ -helices into  $\beta$ -sheets features in the formation of the scrapie prion proteins. *Proc. Natl. Acad. Sci. U.S.A.* **90**, 10962–10966
2. Büeler, H., Aguzzi, A., Sailer, A., Greiner, R. A., Autenried, P., Aguet, M., and Weissmann, C. (1993) Mice devoid of PrP are resistant to scrapie. *Cell* **73**, 1339–1347
3. Taylor, D. R., and Hooper, N. M. (2006) The prion protein and lipid rafts. *Mol. Membr. Biol.* **23**, 89–99
4. Taraboulos, A., Scott, M., Semenov, A., Avrahami, D., Laszlo, L., Prusiner,

- S. B., and Avraham, D. (1995) Cholesterol depletion and modification of COOH-terminal targeting sequence of the prion protein inhibit formation of the scrapie isoform. *J. Cell Biol.* **129**, 121–132
5. Stahl, N., Borchelt, D. R., Hsiao, K., and Prusiner, S. B. (1987) Scrapie prion protein contains a phosphatidylinositol glycolipid. *Cell* **51**, 229–240
  6. Chatterjee, S., and Mayor, S. (2001) The GPI-anchor and protein sorting. *Cell Mol. Life Sci.* **58**, 1969–1987
  7. Chesebro, B., Trifilo, M., Race, R., Meade-White, K., Teng, C., LaCasse, R., Raymond, L., Favara, C., Baron, G., Priola, S., Caughey, B., Masliah, E., and Oldstone, M. (2005) Anchorless prion protein results in infectious amyloid disease without clinical scrapie. *Science* **308**, 1435–1439
  8. McNally, K. L., Ward, A. E., and Priola, S. A. (2009) Cells expressing anchorless prion protein are resistant to scrapie infection. *J. Virol.* **83**, 4469–4475
  9. Bate, C., and Williams, A. (2011) Monoacylated cellular prion protein modifies cell membranes, inhibits cell signaling and reduces prion formation. *J. Biol. Chem.* **286**, 8752–8758
  10. Legler, D. F., Doucey, M. A., Schneider, P., Chapatte, L., Bender, F. C., and Bron, C. (2005) Differential insertion of GPI-anchored GFPs into lipid rafts of live cells. *FASEB J.* **19**, 73–75
  11. Liu, T., Li, R., Pan, T., Liu, D., Petersen, R. B., Wong, B. S., Gambetti, P., and Sy, M. S. (2002) Intercellular transfer of the cellular prion protein. *J. Biol. Chem.* **277**, 47671–47678
  12. Stahl, N., Baldwin, M. A., Hecker, R., Pan, K. M., Burlingame, A. L., and Prusiner, S. B. (1992) Glycosylphospholipid anchors of the scrapie and cellular prion proteins contain sialic acid. *Biochemistry* **31**, 5043–5053
  13. Bate, C., and Williams, A. (2012) Neurodegeneration induced by the clustering of sialylated glycosylphosphatidylinositols of prion proteins. *J. Biol. Chem.* **287**, 7935–7944
  14. Bate, C., Tayebi, M., and Williams, A. (2010) The glycosylphosphatidylinositol anchor is a major determinant of prion binding and replication. *Biochem. J.* **428**, 95–101
  15. Bate, C. A., and Kwiatkowski, D. (1994) A monoclonal antibody that recognizes phosphatidylinositol inhibits induction of tumor necrosis factor alpha by different strains of *Plasmodium falciparum*. *Infect. Immun.* **62**, 5261–5266
  16. Haraguchi, T., Fisher, S., Olofsson, S., Endo, T., Groth, D., Tarentino, A., Borchelt, D. R., Teplow, D., Hood, L., and Burlingame, A. (1989) Asparagine-linked glycosylation of the scrapie and cellular prion proteins. *Arch. Biochem. Biophys.* **274**, 1–13
  17. Sunyach, C., Jen, A., Deng, J., Fitzgerald, K. T., Frobert, Y., Grassi, J., McCaffrey, M. W., and Morris, R. (2003) The mechanism of internalization of glycosylphosphatidylinositol-anchored prion protein. *EMBO J.* **22**, 3591–3601
  18. Wilson, B. S., Steinberg, S. L., Liederman, K., Pfeiffer, J. R., Surviladze, Z., Zhang, J., Samelson, L. E., Yang, L. H., Kotula, P. G., and Oliver, J. M. (2004) Markers for detergent-resistant lipid rafts occupy distinct and dynamic domains in native membranes. *Mol. Biol. Cell* **15**, 2580–2592
  19. Bate, C., Tayebi, M., Diomedea, L., Salmons, M., and Williams, A. (2009) Glimepiride reduces the expression of PrP<sup>C</sup>, prevents PrP<sup>Sc</sup> formation, and protects against prion mediated neurotoxicity. *PLoS One* **4**, e8221
  20. Müller, G., Dearey, E. A., and Pünter, J. (1993) The sulphonylurea drug, glimepiride, stimulates release of glycosylphosphatidylinositol-anchored plasma-membrane proteins from 3T3 adipocytes. *Biochem. J.* **289**, 509–521
  21. Bate, C., Reid, S., and Williams, A. (2004) Phospholipase A<sub>2</sub> inhibitors or platelet-activating factor antagonists prevent prion replication. *J. Biol. Chem.* **279**, 36405–36411
  22. Bate, C., and Williams, A. (2011) Amyloid-β-induced synapse damage is mediated via cross-linkage of the cellular prion protein. *J. Biol. Chem.* **286**, 37955–37963
  23. Nalefski, E. A., Sultzman, L. A., Martin, D. M., Kriz, R. W., Towler, P. S., Knopf, J. L., and Clark, J. D. (1994) Delineation of two functionally distinct domains of cytosolic phospholipase A<sub>2</sub>, a regulatory Ca<sup>2+</sup>-dependent lipid-binding domain and a Ca<sup>2+</sup>-independent catalytic domain. *J. Biol. Chem.* **269**, 18239–18249
  24. Barboni, E., Rivero, B. P., George, A. J., Martin, S. R., Renoup, D. V., Hounsell, E. F., Barber, P. C., and Morris, R. J. (1995) The glycosylphosphatidylinositol anchor affects the conformation of Thy-1 protein. *J. Cell Sci.* **108**, 487–497
  25. Nicholson, T. B., and Stanners, C. P. (2006) Specific inhibition of GPI-anchored protein function by homing and self-association of specific GPI anchors. *J. Cell Biol.* **175**, 647–659
  26. Pike, L. J. (2004) Lipid rafts: heterogeneity on the high seas. *Biochem. J.* **378**, 281–292
  27. Madore, N., Smith, K. L., Graham, C. H., Jen, A., Brady, K., Hall, S., and Morris, R. (1999) Functionally different GPI proteins are organized in different domains on the neuronal surface. *EMBO J.* **18**, 6917–6926
  28. Puig, B., Altmeyer, H. C., Thurm, D., Geissen, M., Conrad, C., Bräulke, T., and Glatzel, M. (2011) N-Glycans and glycosylphosphatidylinositol-anchors act on polarized sorting of mouse PrP(C) in Madin-Darby canine kidney cells. *PLoS One* **6**, e24624
  29. Anderson, R. G., and Jacobson, K. (2002) A role for lipid shells in targeting proteins to caveolae, rafts, and other lipid domains. *Science* **296**, 1821–1825
  30. Homans, S. W., Ferguson, M. A., Dwek, R. A., Rademacher, T. W., Anand, R., and Williams, A. F. (1988) Complete structure of the glycosyl phosphatidylinositol membrane anchor of rat brain Thy-1 glycoprotein. *Nature* **19**, 269–272
  31. Brügger, B., Graham, C., Leibrecht, I., Mombelli, E., Jen, A., Wieland, F., and Morris, R. (2004) The membrane domains occupied by glycosylphosphatidylinositol-anchored prion protein and Thy-1 differ in lipid composition. *J. Biol. Chem.* **279**, 7530–7536
  32. Naslavsky, N., Shmeeda, H., Friedlander, G., Yanai, A., Futerman, A. H., Barenholz, Y., and Taraboulos, A. (1999) Sphingolipid depletion increases formation of the scrapie prion protein in neuroblastoma cells infected with prions. *J. Biol. Chem.* **274**, 20763–20771
  33. Cantù, L., Del Favero, E., Sonnino, S., and Prinetti, A. (2011) Gangliosides and the multiscale modulation of membrane structure. *Chem. Phys. Lipids* **164**, 796–810
  34. Slotte, J. P. (1999) Sphingomyelin-cholesterol interactions in biological and model membranes. *Chem. Phys. Lipids* **102**, 13–27
  35. Ohmi, Y., Ohkawa, Y., Yamauchi, Y., Tajima, O., Furukawa, K., and Furukawa, K. (2012) Essential roles of gangliosides in the formation and maintenance of membrane microdomains in brain tissues. *Neurochem. Res.* **37**, 1185–1191
  36. Brown, D. A., and London, E. (2000) Structure and function of sphingolipid- and cholesterol-rich membrane rafts. *J. Biol. Chem.* **275**, 17221–17224
  37. Simons, M., Friedrichson, T., Schulz, J. B., Pitto, M., Masserini, M., and Kurzchalia, T. V. (1999) Exogenous administration of gangliosides displaces GPI-anchored proteins from lipid microdomains in living cells. *Mol. Biol. Cell* **10**, 3187–3196
  38. Nagafuku, M., Kabayama, K., Oka, D., Kato, A., Tani-ichi, S., Shimada, Y., Ohno-Iwashita, Y., Yamasaki, S., Saito, T., Iwabuchi, K., Hamaoka, T., Inokuchi, J., and Kosugi, A. (2003) Reduction of glycosphingolipid levels in lipid rafts affects the expression state and function of glycosylphosphatidylinositol-anchored proteins but does not impair signal transduction via the T cell receptor. *J. Biol. Chem.* **278**, 51920–51927
  39. Galvan, C., Camoletto, P. G., Dotti, C. G., Aguzzi, A., and Ledesma, M. D. (2005) Proper axonal distribution of PrP<sup>C</sup> depends on cholesterol-sphingomyelin-enriched membrane domains and is developmentally regulated in hippocampal neurons. *Mol. Cell Neurosci.* **30**, 304–315
  40. Taylor, D. R., Watt, N. T., Perera, W. S., and Hooper, N. M. (2005) Assigning functions to distinct regions of the N-terminus of the prion protein that are involved in its copper-stimulated, clathrin-dependent endocytosis. *J. Cell Sci.* **118**, 5141–5153
  41. Schiff, E., Campana, V., Tivodar, S., Lebreton, S., Gousset, K., and Zurzolo, C. (2008) Coexpression of wild-type and mutant prion proteins alters their cellular localization and partitioning into detergent-resistant membranes. *Traffic* **9**, 1101–1115
  42. Montixi, C., Langlet, C., Bernard, A.-M., Thimonier, J., Dubois, C., Wurzel, M.-A., Chauvin, J.-P., Pierres, M., and He, H.-T. (1998) Engagement of T cell receptor triggers its recruitment to low-density detergent-insoluble membrane domains. *EMBO J.* **17**, 5334–5348



## Composition of GPI Affects Prion Formation

43. Eisenberg, S., Shvartsman, D. E., Ehrlich, M., and Henis, Y. I. (2006) Clustering of raft-associated proteins in the external membrane leaflet modulates internal leaflet H-Ras diffusion and signaling. *Mol. Cell Biol.* **26**, 7190–7200
44. Gri, G., Molon, B., Manes, S., Pozzan, T., and Viola, A. (2004) The inner side of T cell lipid rafts. *Immunol. Lett.* **94**, 247–252
45. Field, K. A., Holowka, D., and Baird, B. (1995) Fc epsilon RI-mediated recruitment of p53/56lyn to detergent-resistant membrane domains accompanies cellular signaling. *Proc. Natl. Acad. Sci. U.S.A.* **92**, 9201–9205
46. Bianco, I. D., Fidelio, G. D., and Maggio, B. (1989) Modulation of phospholipase A2 activity by neutral and anionic glycosphingolipids in monolayers. *Biochem. J.* **258**, 95–99
47. Maggio, B., Bianco, I. D., Montich, G. G., Fidelio, G. D., and Yu, R. K. (1994) Regulation by gangliosides and sulfatides of phospholipase A2 activity against dipalmitoyl- and dilauroylphosphatidylcholine in small unilamellar bilayer vesicles and mixed monolayers. *Biochim. Biophys. Acta* **1190**, 137–148
48. Yang, H. C., Farooqui, A. A., and Horrocks, L. A. (1994) Effects of glycosaminoglycans and glycosphingolipids on cytosolic phospholipases A2 from bovine brain. *Biochem. J.* **299**, 91–95
49. McConville, M. J., and Ferguson, M. A. (1993) The structure, biosynthesis and function of glycosylated phosphatidylinositols in the parasitic protozoa and higher eukaryotes. *Biochem. J.* **294**, 305–324

# **Sialic Acid on the Glycosylphosphatidylinositol Anchor Regulates PrP-mediated Cell Signaling and Prion Formation**

Clive Bate, William Nolan and Alun Williams

*J. Biol. Chem.* 2016, 291:160-170.

doi: 10.1074/jbc.M115.672394 originally published online November 9, 2015

---

Access the most updated version of this article at doi: [10.1074/jbc.M115.672394](https://doi.org/10.1074/jbc.M115.672394)

## Alerts:

- [When this article is cited](#)
- [When a correction for this article is posted](#)

[Click here](#) to choose from all of JBC's e-mail alerts

This article cites 49 references, 33 of which can be accessed free at <http://www.jbc.org/content/291/1/160.full.html#ref-list-1>