

This is the peer-reviewed, manuscript version of the following article:

Orriss, I. R. (2015) 'The role of purinergic signalling in the musculoskeletal system', *Autonomic Neuroscience*, 191, 124-134.

The final version is available online via <http://dx.doi.org/10.1016/j.autneu.2015.04.013>.

© 2015. This manuscript version is made available under the CC-BY-NC-ND 4.0 license <http://creativecommons.org/licenses/by-nc-nd/4.0/>.

The full details of the published version of the article are as follows:

TITLE: The role of purinergic signalling in the musculoskeletal system

AUTHORS: Orriss, I. R.

JOURNAL TITLE: *Autonomic Neuroscience*

VOLUME/EDITION: 191

PUBLISHER: Elsevier

PUBLICATION DATE: September 2015 (online)

DOI: 10.1016/j.autneu.2015.04.013

The role of purinergic signalling in the musculoskeletal system

Isabel R Orriss

Department of Comparative Biomedical Sciences
Royal Veterinary College, London, UK

Address correspondence to:

Isabel R Orriss
Department of Comparative Biomedical Sciences,
Royal Veterinary College
Royal College Street,
London, NW1 0TU
Tel: 020 7468 1238 (ext 5468)
Email: iorriss@rvc.ac.uk

Grant support: **Arthritis Research UK**

Key words: ATP, P2 receptors, bone, mineralisation, cartilage, muscle

Abstract

Accumulating evidence now suggests that purinergic signalling exerts significant regulatory effects in the musculoskeletal system. In particular, it has emerged that extracellular nucleotides are key regulators of bone cell differentiation, survival and function. This review discusses our current understanding of the effects of purinergic signalling in bone, cartilage and muscle.

1. Introduction

Adenosine triphosphate (ATP) has long been recognized for its role in intracellular energy metabolism; however, it is also an important extracellular signalling molecule. The potent actions of ATP were first described in 1929, yet it was 1972 before the concept of purinergic neurotransmission was proposed (Burnstock, 1972). Extracellular nucleotides, signalling via purinergic receptors, are now known to participate in numerous biological processes in both neuronal and non-neuronal tissues. The receptors for purines and pyrimidines are classified into two groups; P1 receptors and P2 receptors. There are four P1 receptor subtypes (A_1, A_{2a}, A_{2b}, A_3); these receptors are G-protein coupled and activated by adenosine. P2 receptors respond to a number of different nucleotides including ATP, adenosine diphosphate (ADP), uridine triphosphate (UTP) and uridine diphosphate (UDP). The P2 receptors are further subdivided into the P2X ligand-gated ion channels and P2Y G-protein-coupled receptors (Abbracchio et al., 1994; Burnstock et al., 1985; Ralevic et al., 1998). To date, seven P2X receptors (P2X1-7) and eight P2Y receptors (P2Y_{1,2,4,6,11,12,13,14}) have been identified; each receptor has been cloned, characterised and displays distinct tissue expression and pharmacology (Burnstock, 2007).

Over the last two decades, it has emerged that purinergic signalling plays a key regulatory role in the musculoskeletal system. In particular, the regulation of bone cell function by extracellular nucleotides has emerged as a particularly active and promising area of research. This review will summarise current understanding into the role of purinergic signalling in bone, cartilage and muscle.

2. Bone

Bone is a composite tissue containing inorganic mineral salts deposited within an organic collagenous matrix, and three cell types: osteoblasts, osteoclasts and osteocytes. Osteoblasts, which are derived from mesenchymal stem cells, are the bone-forming cells. When osteoblasts become incorporated within the bone matrix they may undergo a terminal differentiation to form an osteocyte. Osteocytes, the most abundant cells in bone (Jande et al., 1973), form a regular interconnected network of cells that is thought to regulate bone remodelling and mediate the responses to mechanical loading (Bonewald, 2011). Osteoclasts, the bone-resorbing cells, are usually multinuclear, and are formed by the fusion of mononuclear progenitors of the monocyte/macrophage lineage. The coordinated activity of these cells allows the skeleton to grow, adapt and repair itself; abnormalities in this process result in a variety of skeletal disorders.

The first indication that purinergic signalling could be important in modulating bone cell function came over 20 years ago when P1 (A_2 -like) adenosine receptors were reported on osteoblasts (Lerner 1987). The expression of multiple P1 and P2 receptor subtypes in bone cells has now been widely reported (see review by (Burnstock et al., 2013)). More recently, analysis of knockout mouse models has provided evidence for the important role of purinergic signalling in bone remodelling *in vivo* (Gartland et al., 2003b; Ke et al., 2003; Orriss et al., 2011a; Su et al., 2012; Wang et al., 2012) (**Fig. 1**)

2.1. The role of purinergic signalling in osteoblast biology

P2 receptor expression by osteoblasts was first reported in the late 1980's when fluorescence studies demonstrated that extracellular nucleotides could transiently increase $[Ca^{2+}]_i$ and induce IP_3 formation (Kumagai et al., 1991; Kumagai et al., 1989). Subsequent pharmacological studies revealed that extracellular nucleotides interacted with at least two P2 receptors (P2Y₁ and P2Y₂) on rat osteoblast-like cells (Reimer et al., 1992; Yu et al., 1993b). The first molecular evidence for P2Y receptor expression was shown in 1995 by Bowler *et al*, who used RT-PCR and *in situ* hybridisation to demonstrate P2Y₂ receptor expression in human osteoblasts (Bowler et al., 1995). A detailed investigation of both single

cells and populations of human osteoblasts indicated that there was a heterogeneity in P2 receptor expression in any one culture (Dixon et al., 1997). This observation raised the possibility that the differentiation status of the osteoblast influences P2 receptor expression, an idea which was later confirmed in several separate studies (Nishii et al., 2009; Orriss et al., 2012; Orriss et al., 2006).

The expression of multiple P2 receptors by osteoblasts has now been widely reported (**Fig. 2**). Studies using primary rat and mouse calvarial osteoblasts, human osteoblast-like cell lines (MG-63, OHS-4, SaM-1, SaOS-2, Te85) and rat osteosarcoma cell lines (UMR106-01, ROS17/2,8) have all described expression of most or all P2X and P2Y receptor subtypes in osteoblasts (see reviews (Burnstock et al., 2013; Gartland et al., 2012a)). P2 receptor mediated signalling has now been shown to exert many effects on osteoblast proliferation, differentiation, function, gene expression and cell signalling.

Extracellular nucleotides and the regulation of bone mineralisation

Initial functional studies showed that ATP and UTP ($\geq 1\mu\text{M}$), acting via the P2Y₂ receptor, strongly inhibited bone formation by cultured rat osteoblasts (Hoeberz et al., 2002). A follow up investigation demonstrated that ATP and UTP selectively inhibited the mineralisation of the organic matrix and alkaline phosphatase (TNAP) expression and activity (Orriss et al., 2007). It also provided further evidence for the involvement of the P2Y₂ receptor (Orriss et al., 2007). Subsequent skeletal analysis of P2Y₂ receptor knockout mice by dual energy x-ray absorptiometry (DEXA) and micro computed tomography (μCT) demonstrated large increases in trabecular and cortical bone parameters in the long bones (Orriss et al., 2011a; Orriss et al., 2007) (**Fig. 1**). Recently, it was also demonstrated that the P2X₁, P2X₃ and P2X₇ receptor agonists, α,β -meATP, β,γ -methylene ATP and 2'(3')-O-(4-benzoylbenzoyl) adenosine 5'-triphosphate (Bz-ATP) reduced bone mineralization *in vitro* (Orriss et al., 2012). This study used selective P2 receptor antagonists to suggest that the effects were mediated via the P2X₁ and P2X₇ receptor subtypes (Orriss et al., 2012). Furthermore, another investigation demonstrated increased bone mineralisation in osteoblast cultures treated with apyrase, suggesting that endogenous ATP released by osteoblasts acts as an important local brake on mineralisation (Orriss et al., 2013). Combined these data suggest that ATP acts as a key endogenous inhibitor of bone mineralisation.

The ATP concentration in cell cytosol is between 2-5mM. Following membrane damage or necrosis, all cells can release ATP into the extracellular environment, which can then act in an autocrine/paracrine manner to influence local purinergic signalling. Controlled ATP release has been demonstrated from numerous excitatory and non-excitatory cells. In the bone microenvironment, osteoblasts (Buckley et al., 2003; Genetos et al., 2005; Orriss et al., 2009; Romanello et al., 2001; Rumney et al., 2012), osteoclasts (Brandao-Burch et al., 2012) and MLO-Y4 osteocyte-like cells (Genetos et al., 2007; Kringelbach et al., 2014) have all been shown to constitutively release ATP.

Once released, nucleotides are rapidly broken down by an extracellular hydrolysis cascade. Molecular and functional characterisation has shown there are four families of ecto-nucleotidases: (1) the NTPdases (ecto-nucleoside triphosphate diphosphohydrolase); (2) the NPPs (ecto-nucleotide pyrophosphatase/phosphodiesterase); (3) alkaline phosphatases and, (4) ecto-5'-nucleotidase (Zimmermann et al., 2012). Many ecto-nucleotidases have overlapping specificities. For example, NTPdases catalyse the reactions: nucleotide triphosphate (NTP) \rightarrow nucleotide diphosphate (NDP) + phosphate (Pi) and NDP \rightarrow nucleotide monophosphate (NMP) + phosphate (Pi), whereas NPPs hydrolyse NTP \rightarrow NMP + pyrophosphate (PP_i) or NDP \rightarrow NMP + Pi. Thus, the combined activities of these ecto-enzymes limit the actions of extracellular nucleotides to cells within close proximity of the release site. Osteoblasts express three members of the NPP family (NPP1-3) (Hessle et al., 2002;

Johnson et al., 2000; Orriss et al., 2007) and at least six members of the NTPdase family (NTPdase 1-6)(Orriss et al., 2010). This hydrolysis of ATP and other NTPs by NPPs is particularly important in bone because the product, pyrophosphate (PP_i), is the key, local physicochemical inhibitor of mineralisation (Fleisch et al., 1962; Fleisch et al., 1961). Of the NPPs expressed in bone, NPP1 (previously called plasma cell membrane glycoprotein 1 or PC-1), is thought to be the most important in PP_i generation (Johnson et al., 2000). A recent, detailed study of the mouse model lacking NPP1 showed that this enzyme is essential for normal bone development and control of physiological bone mineralization (Mackenzie et al., 2012). There is now significant evidence suggesting that extracellular ATP is a key source of PP_i in bone (Orriss et al., 2012; Orriss et al., 2013; Orriss et al., 2007). Thus, it is likely that nucleotide triphosphates exert a dual inhibitory action on bone mineralisation via both P2 receptor mediated signalling and direct hydrolysis to PP_i (**Fig. 3**)

P2X receptors and osteoblasts

The mitogenic effects of ATP were first reported by Suzuki and colleagues in the early 1990's (Suzuki et al., 1993). Subsequent work showed that ATP acting via the P2X5 receptor promotes DNA synthesis in human osteoblast-like cells (Nakamura et al., 2000).

The role of the P2X7 receptor in osteoblast biology has been the focus of significant research albeit often with conflicting results. Expression of the P2X7 receptor was first demonstrated in a subpopulation of differentiated human osteoblasts (Gartland et al., 2001); later studies also described P2X7 receptor expression in rat (Orriss et al., 2006) and mouse osteoblasts (Ke et al., 2003). Early reports suggested that P2X7 receptor activation caused enhanced osteoblast apoptosis (Gartland et al., 2001). Subsequent studies suggested that P2X7 stimulation leads to increased membrane blebbing and bone formation; an effect thought to be mediated via increased production of lyophosphatidic acid (LPA) and prostaglandin E₂ (PGE₂)(Li et al., 2005; Panupinthu et al., 2008; Panupinthu et al., 2007). In contrast, a recent investigation found that Bz-ATP acting via the P2X7 receptor inhibited bone mineralisation (Orriss et al., 2012).

Work using selective P2X7 receptor antagonists indicated that this receptor also plays a role in mediating ATP release from osteoblasts (Brandao-Burch et al., 2012). Furthermore, the P2X7 receptor is thought to mediate the ERK1/2 activation, NF-κB translocation and PGE₂ synthesis caused by fluid shear stress in osteoblast-like cells (Genetos et al., 2005; Liu et al., 2008). Other suggested roles for the P2X7 receptor in osteoblasts include the regulation of metabolic acid production (Grol et al., 2012) and callus remodelling during fracture repair (Li et al., 2009).

To date, several studies using P2X7 receptor knockout mice from different genetic backgrounds have been performed. The initial investigation by Ke *et al.* reported a reduced bone mineral content and periosteal circumference accompanied by decreased bone formation parameters and increased bone resorption parameters (Ke et al., 2003). A separate study by Gartland *et al.* (Gartland et al., 2003b) reported no differences in bone mineral density or trabecular bone but an increase in cortical bone was observed. It was subsequently reported that the knockout model used by Gartland *et al.* expressed a P2X7 splice variant in some tissues (Nicke et al., 2009); therefore the results from this model should be interpreted with caution. Further complicating the analysis of this knockout mouse model, a recent study found that the genetic background strongly influenced the bone phenotype (Syberg et al., 2012b).

Several P2X receptors (P2X2, P2X3, P2X4, P2X6) expressed by osteoblasts still have no known functional effects and present interesting areas for future study. Of particular note is the P2X4 receptor, which displays the highest expression of all the P2 receptors present on osteoblasts (Orriss et al., 2012). Furthermore, since osteoblasts express all P2X receptor subtypes it is plausible that they may combine

to form functional heteromultimers. Heteromultimers often display the pharmacology of both participating subunits, thus increasing the diversity of ATP-mediated signalling and downstream functional effects. At present, it is not known whether functional heteromultimers are expressed by osteoblasts.

P2Y receptors and osteoblasts

Studies into the role of P2Y receptors in osteoblasts have identified a wide range of functional effects. The P2Y₂ receptor was initially thought to mediate intercellular signalling since an early report demonstrated that mechanically stimulated human osteoblasts propagate fast intercellular Ca²⁺ waves via the autocrine activation of P2Y₂ receptors (Jorgensen et al., 2000). A follow-up study showed that intercellular signalling between osteoblasts and osteoclasts was not mediated via P2Y receptors but instead appeared to require the P2X₇ receptor (Jorgensen et al., 2002). However, in a separate investigation P2Y₂ receptors were shown to mediate the Ca²⁺ mobilisation induced by oscillatory fluid flow in mouse osteoblasts (You et al., 2002).

Activation of the P2Y₂ receptor in osteoblast-like cells has been shown to activate a number of intracellular signalling pathways including protein kinase C (PKC), p38 mitogen-activated protein kinase (p38 MAPK), and c-Jun NH₂-terminal protein kinase (JNK) (Costessi et al., 2005; Katz et al., 2006; Katz et al., 2008; Pines et al., 2005).

Purinergic signalling can also interact with other intracellular pathways to regulate osteoblast function. In early studies, P2 receptors were found to mediate the potentiation of parathyroid hormone (PTH) receptor-mediated increases in [Ca²⁺]_i and IP₃ in osteoblast-like cells (Kaplan et al., 1995; Sistare et al., 1995). Later work found that synergistic co-activation by PTH and ATP, acting via P2Y receptors, increased mRNA levels of *c-fos*, a transcription factor important in osteoblast activation and bone remodelling. The authors suggested that this may be a mechanism to generate strong localised responses to systemic growth and differentiation factors (Bowler et al., 1999). A follow-up study demonstrated that PTH potentiated nucleotide-induced [Ca²⁺]_i release in rat osteoblasts independently of G_q activation or cyclic adenosine monophosphate (cAMP) accumulation (Buckley et al., 2001). ATP can also act via P2Y receptors to induce the release of IL-6 (Ihara et al., 2005).

Clopidogrel (Plavix®), a selective P2Y₁₂ receptor antagonist, acts to inhibit platelet aggregation. It is an antithrombotic widely prescribed to reduce the risk of heart attack and stroke. A recent investigation by Syberg and colleagues found that low micromolar concentrations of clopidogrel inhibit osteoblast proliferation, differentiation and function and reduce cell viability *in vitro* (Syberg et al., 2012a). Reduced trabecular bone was also observed in ovariectomised mice treated with clopidogrel for 4 weeks (Syberg et al., 2012a). In contrast, a paper by Su *et al.* demonstrated increased bone in mice treated with clopidogrel (Su et al., 2012). The two investigations used very different dosing regimens (oral dosing in the former compared to clopidogrel treated water in the latter) and thus the divergent results may reflect variations in the final dose received by the animals. In agreement with this notion, a study looking at a cohort of Danish patients found that clopidogrel was associated with fracture risk; however, these effects were biphasic with high doses (the recommended range) being associated with increased fracture risk and low doses being associated with decreased fracture risk (Jorgensen et al., 2012).

Mice lacking the P2Y₁₃ receptor display reduced trabecular bone mass, decreased osteoblast numbers *in vivo* and a reduced rate of bone remodelling (Wang et al., 2012). Osteoblasts cultured from P2Y₁₃ knockout mice also displayed a down regulation of RhoA/ROCK I signalling and a reduced ratio of receptor activator of nuclear factor kappaB ligand (RANKL) / osteoprotegerin expression (Wang et al., 2012). A follow up study by the same group demonstrated that P2Y₁₃ knockout mice display enhanced

osteogenic responses to mechanical loading which the authors attribute to reduced levels of extracellular ATP metabolism (Wang et al., 2013). The P2Y₁₃ receptor has also been shown to play an important role in phosphate metabolism (Wang et al., 2014) and the terminal differentiation of osteoprogenitor cells into osteoblasts or adipocytes (Biver et al., 2013).

ATP release from osteoblasts

Controlled ATP release from osteoblast-like cells was first described by Romanello *et al* in 2001 (Romanello et al., 2001). Since then several studies have indicated that the primary method of ATP release from osteoblasts is by vesicular exocytosis (Genetos et al., 2005; Orriss et al., 2009; Romanello et al., 2005). However, it has been suggested that the P2X₇ receptor may also be involved (Brandao-Burch et al., 2012). The amount of ATP released from osteoblasts depends on their differentiation state, with mature, bone-forming cells releasing up to seven-fold more than immature, proliferating cells (Orriss et al., 2009). Several studies have demonstrated enhanced ATP release in response to different external stimuli including hypoxia (Orriss et al., 2009), mechanical stress (Hecht et al., 2013), fluid flow (Genetos et al., 2005; Romanello et al., 2005), vitamin D (Biswas et al., 2009) and ultrasound (Alvarenga et al., 2010; Hayton et al., 2005). A recent investigation examined the effect of different forms of mechanical stimulation on ATP release from human osteoblast-like SaOS-2 cells, including turbulent fluid flow, laminar fluid flow, substrate strain and 3D compressive loading (Rumney et al., 2012). Rumney *et al* concluded that the concentration of ATP released in response to mechanical loading varied in a time-, direction- and strain-dependent manner representing a local mechanostat in bone that could influence bone mineralisation (Rumney et al., 2012). Osteoblasts cultured from P2Y₁₃ receptor knockout mice display increased levels of extracellular ATP suggesting a role for this receptor in the modulation of ATP metabolism (Wang et al., 2013).

P1 receptor signalling and osteoblasts

The role of adenosine and P1 receptor-mediated signalling in the regulation of bone cell function has become an area of increasing interest in recent years. Osteoblasts have been shown to express all four P1 receptor subtypes (Gharibi et al., 2011; Vincenzi et al., 2013); however, the actions of extracellular adenosine on osteoblasts appear to be less clear-cut than those of ATP. Initial work found that synthetic adenosine analogues elicited a receptor-mediated rise in cAMP levels in calvarial osteoblasts (Lerner et al., 1987). Subsequently, it was reported that adenosine acts as a mitogen for osteoblast-like cells by stimulating DNA synthesis (Shimegi, 1996). An important early study found that human osteoblast precursors produced extracellular adenosine, probably via the breakdown of released ATP, which modulated their secretion of Il-6 and osteoprotegerin (Evans et al., 2006).

Since the late 1990's, several studies have failed to find an effect of adenosine on mineralised bone nodule formation by rat calvarial osteoblasts (Jones et al., 1997; Hoebertz et al., 2002). However, another study indicated that adenosine, acting via the A_{2B} receptor, may increase the osteogenic differentiation of rat long bone mesenchymal stem cells (Gharibi et al., 2011). Furthermore, a synthetic A_{2B} receptor agonist has been shown to increase bone formation, and osteoblasts from A_{2B} receptor knockout mice display reduced activity *in vitro* (Carroll et al., 2012). Recently, it was reported that the adenosine generated by ecto-5'-nucleotidase (CD-73) was important in promoting osteoblast differentiation (Takedachi et al., 2012). In contrast, adenosine analogues, acting via the A₁ or A_{2A} receptors, have been shown to inhibit the differentiation of rodent osteoblast-like cells (Gharibi et al., 2012).

2.2. Purinergic signalling and mesenchymal stem cells

Osteoblasts are derived from mesenchymal stem cells (MSCs) and thus these cells play a key role within the bone. Although information is limited a role for extracellular nucleotides in the regulation of MSC proliferation and differentiation is emerging. Initial work showed that fluid flow induced vesicular release of ATP from human MSCs, which then stimulated proliferation via a P2 receptor-mediated increase in $[Ca^{2+}]_i$ (Riddle et al., 2007). More recent studies have shown that human MSCs express the P2X3-7 receptors and all P2Y receptors (Noronha-Matos et al., 2012; Zippel et al., 2012). P2 receptors and extracellular nucleotides are now thought to be important modulators of human MSC differentiation, with the P2X6, P2Y₄, P2Y₆ and P2Y₁₄ receptors identified as playing pivotal roles in this process (Ferrari et al., 2011; Zippel et al., 2012; Noronha-Matos et al., 2012). Furthermore, a detailed investigation using knockout mice identified that the P2Y₁₃ receptor regulates the terminal differentiation of MSCs into osteoblasts or adipocytes (Biver et al., 2013). It has also been reported that NTPDases influence which osteoblast progenitors are driven into proliferation or differentiation (Noronha-Matos et al., 2012).

To date, many of the studies investigating the role of adenosine in osteoblasts have utilised MSCs (as described above). In addition, A₁, A_{2A}, A_{2B} and A₃ receptor activation has been shown to increase the proliferation (mainly via A₁ and A_{2A} receptors) and differentiation (via A_{2B} receptors) of MSCs (Costa et al., 2011).

2.3. P2 receptors and osteocytes

Osteocytes are the most abundant cell type within bone (Jande et al., 1973) yet little is known about the role of purinergic signalling in their survival and function. Because they are embedded within the bone matrix, primary osteocytes are difficult to isolate in significant amounts; therefore most work to date has been performed on osteocyte-like cell lines. An early study found that the calcium signals induced by fluid flow were decreased by suramin and thapsigargin, suggesting involvement of ATP acting via P2Y receptors (Huo et al., 2008). Expression of functional P2X2, P2X7, P2Y₂, P2Y₄, P2Y₁₂ and P2Y₁₃ receptors on MLO-Y4 osteocyte-like cells has recently been described (Kringelbach et al., 2014)

Several studies have also reported that osteocytes release ATP in a controlled manner (Genetos et al., 2007; Kringelbach et al., 2014; Thompson et al., 2011). This release is enhanced by UTP and mechanical stimulation and is thought to be mediated primarily by vesicular exocytosis (Kringelbach et al., 2014). Mechanically induced ATP release can also be regulated by the $\alpha_2\delta_1$ auxiliary subunit of T-type voltage sensitive calcium channels (Thompson et al., 2011).

Since ATP can inhibit bone mineralisation it is possible that release from osteocytes *in vivo* could act to reduce progressive, age-related mineral encroachment from the surrounding bone, thus preventing eventual cell 'fossilisation' and death.

2.4. Purinergic signalling and osteoclasts

The first description of P2 receptor expression by osteoclasts was in the early 1990's, when it was shown that ATP induced a rapid and transient increase in $[Ca^{2+}]_i$ in rabbit osteoclasts (Yu et al., 1993a). The human P2U (P2Y₂/P2Y₄) receptor was subsequently cloned and sequenced from osteoclastoma, indicating that this receptor was probably expressed by osteoclasts (Bowler et al., 1995). Early investigations also provided evidence for the presence of P2X receptors, giving rise to non-selective cation currents, and P2Y receptors, mediating Ca²⁺ release from intracellular stores (Weidema et al., 1997; Wiebe et al., 1999). Initially it was thought that P2X4 receptors mediated the ATP-activated non-selective cation currents in rabbit osteoclasts (Naemsch et al., 1999). However, it was subsequently found that the nucleotide-induced elevation of $[Ca^{2+}]_i$ arose primarily from activation of P2Y receptors (Weidema et al., 2001)(Weidema AF 2001).

Expression of multiple P2 receptor subtypes by osteoclasts has now been described (see review (Burnstock et al., 2013)). Initial studies demonstrated expression of mRNA and protein for the P2X2, P2X4, P2X7, P2Y₁ and P2Y₂ receptors by rat long bone-derived osteoclasts (Hoebertz et al., 2000). Osteoclasts generated from human peripheral blood were later shown to express the P2X1, P2X4, P2X7, P2Y₁, P2Y₂, P2Y₄, P2Y₆ and P2Y₁₁ receptors (Buckley et al., 2002). Recently, osteoclasts derived from mouse bone marrow were found to express all P2 receptors, with the exception of the P2X6 and P2Y₄ receptors (Orriss et al., 2011b). This study also demonstrated large differences in the expression levels of the different P2 receptors, with the P2Y₆ receptor being the most abundant (Orriss et al., 2011b).

Functional effects of P2 receptor-mediated signalling in osteoclasts

The first indication that purinergic signalling could play a role in the regulation of osteoclast function came in 1995 when Bowler *et al.* reported that ATP stimulated resorption by cells from human osteoclastoma (Bowler et al., 1995). Originally, the P2Y₂ receptor was thought to mediate this effect; however, in a follow up study UTP failed to stimulate resorption suggesting this was not the case (Bowler et al., 1998). Subsequently ATP was found to increase the formation and activity of rodent osteoclasts, effects that were inhibited by suramin or apyrase (Morrison et al., 1998). These stimulatory actions on resorption were further enhanced when osteoclasts were first activated by culture in acidified medium (Morrison et al., 1998). Since the P2X2 receptor is the only P2 receptor that requires extracellular acidification to show its full sensitivity to ATP (Wildman et al., 1998) it was suggested that the P2X2 receptor could mediate these pro-resorptive effects. Further investigation revealed that ADP, 2-meththioADP and ATP all stimulated osteoclast formation and activity, suggesting involvement of P2Y receptors. Pharmacological and cytochemical evidence indicated that these effects were mediated by the P2Y₁ receptor (Hoebertz et al., 2001).

DEXA and μ CT analysis of P2Y₁ receptor knockout mice showed reduced trabecular bone in the long bones (Orriss et al., 2011a; Orriss et al., 2008) (**Fig. 1**). Given the suggested role for the P2Y₁ receptor in promoting osteoclast activity this decrease in bone mass was unexpected. However, a recent study found that ADP still stimulated osteoclast formation and resorption in cells derived from P2Y₁ receptor knockout mice (Su et al., 2012), indicating the involvement of another P2Y receptor. Based on pharmacology, the P2Y₁₂ receptor was the logical alternative candidate for mediating these effects. The P2Y₁₂ receptor was only identified in 2001 (Hollopeter et al., 2001), the same year as the study by Hoebertz *et al.*, and thus it was not considered in the original investigation. Expression of the P2Y₁₂ receptor by osteoclasts has recently been reported (Orriss et al., 2011b) and P2Y₁₂ receptor knockout mice display impaired responses to ADP, increased trabecular bone and decreased arthritis associated bone loss (Su et al., 2012). Furthermore, the P2Y₁₂ receptor antagonist, clopidogrel, inhibits osteoclast formation, viability and resorptive activity (Syberg et al., 2012a). Taken together these data suggest a role for the P2Y₁₂ receptor in mediating the effects of ADP on osteoclast formation and activity.

The P2Y₆ receptor also appears to regulate osteoclast formation, activity and survival. UDP, acting via the P2Y₆ receptor, has been shown to prevent the apoptosis induced by tumour necrosis factor- α (TNF- α), induce the translocation and activation of NF- κ B and stimulate osteoclast formation and activity (Korcok et al., 2005; Orriss et al., 2011b). Consistent with a role for this receptor in bone resorption, osteoclasts derived from P2Y₆ receptor knockout mice display defective activity (Orriss et al., 2011b). Analysis of P2Y₆ receptor knockout mice by μ CT revealed increased cortical bone in the long bones and spine (Orriss et al., 2011b) (**Fig. 1**).

The role of the P2Y₁₃ receptor in regulating osteoclast function is still unclear. However, P2Y₁₃ receptor knockout mice display reduced osteoclast numbers *in vivo* and protection from ovariectomy-induced bone loss in females (Wang et al., 2012).

The regulation of osteoclast formation, activity and survival by the P2X₇ receptor has been the focus of many, often conflicting, studies. P2X₇ receptor blockade with selective antagonists or a monoclonal antibody has been shown to inhibit osteoclast formation (Agrawal et al., 2010; Gartland et al., 2003a). Analogues of the P2X₇ receptor antagonist, KN-62, have also been shown to induce osteoclast apoptosis (Penolazzi et al., 2005). Combined, these studies suggest a role for the P2X₇ receptor in osteoclast formation. However, other investigations have demonstrated that P2X₇ receptor knockout mice possess functional osteoclasts *in vivo* and that osteoclasts can be generated *in vitro*, using knockout precursor cells (Gartland et al., 2003b; Ke et al., 2003). Furthermore, prolonged exposure to ATP has also been shown to down-regulate P2X₇ receptor function and inhibit osteoclast formation and activity (Hiken et al., 2004; Naemsch et al., 2001).

Other work has demonstrated that P2X₇ receptor activation induces the translocation of NFκB (Korcok et al., 2004) and an isoform specific PKC in osteoclasts and their precursors (Armstrong et al., 2009). The P2X₇ receptor may additionally play a role in the intercellular communication between osteoblasts and osteoclasts, cytoskeletal reorganisation at the sealing zone and the delivery and secretion of lytic granules into the resorption lacunae (Hazama et al., 2009; Jorgensen et al., 2002). Recently, it was reported that ATP release from osteoclasts is mediated via the P2X₇ receptor (Brandao-Burch et al., 2012; Pellegatti et al., 2011). Thus, this receptor may promote osteoclast fusion by increasing the concentration of extracellular adenosine (via breakdown of released ATP) (Pellegatti et al., 2011). It has also been suggested that ATP stimulates human osteoclast activity indirectly through upregulation of osteoblast-expressed RANKL (Buckley et al., 2002).

At present, it is not known how activation of the other P2 receptor subtypes influences osteoclast formation and activity and this presents an interesting area for further study.

P1 receptors, adenosine and osteoclasts

A potential role for adenosine in regulating osteoclast function was first suggested in the 1980s when the P1 receptor agonist, 2-chloroadenosine, was shown to stimulate bone resorption in calvarial bones in organ culture (Lerner et al., 1983). However, a later investigation found that adenosine analogues had no effects on resorption in this culture system (Lerner et al., 1987). In agreement, several studies have reported that adenosine has no effect on the formation or resorptive activity of rodent osteoclasts *in vitro* (Hoebertz et al., 2001; Morrison et al., 1998). Adenosine was also reported to have no effect on intracellular calcium levels in rabbit osteoclasts (Korcok et al., 2004). However, more recent work has suggested that stimulation of the A_{2A} receptor by adenosine or selective agonists can both stimulate (Pellegatti et al., 2011) and inhibit (Mediero et al., 2012b) osteoclast formation. A_{2A} receptor activation has also been shown to prevent the osteolysis that occurs during prosthesis loosening, a common cause of joint implant failure (Mediero et al., 2012a).

Pellegatti *et al* (2011) reported that the A₁ receptor was only weakly expressed by osteoclasts and activation of the A₁ receptor was recently shown to have no effect on mouse osteoclasts (Pellegatti et al., 2011; He et al., 2012). In contrast, blockade or deletion of the A₁ receptor can reduce the formation of mouse osteoclasts in culture (Kara et al., 2010a) and A₁ receptor knockout mice display increased bone mineral density and resistance to ovariectomy-induced bone loss (Kara et al., 2010b)

Adenosine has also been reported to stimulate osteoclastogenesis indirectly. Evans *et al* (2006) suggested that the adenosine produced from the hydrolysis of released ATP acts on P1 receptors to cause IL-6 release and inhibition of osteoprotegerin secretion (Evans *et al.*, 2006).

2.5. P2 receptor polymorphisms and bone

Single nucleotide polymorphisms (SNPs) have been described in several P2 receptor genes (P2X4, P2X7, P2Y₁, P2Y₂, P2Y₆, P2Y₁₃). To date, only SNPs in the genes for the P2Y₂, P2X4 and P2X7 receptors have been studied for an association with osteoporosis risk or bone turnover (see review (Wesselius *et al.*, 2012)).

At present, two SNPs have been identified which alter P2Y₂ receptor activity. The first of these is located at position 1000 (cytosine to thymine), changing arginine to cysteine at residue 334 (Arg³³⁴Cys) (Janssens *et al.*, 1999). The second SNP is located at position 936 (guanine to cytosine) and changes arginine to serine at residue 312 (Arg³¹²Ser) (Buscher *et al.*, 2006). A recent study examining the association between the Arg³¹²Ser SNP and bone status found that subjects homozygous for the C (variant) allele of the P2Y₂ gene had a significantly increased bone mineral density (BMD) at menopause (Wesselius *et al.*, 2013a). Furthermore, the rate of bone loss for up to ten years post-menopause was 20-30% lower than both the heterozygous and the homozygous for the common allele (Wesselius *et al.*, 2013a).

Many SNPs have been identified in the P2X7 receptor gene (see the recent review by Wesselius *et al* (Wesselius *et al.*, 2012) for detailed information). Loss of function polymorphisms in the P2X7 receptor have been associated with increased fracture risk, reduced BMD and osteoporosis (Gartland *et al.*, 2012b; Husted *et al.*, 2013; Ohlendorff *et al.*, 2007). Recently, it was reported that a single SNP in the P2X4 receptor gene (tyrosine to cysteine at position 315) is associated with an increased risk of osteoporosis and lower lumbar spine BMD (Wesselius *et al.*, 2013b).

3. Cartilage

Cartilage is a flexible connective tissue that is found throughout the body, including the surface of joints. It is composed of specialised cells called chondrocytes that synthesize large amounts of collagen, abundant ground substance rich in proteoglycans and elastin fibres. Unlike other connective tissues, cartilage does not have a blood supply and chondrocytes receive their nutrients by diffusion. This lack of vasculature means that, compared to other tissues, cartilage grows and repairs itself more slowly.

P2 receptor expression by chondrocytes

Purinergic receptor expression in cartilage was first suggested in 1990 when ATP was found to stimulate resorption of bovine nasal cartilage (Leong *et al.*, 1990). The following year, ATP was found to increase the production of PGE₂ from articular chondrocytes (Caswell *et al.*, 1991). Several early studies reported that chondrocytes were responsive to UTP as well as ATP indicating the presence of P2Y receptors (Caswell *et al.*, 1992; Kaplan *et al.*, 1996; Koolpe *et al.*, 1997). Chondrocytes are now known to express multiple P2 receptor subtypes including the P2X1, P2X2, P2X3, P2X4, P2X7, P2Y₁ and P2Y₂ receptors (Koolpe *et al.*, 1999; Knight *et al.*, 2009; Varani *et al.*, 2008b; Kudirka *et al.*, 2007). Many of these receptors (P2X2, P2X4, P2X7 and P2Y₁) are expressed throughout all the zones of articular cartilage, whilst others (e.g. the P2Y₂ receptor) show more limited spatial expression (Knight *et al.*, 2009). A number of investigations have shown increased [Ca²⁺] levels following treatment with ATP, ADP or UDP, indicating the presence of functional P2 receptors on chondrocytes (Bulman *et al.*, 1995; Hung *et al.*, 1997; Kono *et al.*, 2006).

Functional effects of P2 receptor signalling in chondrocytes

To date, many of the studies describing the effects of purinergic signalling on chondrocytes have been conflicting with some reports suggesting that extracellular nucleotides negatively regulate cartilage metabolism whilst others describe beneficial effects of ATP. ATP has been reported to inhibit cartilage formation in micromass cultures (Meyer et al., 2001), promote proteoglycan breakdown and glycosaminoglycan release in bovine nasal cartilage (Brown et al., 1997) and increase the production of inflammatory mediators, nitric oxide (NO) and PGE₂ (Varani et al., 2008b). Furthermore, P2Y₂ receptor stimulation increased IL-1-mediated PGE₂ release from articular chondrocytes (Koolpe et al., 1999) and induced a rapid rise in PGE₂ synthesis via the ERK1/2 and p38 MAPK signalling pathways (Berenbaum et al., 2003). In contrast, early work suggested that ATP caused articular cartilage mineralisation by promoting matrix vesicle-mediated calcium deposition (Ryan et al., 1992; Hsu, 1992). Extracellular ATP and UTP have also been shown to stimulate cartilage proteoglycan and collagen accumulation and suppress inflammatory mediator (NO) production (Chowdhury et al., 2006; Croucher et al., 2000).

Other reported effects of purinergic signalling include increased chondrocyte differentiation (Fodor et al., 2009), elevated fibroblast growth factor-induced proliferation (Kaplan et al., 1996; Koolpe et al., 1997) and increased responsiveness to IL-1 β and TNF- α (Caswell et al., 1992; Leong et al., 1993).

ATP release and breakdown in cartilage

The first direct evidence for constitutive ATP release from chondrocytes came in 2000 from a study by Graff *et al.* (Graff et al., 2000). Subsequent studies have shown that P2Y receptor stimulation increases [Ca²⁺]_i levels which trigger further release of ATP from adjacent cells (Kono et al., 2006; Millward-Sadler et al., 2004).

Ecto-nucleotidase activity in chondrocytes was first reported by Kanabe *et al.* in 1983 (Kanabe et al., 1983). It was later shown that NTPDases, NPPs, alkaline phosphatase and ecto-5'-nucleotidase, as well as adenylate kinase and nucleoside diphosphokinase which mediate regeneration of ATP, are all present in cartilage (Graff et al., 2003). A recent study found that P1 receptor agonists decreased extracellular PP_i production in chondrocytes and cartilage explants, while ADP increased PP_i levels, suggesting a homeostatic role for P1 and P2 receptors in stabilizing concentrations of PP_i (Rosenthal et al., 2010).

P1 receptors and chondrocytes

Chondrocytes are known to express all the P1 receptors (Tesch et al., 2002; Varani et al., 2008a); however, information regarding the functional effects of receptor activation remains limited. Pulsed electro-magnetic fields have been shown to promote chondrocyte proliferation via the up-regulation of A_{2A} and A₃ receptor expression (Varani et al., 2008a). Adenosine, acting via A_{2A} receptors, can also reduce IL-1 β -induced inflammation (Campo et al., 2012) and VEGF secretion (Vincenzi et al., 2013).

4. Muscle

There are three types of muscle, skeletal, cardiac and smooth within the body. Cardiac and smooth muscle are often termed involuntary as these muscles contract without conscious thought. Whilst cardiac muscle is only found within the heart, smooth muscle is more widely distributed being found within the walls of many internal organs. Skeletal (or voluntary) muscle is anchored to bones by tendons. It plays a key role in locomotion and posture and thus is a critical component of the musculoskeletal system. Within muscle, extracellular ATP comes from a number of sources including

motor nerves, stressed or exercising muscle and damaged or dying muscle and vascular endothelial cells (Buvinic et al., 2009; Mortensen et al., 2009; Tu et al., 2012).

Skeletal muscle and purinergic signalling

The first evidence for involvement of purinergic signalling in skeletal muscle came in the 1980's when ATP analogues were found to induce the lengthening of muscle fibres in rigor (Schoenberg, 1989). Activation of P2 receptors was later shown to induce IP₃ accumulation in cultured skeletal muscle cells (Keresztes et al., 1991). Many studies have demonstrated the expression of multiple P2 receptor subtypes by skeletal muscle (Banachewicz et al., 2005; Cheung et al., 2003; Deli et al., 2007; Janssens et al., 1996); however, the pattern of purinergic receptor expression is strongly influenced by developmental stage and species (see (Burnstock et al., 2013)).

There is now abundant evidence showing that purinergic signalling plays an important role in modulating the development and function of skeletal muscle (see review by (Burnstock et al., 2013) for detailed information). For example, several studies have shown that extracellular ATP can regulate myoblast proliferation and the differentiation of mammalian skeletal muscle (Martinello et al., 2011; Ryten et al., 2002; Sciancalepore et al., 2012). Extracellular nucleotides, acting via the P2Y₁ receptor, can also enhance muscle contraction and influence muscle excitability (Choi et al., 2003; Voss, 2009).

5. Summary and future directions

Our knowledge about the role purinergic signalling in the musculoskeletal system has increased significantly in the past twenty years. In particular, it is now evident that P2 receptor-mediated signalling exerts complex, local effects on bone cell function. The effects of this signalling system are influenced by a number of factors including the receptor subtype, the extracellular nucleotides present locally and the expression of ecto-nucleotidases. At present, there is abundant evidence for the negative actions of extracellular nucleotides, with particularly notable effects on osteoblast function and matrix mineralisation. However, given that purinergic signalling also exerts some positive effects on bone cells, it is likely that a balance exists *in vivo* between the positive and negative actions of this complex signalling system. Selective receptor agonists and antagonists for the P2 receptor subtypes involved in bone remodelling are being developed, which ultimately could lead to new ways to treat bone disease.

Bone, cartilage and muscle are often studied in isolation and the ability of one tissue to influence another is often overlooked. However, given the significant ATP release and widespread expression of purinergic receptors in the musculoskeletal system crosstalk between these tissues is likely and presents an important area for future work.

Acknowledgements

The author is grateful for funding from Arthritis Research UK and would like to thank Prof. Tim Arnett, University College London, for his support and expertise.

Figure Legends

Figure 1. Bone phenotypes of P2Y₁, P2Y₂ and P2Y₆ receptor knockout mice

Representative μ CT images of the femurs of 8-week old P2Y₁, P2Y₂ and P2Y₆ receptor knockout mice show that receptor deletion causes significant changes in the bone structure. **(A)** P2Y₁ null mice have decreased trabecular bone volume and trabecular number. **(B)** P2Y₂ receptor knockout animals display increased trabecular bone volume, thickness and number. **(C)** P2Y₆ null mice have a higher cortical bone volume and thickness.

Figure 2. Expression of P2 receptors by osteoblasts

The expression of P2 receptors by primary rat osteoblasts was studied by immunofluorescence using specific primary polyclonal antibodies, Cy3-labelled anti-rabbit secondary antibody (red) and DAPI nuclear stain (blue). Expression of protein for all the P2 receptors was detected in in rat osteoblasts. Scale bar = 25 μ m.

Figure 3. Extracellular ATP acts via P2-receptor dependent and independent mechanisms to inhibit bone mineralisation.

Osteoblasts constitutively release ATP into the bone microenvironment, where it can act via two different mechanisms to inhibit mineralisation. (1) *P2-receptor dependent*: activation of the P2Y₂, P2X₁ or P2X₇ receptor decreases TNAP activity and PP_i breakdown. (2) *Receptor independent*: hydrolysis of ATP by NPP1 produces PP_i. The overall effect of either mechanism is increased levels of extracellular PP_i, the key physicochemical inhibitor of mineralisation.

References

- Abbracchio, M.P., Burnstock, G. 1994. Purinoceptors: are there families of P2X and P2Y purinoceptors? *Pharmacol Ther* 64, 445-475.
- Agrawal, A., Buckley, K.A., Bowers, K., Furber, M., Gallagher, J.A., Gartland, A. 2010. The effects of P2X7 receptor antagonists on the formation and function of human osteoclasts in vitro. *Purinergic Signal* 6, 307-315.
- Alvarenga, E.C., Rodrigues, R., Caricati-Neto, A., Silva-Filho, F.C., Paredes-Gamero, E.J., Ferreira, A.T. 2010. Low-intensity pulsed ultrasound-dependent osteoblast proliferation occurs by via activation of the P2Y receptor: role of the P2Y1 receptor. *Bone* 46, 355-362.
- Armstrong, S., Pereverzev, A., Dixon, S.J., Sims, S.M. 2009. Activation of P2X7 receptors causes isoform-specific translocation of protein kinase C in osteoclasts. *J Cell Sci* 122, 136-144.
- Banachewicz, W., Suplat, D., Krzeminski, P., Pomorski, P., Baranska, J. 2005. P2 nucleotide receptors on C2C12 satellite cells. *Purinergic Signal* 1, 249-257.
- Berenbaum, F., Humbert, L., Bereziat, G., Thirion, S. 2003. Concomitant recruitment of ERK1/2 and p38 MAPK signalling pathway is required for activation of cytoplasmic phospholipase A2 via ATP in articular chondrocytes. *J Biol Chem* 278, 13680-13687.
- Biswas, P., Zanello, L.P. 2009. 1 α ,25(OH) $_2$ vitamin D(3) induction of ATP secretion in osteoblasts. *J Bone Miner Res* 24, 1450-1460.
- Biver, G., Wang, N., Gartland, A., Orriss, I., Arnett, T.R., Boeynaems, J.M., Robaye, B. 2013. Role of the P2Y $_{13}$ receptor in the differentiation of bone marrow stromal cells into osteoblasts and adipocytes. *Stem Cells* 31, 2747-2758.
- Bonewald, L.F. 2011. The amazing osteocyte. *J Bone Miner Res* 26, 229-238.
- Bowler, W.B., Birch, M.A., Gallagher, J.A., Bilbe, G. 1995. Identification and cloning of human P2U purinoceptor present in osteoclastoma, bone, and osteoblasts. *J Bone Miner Res* 10, 1137-1145.
- Bowler, W.B., Dixon, C.J., Halleux, C., Maier, R., Bilbe, G., Fraser, W.D., Gallagher, J.A., Hipskind, R.A. 1999. Signaling in human osteoblasts by extracellular nucleotides. Their weak induction of the c-fos proto-oncogene via Ca $^{2+}$ mobilization is strongly potentiated by a parathyroid hormone/cAMP-dependent protein kinase pathway independently of mitogen-activated protein kinase. *J Biol Chem* 274, 14315-14324.
- Bowler, W.B., Littlewood-Evans, A., Bilbe, G., Gallagher, J.A., Dixon, C.J. 1998. P2Y $_2$ receptors are expressed by human osteoclasts of giant cell tumor but do not mediate ATP-induced bone resorption. *Bone* 22, 195-200.
- Brandao-Burch, A., Key, M.L., Patel, J.J., Arnett, T.R., Orriss, I.R. 2012. The P2X7 Receptor is an important regulator of extracellular ATP levels. *Front Endocrinol (Lausanne)* 3, 41.
- Brown, C.J., Caswell, A.M., Rahman, S., Russell, R.G., Buttle, D.J. 1997. Proteoglycan breakdown from bovine nasal cartilage is increased, and from articular cartilage is decreased, by extracellular ATP. *Biochim Biophys Acta* 1362, 208-220.
- Buckley, K.A., Golding, S.L., Rice, J.M., Dillon, J.P., Gallagher, J.A. 2003. Release and interconversion of P2 receptor agonists by human osteoblast-like cells. *FASEB J* 17, 1401-1410.
- Buckley, K.A., Hipskind, R.A., Gartland, A., Bowler, W.B., Gallagher, J.A. 2002. Adenosine triphosphate stimulates human osteoclast activity via upregulation of osteoblast-expressed receptor activator of nuclear factor-kappa B ligand. *Bone* 31, 582-590.

- Buckley, K.A., Wagstaff, S.C., McKay, G., Gaw, A., Hipskind, R.A., Bilbe, G., Gallagher, J.A., Bowler, W.B. 2001. Parathyroid hormone potentiates nucleotide-induced $[Ca^{2+}]_i$ release in rat osteoblasts independently of Gq activation or cyclic monophosphate accumulation. A mechanism for localizing systemic responses in bone. *J Biol Chem* 276, 9565-9571.
- Bulman, W.A., Iannotti, J.P., Glowacki, K., Bleuit, J., Clark, C.C. 1995. Serum fractions and related agonists with calcium-mobilizing activity in the bovine growth plate chondrocyte. *J Orthop Res* 13, 220-229.
- Burnstock, G. 1972. Purinergic nerves. *Pharmacol Rev* 24, 509-581.
- Burnstock, G. 2007. Purine and pyrimidine receptors. *Cell Mol Life Sci* 64, 1471-1483.
- Burnstock, G., Arnett, T.R., Orriss, I.R. 2013. Purinergic signalling in the musculoskeletal system. *Purinergic Signal* 9, 541-572.
- Burnstock, G., Kennedy, C. 1985. Is there a basis for distinguishing two types of P2-purinoceptor? *Gen Pharmacol* 16, 433-440.
- Buscher, R., Hoerning, A., Patel, H.H., Zhang, S., Arthur, D.B., Grasemann, H., Ratjen, F., Insel, P.A. 2006. P2Y₂ receptor polymorphisms and haplotypes in cystic fibrosis and their impact on Ca²⁺ influx. *Pharmacogenet Genomics* 16, 199-205.
- Buvinic, S., Almarza, G., Bustamante, M., Casas, M., Lopez, J., Riquelme, M., Saez, J.C., Huidobro-Toro, J.P., Jaimovich, E. 2009. ATP released by electrical stimuli elicits calcium transients and gene expression in skeletal muscle. *J Biol Chem* 284, 34490-34505.
- Campo, G.M., Avenoso, A., D'Ascola, A., Prestipino, V., Scuruchi, M., Nastasi, G., Calatroni, A., Campo, S. 2012. The stimulation of adenosine 2A receptor reduces inflammatory response in mouse articular chondrocytes treated with hyaluronan oligosaccharides. *Matrix Biol* 31, 338-351.
- Carroll, S.H., Wigner, N.A., Kulkarni, N., Johnston-Cox, H., Gerstenfeld, L.C., Ravid, K. 2012. A2B adenosine receptor promotes mesenchymal stem cell differentiation to osteoblasts and bone formation in vivo. *J Biol Chem* 287, 15718-15727.
- Caswell, A.M., Leong, W.S., Russell, R.G. 1991. Evidence for the presence of P2-purinoceptors at the surface of human articular chondrocytes in monolayer culture. *Biochim Biophys Acta* 1074, 151-158.
- Caswell, A.M., Leong, W.S., Russell, R.G. 1992. Interleukin-1 beta enhances the response of human articular chondrocytes to extracellular ATP. *Biochim Biophys Acta* 1137, 52-58.
- Cheung, K.K., Ryten, M., Burnstock, G. 2003. Abundant and dynamic expression of G protein-coupled P2Y receptors in mammalian development. *Dev Dyn* 228, 254-266.
- Choi, R.C., Siow, N.L., Cheng, A.W., Ling, K.K., Tung, E.K., Simon, J., Barnard, E.A., Tsim, K.W. 2003. ATP acts via P2Y₁ receptors to stimulate acetylcholinesterase and acetylcholine receptor expression: transduction and transcription control. *J Neurosci* 23, 4445-4456.
- Chowdhury, T.T., Knight, M.M. 2006. Purinergic pathway suppresses the release of .NO and stimulates proteoglycan synthesis in chondrocyte/agarose constructs subjected to dynamic compression. *J Cell Physiol* 209, 845-853.
- Costa, M.A., Barbosa, A., Neto, E., Sa-e-Sousa, A., Freitas, R., Neves, J.M., Magalhaes-Cardoso, T., Ferreirinha, F., Correia-de-Sa, P. 2011. On the role of subtype selective adenosine receptor agonists during proliferation and osteogenic differentiation of human primary bone marrow stromal cells. *J Cell Physiol* 226, 1353-1366.
- Costessi, A., Pines, A., D'andrea, P., Romanello, M., Damante, G., Cesaratto, L., Quadrifoglio, F., Moro, L., Tell, G. 2005. Extracellular nucleotides activate Runx2 in the osteoblast-like HOBIT cell line: a possible molecular link between mechanical stress and osteoblasts' response. *Bone* 36, 418-432.

Croucher, L.J., Crawford, A., Hatton, P.V., Russell, R.G., Buttle, D.J. 2000. Extracellular ATP and UTP stimulate cartilage proteoglycan and collagen accumulation in bovine articular chondrocyte pellet cultures. *Biochim Biophys Acta* 1502, 297-306.

Deli, T., Szappanos, H., Szigeti, G.P., Cseri, J., Kovacs, L., Csernoch, L. 2007. Contribution from P2X and P2Y purinoreceptors to ATP-evoked changes in intracellular calcium concentration on cultured myotubes. *Pflugers Arch* 453, 519-529.

Dixon, C.J., Bowler, W.B., Walsh, C.A., Gallagher, J.A. 1997. Effects of extracellular nucleotides on single cells and populations of human osteoblasts: contribution of cell heterogeneity to relative potencies. *Br J Pharmacol.* 120, 777-780.

Evans, B.A., Elford, C., Pexa, A., Francis, K., Hughes, A.C., Deussen, A., Ham, J. 2006. Human osteoblast precursors produce extracellular adenosine, which modulates their secretion of IL-6 and osteoprotegerin. *J Bone Miner Res* 21, 228-236.

Ferrari, D., Gulinelli, S., Salvestrini, V., Lucchetti, G., Zini, R., Manfredini, R., Caione, L., Piacibello, W., Ciciarello, M., Rossi, L., Idzko, M., Ferrari, S., Di Virgilio, F., Lemoli, R.M. 2011. Purinergic stimulation of human mesenchymal stem cells potentiates their chemotactic response to CXCL12 and increases the homing capacity and production of proinflammatory cytokines. *Exp Hematol* 39, 360-374, 374 e361-365.

Fleisch, H., Bisaz, S. 1962. Isolation from urine of pyrophosphate, a calcification inhibitor. *Am J Physiol* 203, 671-675.

Fleisch, H., Neuman, W. 1961. The role of phosphatase and polyphosphates in calcification of collagen. *Helv Physiol Pharmacol Acta* 19, C17-C18.

Fodor, J., Matta, C., Juhasz, T., Olah, T., Gonczi, M., Szijgyarto, Z., Gergely, P., Csernoch, L., Zakany, R. 2009. Ionotropic purinergic receptor P2X4 is involved in the regulation of chondrogenesis in chicken micromass cell cultures. *Cell Calcium* 45, 421-430.

Gartland, A., Buckley, K.A., Hipskind, R.A., Bowler, W.B., Gallagher, J.A. 2003a. P2 receptors in bone-modulation of osteoclast formation and activity via P2X7 activation. *Crit Rev Eukaryot Gene Expr* 13, 237-242.

Gartland, A., Buckley, K.A., Hipskind, R.A., Perry, M.J., Tobias, J.H., Buell, G., Chessell, I., Bowler, W.B., Gallagher, J.A. 2003b. Multinucleated osteoclast formation in vivo and in vitro by P2X7 receptor-deficient mice. *Crit Rev Eukaryot Gene Expr* 13, 243-253.

Gartland, A., Hipskind, R.A., Gallagher, J.A., Bowler, W.B. 2001. Expression of a P2X7 receptor by a subpopulation of human osteoblasts. *J Bone Miner Res* 16, 846-856.

Gartland, A., Orriss, I.R., Rumney, R.M., Bond, A.P., Arnett, T., Gallagher, J.A. 2012a. Purinergic signalling in osteoblasts. *Front Biosci* 17, 16-29.

Gartland, A., Skarratt, K.K., Hocking, L.J., Parsons, C., Stokes, L., Jorgensen, N.R., Fraser, W.D., Reid, D.M., Gallagher, J.A., Wiley, J.S. 2012b. Polymorphisms in the P2X7 receptor gene are associated with low lumbar spine bone mineral density and accelerated bone loss in post-menopausal women. *Eur J Hum Genet* 20, 559-564.

Genetos, D.C., Geist, D.J., Liu, D., Donahue, H.J., Duncan, R.L. 2005. Fluid Shear-Induced ATP Secretion Mediates Prostaglandin Release in MC3T3-E1 Osteoblasts. *J Bone Miner Res* 20, 41-49.

Genetos, D.C., Kephart, C.J., Zhang, Y., Yellowley, C.E., Donahue, H.J. 2007. Oscillating fluid flow activation of gap junction hemichannels induces ATP release from MLO-Y4 osteocytes. *J Cell Physiol* 212, 207-214.

- Gharibi, B., Abraham, A.A., Ham, J., Evans, B.A. 2011. Adenosine receptor subtype expression and activation influence the differentiation of mesenchymal stem cells to osteoblasts and adipocytes. *J Bone Miner Res* 26, 2112-2124.
- Gharibi, B., Abraham, A.A., Ham, J., Evans, B.A. 2012. Contrasting effects of A1 and A2b adenosine receptors on adipogenesis. *Int J Obes* 36, 397-406.
- Graff, R.D., Lazarowski, E.R., Banes, A.J., Lee, G.M. 2000. ATP release by mechanically loaded porcine chondrons in pellet culture. *Arthritis Rheum.* 43, 1571-1579.
- Graff, R.D., Picher, M., Lee, G.M. 2003. Extracellular nucleotides, cartilage stress, and calcium crystal formation. *Curr Opin Rheumatol* 15, 315-320.
- Grol, M.W., Zelner, I., Dixon, S.J. 2012. P2X(7)-mediated calcium influx triggers a sustained, PI3K-dependent increase in metabolic acid production by osteoblast-like cells. *Am J Physiol Endocrinol Metab* 302, E561-575.
- Hayton, M.J., Dillon, J.P., Glynn, D., Curran, J.M., Gallagher, J.A., Buckley, K.A. 2005. Involvement of adenosine 5'-triphosphate in ultrasound-induced fracture repair. *Ultrasound Med Biol* 31, 1131-1138.
- Hazama, R., Qu, X., Yokoyama, K., Tanaka, C., Kinoshita, E., He, J., Takahashi, S., Tohyama, K., Yamamura, H., Tohyama, Y. 2009. ATP-induced osteoclast function: the formation of sealing-zone like structure and the secretion of lytic granules via microtubule-deacetylation under the control of Syk. *Genes Cells* 14, 871-884.
- He, W., Cronstein, B.N. 2012. Adenosine A1 receptor regulates osteoclast formation by altering TRAF6/TAK1 signaling. *Purinergic Signal* 8, 327-337.
- Hecht, E., Liedert, A., Ignatius, A., Mizaikoff, B., Kranz, C. 2013. Local detection of mechanically induced ATP release from bone cells with ATP microbiosensors. *Biosens Bioelectron* 44, 27-33.
- Hessle, L., Johnson, K.A., Anderson, H.C., Narisawa, S., Sali, A., Goding, J.W., Terkeltaub, R., Millan, J.L. 2002. Tissue-nonspecific alkaline phosphatase and plasma cell membrane glycoprotein-1 are central antagonistic regulators of bone mineralization. *Proc Natl Acad Sci USA* 99, 9445-9449.
- Hiken, J.F., Steinberg, T.H. 2004. ATP downregulates P2X7 and inhibits osteoclast formation in RAW cells. *Am J Physiol Cell Physiol* 287, C403-412.
- Hoebertz, A., Mahendran, S., Burnstock, G., Arnett, T.R. 2002. ATP and UTP at low concentrations strongly inhibit bone formation by osteoblasts: a novel role for the P2Y₂ receptor in bone remodeling. *J Cell Biochem* 86, 413-419.
- Hoebertz, A., Meghji, S., Burnstock, G., Arnett, T.R. 2001. Extracellular ADP is a powerful osteolytic agent: evidence for signaling through the P2Y₁ receptor on bone cells. *FASEB J* 15, 1139-1148.
- Hoebertz, A., Townsend-Nicholson, A., Glass, R., Burnstock, G., Arnett, T.R. 2000. Expression of P2 receptors in bone and cultured bone cells. *Bone* 27, 503-510.
- Hollopeter, G., Jantzen, H.M., Vincent, D., Li, G., England, L., Ramakrishnan, V., Yang, R.B., Nurden, P., Nurden, A., Julius, D., Conley, P.B. 2001. Identification of the platelet ADP receptor targeted by antithrombotic drugs. *Nature* 409, 202-207.
- Hsu, H.H. 1992. In vitro calcium deposition by rachitic rat matrix vesicles: nucleoside triphosphate supported calcium deposition. *Biochim Biophys Acta* 1116, 227-233.
- Hung, C.T., Allen, F.D., Mansfield, K.D., Shapiro, I.M. 1997. Extracellular ATP modulates [Ca²⁺]_i in retinoic acid-treated embryonic chondrocytes. *Am Journal Physiol* 272, C1611-1617.
- Huo, B., Lu, X.L., Hung, C.T., Costa, K.D., Xu, Q., Whitesides, G.M., Guo, X.E. 2008. Fluid Flow Induced Calcium Response in Bone Cell Network. *Cell Mol Bioeng* 1, 58-66.

- Husted, L.B., Harslof, T., Stenkjaer, L., Carstens, M., Jorgensen, N.R., Langdahl, B.L. 2013. Functional polymorphisms in the P2X7 receptor gene are associated with osteoporosis. *Osteoporos Int* 24, 949-959.
- Ihara, H., Hirukawa, K., Goto, S., Togari, A. 2005. ATP-stimulated interleukin-6 synthesis through P2Y receptors on human osteoblasts. *Biochem Biophys Res Commun* 326, 329-334.
- Jande, S.S., Belanger, L.F. 1973. The life cycle of the osteocyte. *Clin Orthop Relat Res*, 281-305.
- Janssens, R., Communi, D., Piroton, S., Samson, M., Parmentier, M., Boeynaems, J.M. 1996. Cloning and tissue distribution of the human P2Y₁ receptor. *Biochem Biophys Res Commun* 221, 588-593.
- Janssens, R., Painsavoine, P., Parmentier, M., Boeynaems, J.M. 1999. Human P2Y₂ receptor polymorphism: identification and pharmacological characterization of two allelic variants. *Br J Pharmacol* 127, 709-716.
- Johnson, K.A., Hesse, L., Vaingankar, S., Wennberg, C., Mauro, S., Narisawa, S., Goding, J.W., Sano, K., Millan, J.L., Terkeltaub, R. 2000. Osteoblast tissue-nonspecific alkaline phosphatase antagonizes and regulates PC-1. *Am J Physiol Regul Integr Comp Physiol* 279, R1365-R1377.
- Jones, S.J., Gray, C., Boyde, A., Burnstock, G. 1997. Purinergic transmitters inhibit bone formation by cultured osteoblasts. *Bone* 21, 393-399.
- Jorgensen, N.R., Grove, E.L., Schwarz, P., Vestergaard, P. 2012. Clopidogrel and the risk of osteoporotic fractures: a nationwide cohort study. *J Intern Med* 272, 385-393.
- Jorgensen, N.R., Henriksen, Z., Brot, C., Eriksen, E.F., Sorensen, O.H., Civitelli, R., Steinberg, T.H. 2000. Human osteoblastic cells propagate intercellular calcium signals by two different mechanisms. *J Bone Miner Res* 15, 1024-1032.
- Jorgensen, N.R., Henriksen, Z., Sorensen, O.H., Eriksen, E.F., Civitelli, R., Steinberg, T.H. 2002. Intercellular calcium signaling occurs between human osteoblasts and osteoclasts and requires activation of osteoclast P2X7 receptors. *J Biol Chem* 277, 7574-7580.
- Kanabe, S., Hsu, H.H., Cecil, R.N., Anderson, H.C. 1983. Electron microscopic localization of adenosine triphosphate (ATP)-hydrolyzing activity in isolated matrix vesicles and reconstituted vesicles from calf cartilage. *J Histochem Cytochem* 31, 462-470.
- Kaplan, A.D., Kilkenny, D.M., Hill, D.J., Dixon, S.J. 1996. Extracellular nucleotides act through P2U purinoceptors to elevate [Ca²⁺]_i and enhance basic fibroblast growth factor-induced proliferation in sheep chondrocytes. *Endocrinology* 137, 4757-4766.
- Kaplan, A.D., Reimer, W.J., Feldman, R.D., Dixon, S.J. 1995. Extracellular nucleotides potentiate the cytosolic Ca²⁺, but not cyclic adenosine 3', 5'-monophosphate response to parathyroid hormone in rat osteoblastic cells. *Endocrinology* 136, 1674-1685.
- Kara, F.M., Chitu, V., Sloane, J., Axelrod, M., Fredholm, B.B., Stanley, E.R., Cronstein, B.N. 2010a. Adenosine A1 receptors (A1Rs) play a critical role in osteoclast formation and function. *FASEB J* 24, 2325-2333.
- Kara, F.M., Doty, S.B., Boskey, A., Goldring, S., Zaidi, M., Fredholm, B.B., Cronstein, B.N. 2010b. Adenosine A(1) receptors regulate bone resorption in mice: adenosine A(1) receptor blockade or deletion increases bone density and prevents ovariectomy-induced bone loss in adenosine A(1) receptor-knockout mice. *Arthritis Rheum* 62, 534-541.
- Katz, S., Boland, R., Santillan, G. 2006. Modulation of ERK 1/2 and p38 MAPK signaling pathways by ATP in osteoblasts: involvement of mechanical stress-activated calcium influx, PKC and Src activation. *Int J Biochem Cell Biol* 38, 2082-2091.

Katz, S., Boland, R., Santillan, G. 2008. Purinergic (ATP) signaling stimulates JNK1 but not JNK2 MAPK in osteoblast-like cells: contribution of intracellular Ca²⁺ release, stress activated and L-voltage-dependent calcium influx, PKC and Src kinases. *Arch Biochem Biophys* 477, 244-252.

Ke, H.Z., Qi, H., Weidema, A.F., Zhang, Q., Panupinthu, N., Crawford, D.T., Grasser, W.A., Paralkar, V.M., Li, M., Audoly, L.P., Gabel, C.A., Jee, W.S., Dixon, S.J., Sims, S.M., Thompson, D.D. 2003. Deletion of the P2X7 nucleotide receptor reveals its regulatory roles in bone formation and resorption. *Mol Endocrinol* 17, 1356-1367.

Keresztes, M., Haggblad, J., Heilbronn, E. 1991. Basal and ATP-stimulated phosphoinositol metabolism in fusing rat skeletal muscle cells in culture. *Exp Cell Res* 196, 362-364.

Knight, M.M., McGlashan, S.R., Garcia, M., Jensen, C.G., Poole, C.A. 2009. Articular chondrocytes express connexin 43 hemichannels and P2 receptors - a putative mechanoreceptor complex involving the primary cilium? *J Anat* 214, 275-283.

Kono, T., Nishikori, T., Kataoka, H., Uchio, Y., Ochi, M., Enomoto, K. 2006. Spontaneous oscillation and mechanically induced calcium waves in chondrocytes. *Cell Biochem Func* 24, 103-111.

Koolpe, M., Benton, H.P. 1997. Calcium-mobilizing purine receptors on the surface of mammalian articular chondrocytes. *J Orthop Res* 15, 204-212.

Koolpe, M., Pearson, D., Benton, H.P. 1999. Expression of both P1 and P2 purine receptor genes by human articular chondrocytes and profile of ligand-mediated prostaglandin E2 release. *Arthritis Rheum* 42, 258-267.

Korcok, J., Raimundo, L.N., Du, X., Sims, S.M., Dixon, S.J. 2005. P2Y₆ Nucleotide Receptors Activate NF- κ B and Increase Survival of Osteoclasts. *J Biol Chem* 280, 16909-16915.

Korcok, J., Raimundo, L.N., Ke, H.Z., Sims, S.M., Dixon, S.J. 2004. Extracellular nucleotides act through P2X7 receptors to activate NF- κ B in osteoclasts. *J Bone Miner Res* 19, 642-651.

Kringelbach, T.M., Aslan, D., Novak, I., Schwarz, P., Jorgensen, N.R. 2014. UTP-induced ATP release is a fine-tuned signalling pathway in osteocytes. *Purinergic Signal* 10, 337-347.

Kudirka, J.C., Panupinthu, N., Tesseyman, M.A., Dixon, S.J., Bernier, S.M. 2007. P2Y nucleotide receptor signaling through MAPK/ERK is regulated by extracellular matrix: involvement of beta3 integrins. *J Cell Physiol* 213, 54-64.

Kumagai, H., Sacktor, B., Filburn, C.R. 1991. Purinergic regulation of cytosolic calcium and phosphoinositide metabolism in rat osteoblast-like osteosarcoma cells. *J Bone Miner Res* 6, 697-708.

Kumagai, H., Sakamoto, H., Guggino, S., Filburn, C.R., Sacktor, B. 1989. Neurotransmitter regulation of cytosolic calcium in osteoblast-like bone cells. *Calcif Tissue Int* 45, 251-254.

Leong, W.S., Russell, R.G., Caswell, A.M. 1990. Extracellular ATP stimulates resorption of bovine nasal cartilage. *Biochem Soc Trans* 18, 951-952.

Leong, W.S., Russell, R.G., Caswell, A.M. 1993. Induction of enhanced responsiveness of human articular chondrocytes to extracellular ATP by tumour necrosis factor-alpha. *Clin Sci* 85, 569-575.

Lerner, U., Fredholm, B.B. 1983. Studies on the mechanisms by which 2-chloroadenosine stimulates bone resorption in tissue culture. *Biochim Biophys Acta* 757, 226-234.

Lerner, U.H., Sahlberg, K., Fredholm, B.B. 1987. Characterization of adenosine receptors in bone. Studies on the effect of adenosine analogues on cyclic AMP formation and bone resorption in cultured mouse calvaria. *Acta Physiol Scand*. 131, 287-296.

Li, J., Liu, D., Ke, H.Z., Duncan, R.L., Turner, C.H. 2005. The P2X7 nucleotide receptor mediates skeletal mechanotransduction. *J Biol Chem* 280, 42952-42959.

- Li, J., Meyer, R., Duncan, R.L., Turner, C.H. 2009. P2X7 nucleotide receptor plays an important role in callus remodeling during fracture repair. *Calcif Tissue Int* 84, 405-412.
- Liu, D., Genetos, D.C., Shao, Y., Geist, D.J., Li, J., Ke, H.Z., Turner, C.H., Duncan, R.L. 2008. Activation of extracellular-signal regulated kinase (ERK1/2) by fluid shear is Ca⁽²⁺⁾- and ATP-dependent in MC3T3-E1 osteoblasts. *Bone* 42, 644-652.
- Mackenzie, N.C., Zhu, D., Milne, E.M., van 't, H.R., Martin, A., Darryl, Q.L., Millan, J.L., Farquharson, C., Macrae, V.E. 2012. Altered bone development and an increase in FGF-23 expression in Enpp1(-/-) mice. *PLoS ONE* 7, e32177.
- Martinello, T., Baldoin, M.C., Morbiato, L., Paganin, M., Tarricone, E., Schiavo, G., Bianchini, E., Sandona, D., Betto, R. 2011. Extracellular ATP signaling during differentiation of C2C12 skeletal muscle cells: role in proliferation. *Mol Cell Biochem* 351, 183-196.
- Mediero, A., Frenkel, S.R., Wilder, T., He, W., Mazumder, A., Cronstein, B.N. 2012a. Adenosine A2A receptor activation prevents wear particle-induced osteolysis. *Sci Transl Med* 4, 135ra165.
- Mediero, A., Kara, F.M., Wilder, T., Cronstein, B.N. 2012b. Adenosine A(2A) receptor ligation inhibits osteoclast formation. *Am J Pathol* 180, 775-786.
- Meyer, M.P., Swann, K., Burnstock, G., Clarke, J.D. 2001. The extracellular ATP receptor, cP2Y₁, inhibits cartilage formation in micromass cultures of chick limb mesenchyme. *Dev Dyn* 222, 494-505.
- Millward-Sadler, S.J., Wright, M.O., Flatman, P.W., Salter, D.M. 2004. ATP in the mechanotransduction pathway of normal human chondrocytes. *Biorheology* 41, 567-575.
- Morrison, M.S., Turin, L., King, B.F., Burnstock, G., Arnett, T.R. 1998. ATP is a potent stimulator of the activation and formation of rodent osteoclasts. *J Physiol* 511 (Pt 2), 495-500.
- Mortensen, S.P., Gonzalez-Alonso, J., Nielsen, J.J., Saltin, B., Hellsten, Y. 2009. Muscle interstitial ATP and norepinephrine concentrations in the human leg during exercise and ATP infusion. *J Appl Physiol* 107, 1757-1762.
- Naemsch, L.N., Dixon, S.J., Sims, S.M. 2001. Activity-dependent development of P2X7 current and Ca²⁺ entry in rabbit osteoclasts. *J Biol Chem* 276, 39107-39114.
- Naemsch, L.N., Weidema, A.F., Sims, S.M., Underhill, T.M., Dixon, S.J. 1999. P2X4 purinoceptors mediate an ATP-activated, non-selective cation current in rabbit osteoclasts. *J Cell Sci* 112 (Pt 23), 4425-4435.
- Nakamura, E., Uezono, Y., Narusawa, K., Shibuya, I., Oishi, Y., Tanaka, M., Yanagihara, N., Nakamura, T., Izumi, F. 2000. ATP activates DNA synthesis by acting on P2X receptors in human osteoblast-like MG-63 cells. *Am J Physiol Cell Physiol* 279, C510-C519.
- Nicke, A., Kuan, Y.H., Masin, M., Rettinger, J., Marquez-Klaka, B., Bender, O., Gorecki, D.C., Murrell-Lagnado, R.D., Soto, F. 2009. A functional P2X7 splice variant with an alternative transmembrane domain 1 escapes gene inactivation in P2X7 knock-out mice. *J Biol Chem* 284, 25813-25822.
- Nishii, N., Nejime, N., Yamauchi, C., Yanai, N., Shinozuka, K., Nakabayashi, T. 2009. Effects of ATP on the intracellular calcium level in the osteoblastic TBR31-2 cell line. *Biol Pharm Bull* 32, 18-23.
- Noronha-Matos, J.B., Costa, M.A., Magalhaes-Cardoso, M.T., Ferreira, F., Pelletier, J., Freitas, R., Neves, J.M., Sevigny, J., Correia-de-Sa, P. 2012. Role of ecto-NTPDases on UDP-sensitive P2Y₆ receptor activation during osteogenic differentiation of primary bone marrow stromal cells from postmenopausal women. *J Cell Physiol* 227, 2694-2709.
- Ohlendorff, S.D., Tofteng, C.L., Jensen, J.E., Petersen, S., Civitelli, R., Fenger, M., Abrahamsen, B., Hermann, A.P., Eiken, P., Jorgensen, N.R. 2007. Single nucleotide polymorphisms in the P2X7 gene

are associated to fracture risk and to effect of estrogen treatment. *Pharmacogenet Genomics* 17, 555-567.

Orriss, I., Syberg, S., Wang, N., Robaye, B., Gartland, A., Jorgensen, N., Arnett, T., Boeynaems, J.M. 2011a. Bone phenotypes of P2 receptor knockout mice. *Front Biosci (Schol Ed)* 3, 1038-1046.

Orriss, I.R., Arnett, T.R. 2010. Extracellular nucleotides regulate the expression and activity of ecto-nucleotidases by osteoblasts, *Bone*, Vol. 46 S1. pp. S55-S55.

Orriss, I.R., Evans, H.R., Gartland, A., Arnett, T.R. 2008. Microct analysis of P2Y₁ and P2Y₂ receptor knockout mice demonstrates significant changes in bone phenotype. *Calcif Tissue Int* 83, 2-3.

Orriss, I.R., Key, M.L., Brandao-Burch, A., Patel, J.J., Burnstock, G., Arnett, T.R. 2012. The regulation of osteoblast function and bone mineralisation by extracellular nucleotides: The role of P2X receptors. *Bone* 51, 389-400.

Orriss, I.R., Key, M.L., Hajjawi, M.O., Arnett, T.R. 2013. Extracellular ATP released by osteoblasts is a key local inhibitor of bone mineralisation. *PLoS one* 8, e69057.

Orriss, I.R., Knight, G.E., Ranasinghe, S., Burnstock, G., Arnett, T.R. 2006. Osteoblast responses to nucleotides increase during differentiation. *Bone* 39, 300-309.

Orriss, I.R., Knight, G.E., Utting, J.C., Taylor, S.E., Burnstock, G., Arnett, T.R. 2009. Hypoxia stimulates vesicular ATP release from rat osteoblasts. *J Cell Physiol* 220, 155-162.

Orriss, I.R., Utting, J.C., Brandao-Burch, A., Colston, K., Grubb, B.R., Burnstock, G., Arnett, T.R. 2007. Extracellular nucleotides block bone mineralization in vitro: evidence for dual inhibitory mechanisms involving both P2Y₂ receptors and pyrophosphate. *Endocrinology* 148, 4208-4216.

Orriss, I.R., Wang, N., Burnstock, G., Arnett, T.R., Gartland, A., Robaye, B., Boeynaems, J.M. 2011b. The P2Y₆ receptor stimulates bone resorption by osteoclasts. *Endocrinology* 152, 3706-3716.

Panupinthu, N., Rogers, J.T., Zhao, L., Solano-Flores, L.P., Possmayer, F., Sims, S.M., Dixon, S.J. 2008. P2X7 receptors on osteoblasts couple to production of lysophosphatidic acid: a signaling axis promoting osteogenesis. *J Cell Biol* 181, 859-871.

Panupinthu, N., Zhao, L., Possmayer, F., Ke, H.Z., Sims, S.M., Dixon, S.J. 2007. P2X7 nucleotide receptors mediate blebbing in osteoblasts through a pathway involving lysophosphatidic acid. *J Biol Chem* 282, 3403-3412.

Pellegatti, P., Falzoni, S., Donvito, G., Lemaire, I., Di Virgilio, F. 2011. P2X7 receptor drives osteoclast fusion by increasing the extracellular adenosine concentration. *FASEB J.* 25, 1264-1274.

Penolazzi, L., Bianchini, E., Lambertini, E., Baraldi, P.G., Romagnoli, R., Piva, R., Gambari, R. 2005. N-Arylpiperazine modified analogues of the P2X7 receptor KN-62 antagonist are potent inducers of apoptosis of human primary osteoclasts. *J Biomed Sci* 12, 1013-1020.

Pines, A., Bivi, N., Romanello, M., Damante, G., Kelley, M.R., Adamson, E.D., D'andrea, P., Quadrioglio, F., Moro, L., Tell, G. 2005. Cross-regulation between Egr-1 and APE/Ref-1 during early response to oxidative stress in the human osteoblastic HOBIT cell line: evidence for an autoregulatory loop. *Free Radic Res* 39, 269-281.

Ralevic, V., Burnstock, G. 1998. Receptors for purines and pyrimidines. *Pharmacol Rev* 50, 413-492.

Reimer, W.J., Dixon, S.J. 1992. Extracellular nucleotides elevate [Ca²⁺]_i in rat osteoblastic cells by interaction with two receptor subtypes. *Am J Physiol* 263, C1040-C1048.

Riddle, R.C., Taylor, A.F., Rogers, J.R., Donahue, H.J. 2007. ATP release mediates fluid flow-induced proliferation of human bone marrow stromal cells. *J Bone Miner Res* 22, 589-600.

Romanello, M., Codognotto, A., Bicego, M., Pines, A., Tell, G., D'andrea, P. 2005. Autocrine/paracrine stimulation of purinergic receptors in osteoblasts: Contribution of vesicular ATP release. *Biochem Biophys Res Commun* 331, 1429-1438.

Romanello, M., Pani, B., Bicego, M., D'andrea, P. 2001. Mechanically induced ATP release from human osteoblastic cells. *Biochem Biophys Res Commun*. 289, 1275-1281.

Rosenthal, A.K., Hempel, D., Kurup, I.V., Masuda, I., Ryan, L.M. 2010. Purine receptors modulate chondrocyte extracellular inorganic pyrophosphate production. *Osteoarthritis Cartilage* 18, 1496-1501.

Rumney, R.M., Sunters, A., Reilly, G.C., Gartland, A. 2012. Application of multiple forms of mechanical loading to human osteoblasts reveals increased ATP release in response to fluid flow in 3D cultures and differential regulation of immediate early genes. *J Biomech* 45, 549-554.

Ryan, L.M., Kurup, I.V., Derfus, B.A., Kushnaryov, V.M. 1992. ATP-induced chondrocalcinosis. *Arthritis Rheum* 35, 1520-1525.

Ryten, M., Dunn, P.M., Neary, J.T., Burnstock, G. 2002. ATP regulates the differentiation of mammalian skeletal muscle by activation of a P2X5 receptor on satellite cells. *J Cell Biol* 158, 345-355.

Schoenberg, M. 1989. Effect of adenosine triphosphate analogues on skeletal muscle fibers in rigor. *Biophys J* 56, 33-41.

Sciancalepore, M., Luin, E., Parato, G., Ren, E., Giniatullin, R., Fabbretti, E., Lorenzon, P. 2012. Reactive oxygen species contribute to the promotion of the ATP-mediated proliferation of mouse skeletal myoblasts. *Free Radical Biol Med* 53, 1392-1398.

Shimegi, S. 1996. ATP and adenosine act as a mitogen for osteoblast-like cells (MC3T3-E1). *Calcif Tissue Int* 58, 109-113.

Sistare, F.D., Rosenzweig, B.A., Contrera, J.G. 1995. P2 purinergic receptors potentiate parathyroid hormone receptor-mediated increases in intracellular calcium and inositol trisphosphate in UMR-106 rat osteoblasts. *Endocrinology* 136, 4489-4497.

Su, X., Floyd, D.H., Hughes, A., Xiang, J., Schneider, J.G., Uluckan, O., Heller, E., Deng, H., Zou, W., Craft, C.S., Wu, K., Hirbe, A.C., Grabowska, D., Egleton, M.C., Townsley, S., Collins, L., Piwnicka-Worms, D., Steinberg, T.H., Novack, D.V., Conley, P.B., Hurchla, M.A., Rogers, M., Weilbaecher, K.N. 2012. The ADP receptor P2RY₁₂ regulates osteoclast function and pathologic bone remodeling. *J Clin Invest* 122, 3579-3592.

Suzuki, A., Kotoyori, J., Oiso, Y., Kozawa, O. 1993. Prostaglandin E2 is a potential mediator of extracellular ATP action in osteoblast-like cells. *Cell Adhes Commun* 1, 113-118.

Syberg, S., Brandao-Burch, A., Patel, J.J., Hajjawi, M., Arnett, T.R., Schwarz, P., Jorgensen, N.R., Orriss, I.R. 2012a. Clopidogrel (Plavix(R)), a P2Y₁₂ receptor antagonist, inhibits bone cell function in vitro and decreases trabecular bone in vivo. *J Bone Miner Res* 27, 2373-2386.

Syberg, S., Petersen, S., Beck Jensen, J.E., Gartland, A., Teilmann, J., Chessell, I., Steinberg, T.H., Schwarz, P., Jorgensen, N.R. 2012b. Genetic background strongly influences the bone phenotype of P2X7 receptor knockout mice. *J Osteoporos* 2012, 391097.

Takedachi, M., Oohara, H., Smith, B.J., Iyama, M., Kobashi, M., Maeda, K., Long, C.L., Humphrey, M.B., Stoecker, B.J., Toyosawa, S., Thompson, L.F., Murakami, S. 2012. CD73-generated adenosine promotes osteoblast differentiation. *J Cell Physiol* 227, 2622-2631.

Tesch, A.M., MacDonald, M.H., Kollias-Baker, C., Benton, H.P. 2002. Chondrocytes respond to adenosine via A(2)receptors and activity is potentiated by an adenosine deaminase inhibitor and a phosphodiesterase inhibitor. *Osteoarthritis Cartilage* 10, 34-43.

- Thompson, W.R., Majid, A.S., Czymmek, K.J., Ruff, A.L., Garcia, J., Duncan, R.L., Farach-Carson, M.C. 2011. Association of the $\alpha(2)\delta(1)$ subunit with $Ca(v)3.2$ enhances membrane expression and regulates mechanically induced ATP release in MLO-Y4 osteocytes. *J Bone Miner Res* 26, 2125-2139.
- Tu, J., Lu, L., Cai, W., Ballard, H.J. 2012. cAMP/protein kinase A activates cystic fibrosis transmembrane conductance regulator for ATP release from rat skeletal muscle during low pH or contractions. *PloS one* 7, e50157.
- Varani, K., De, M.M., Vincenzi, F., Gessi, S., Merighi, S., Pellati, A., Ongaro, A., Caruso, A., Cadossi, R., Borea, P.A. 2008a. Characterization of adenosine receptors in bovine chondrocytes and fibroblast-like synoviocytes exposed to low frequency low energy pulsed electromagnetic fields. *Osteoarthritis Cartilage* 16, 292-304.
- Varani, K., De, M.M., Vincenzi, F., Tosi, A., Gessi, S., Merighi, S., Pellati, A., Masieri, F., Ongaro, A., Borea, P.A. 2008b. Pharmacological characterization of P2X1 and P2X3 purinergic receptors in bovine chondrocytes. *Osteoarthritis Cartilage* 16, 1421-1429.
- Vincenzi, F., Targa, M., Corciulo, C., Gessi, S., Merighi, S., Setti, S., Cadossi, R., Goldring, M.B., Borea, P.A., Varani, K. 2013. Pulsed electromagnetic fields increased the anti-inflammatory effect of A(2)A and A(3) adenosine receptors in human T/C-28a2 chondrocytes and hFOB 1.19 osteoblasts. *PloS one* 8, e65561.
- Voss, A.A. 2009. Extracellular ATP inhibits chloride channels in mature mammalian skeletal muscle by activating P2Y₁ receptors. *J Physiol* 587, 5739-5752.
- Wang, N., Robaye, B., Agrawal, A., Skerry, T.M., Boeynaems, J.M., Gartland, A. 2012. Reduced bone turnover in mice lacking the P2Y₁₃ receptor of ADP. *Mol Endocrinol* 26, 142-152.
- Wang, N., Robaye, B., Gossiel, F., Boeynaems, J.M., Gartland, A. 2014. The P2Y₁₃ receptor regulates phosphate metabolism and FGF-23 secretion with effects on skeletal development. *FASEB J* 28, 2249-2259.
- Wang, N., Rumney, R.M., Yang, L., Robaye, B., Boeynaems, J.M., Skerry, T.M., Gartland, A. 2013. The P2Y₁₃ receptor regulates extracellular ATP metabolism and the osteogenic response to mechanical loading. *J Bone Miner Res* 28, 1446-1456.
- Weidema, A.F., Barbera, J., Dixon, S.J., Sims, S.M. 1997. Extracellular nucleotides activate non-selective cation and Ca^{2+} -dependent K^+ channels in rat osteoclasts. *J Physiol* 503 (Pt 2), 303-315.
- Weidema, A.F., Dixon, S.J., Sims, S.M. 2001. Activation of P2Y but not P2X(4) nucleotide receptors causes elevation of $[Ca^{2+}]_i$ in mammalian osteoclasts. *Am J Physiol Cell Physiol* 280, C1531-C1539.
- Wesselius, A., Bours, M.J., Agrawal, A., Gartland, A., Dagnelie, P.C., Schwarz, P., Jorgensen, N.R. 2012. Role of purinergic receptor polymorphisms in human bone. *Front Biosci* 17, 2572-2585.
- Wesselius, A., Bours, M.J., Henriksen, Z., Syberg, S., Petersen, S., Schwarz, P., Jorgensen, N.R., van Helden, S., Dagnelie, P.C. 2013a. Association of P2Y₂ receptor SNPs with bone mineral density and osteoporosis risk in a cohort of Dutch fracture patients. *Purinergic Signal* 9, 41-49.
- Wesselius, A., Bours, M.J., Jorgensen, N.R., Wiley, J., Gu, B., van Helden, S., van Rhijn, L., Dagnelie, P.C. 2013b. Non-synonymous polymorphisms in the P2RX4 are related to bone mineral density and osteoporosis risk in a cohort of Dutch fracture patients. *Purinergic Signal* 9, 123-130.
- Wiebe, S.H., Sims, S.M., Dixon, S.J. 1999. Calcium signalling via multiple P2 purinoceptor subtypes in rat osteoclasts. *Cellular Physiol Biochem* 9, 323-337.
- Wildman, S.S., King, B.F., Burnstock, G. 1998. Zn^{2+} modulation of ATP-responses at recombinant P2X2 receptors and its dependence on extracellular pH. *Br J Pharmacol* 123, 1214-1220.

You, J., Jacobs, C.R., Steinberg, T.H., Donahue, H.J. 2002. P2Y purinoceptors are responsible for oscillatory fluid flow-induced intracellular calcium mobilization in osteoblastic cells. *J Biol Chem* 277, 48724-48729.

Yu, H., Ferrier, J. 1993a. ATP induces an intracellular calcium pulse in osteoclasts. *Biochem Biophys Res Commun* 191, 357-363.

Yu, H., Ferrier, J. 1993b. Osteoblast-like cells have a variable mixed population of purino/nucleotide receptors. *FEBS Lett.* 328, 209-214.

Zimmermann, H., Zebisch, M., Strater, N. 2012. Cellular function and molecular structure of ecto-nucleotidases. *Purinergic Signal* 8, 437-502.

Zippel, N., Limbach, C.A., Ratajski, N., Urban, C., Luparello, C., Pansky, A., Kassack, M.U., Tobiasch, E. 2012. Purinergic receptors influence the differentiation of human mesenchymal stem cells. *Stem Cells Dev* 21, 884-900.

Figure 1




Figure 2




Figure 3

