


Contents lists available at [ScienceDirect](#)

Animal

The international journal of animal biosciences



Dairy vs beef production – expert views on welfare of cattle in common food production systems

Roi Mandel ^{a,b,*}, Marc B.M. Bracke ^{c,1}, Christine J. Nicol ^{d,1}, John A. Webster ^e, Lorenz Gyax ^f

^a Section of Animal Welfare and Disease Control, Department of Veterinary and Animal Sciences, University of Copenhagen, 1870 Frederiksberg C, Denmark

^b Behavioural Ecology Group, Section for Ecology & Evolution, Department of Biology, University of Copenhagen, 2100 Copenhagen, Denmark

^c Wageningen Livestock Research, Wageningen University and Research, 6708 WD Wageningen, The Netherlands

^d Royal Veterinary College, AL9 7TA Hatfield, United Kingdom

^e Professor Emeritus at the University of Bristol and Former Head of the Bristol Vet School, BS40 5DU Langford, United Kingdom

^f Animal Husbandry & Ethology, Albrecht Daniel Thaer-Institute of Agricultural and Horticultural Sciences, Humboldt-University of Berlin, 10099 Berlin, Germany

ARTICLE INFO

Article history:

Received 29 January 2022

Revised 18 July 2022

Accepted 22 July 2022

Available online xxxxx

Keywords:

Animal welfare

Expert opinion

Meat

Milk

Welfare risk

ABSTRACT

Consumers' views and concerns about the welfare of farm animals may play an important role in their decision to consume dairy, meat and/or plants as their primary protein source. As animals are killed prematurely in both dairy and beef industries, it is important to quantify and compare welfare compromises in these two sectors before the point of death. Seventy world-leading bovine welfare experts based in 23 countries were asked to evaluate the likelihood of a bovine to experience 12 states of potential welfare concern, inspired by the Welfare Quality[®] protocol. The evaluation focused on the most common beef and dairy production systems in the experts' country and was carried out separately for dairy/beef calves raised for red meat, dairy/beef calves raised for veal, dairy/beef calves raised as a replacement, and for dairy/beef cows. The results show experts rated the overall likelihood of a negative welfare state (i.e. welfare risk) to be higher in animals from dairy herds than from beef herds, for all animal categories, regardless of whether they were used to produce milk, red meat or veal. These findings suggest that consuming food products derived from common dairy production systems (dairy or meat) may be more harmful to the welfare of animals than consuming products derived from common beef production systems (i.e. from animals solely raised for their meat). Raising awareness about the linkage between dairy and meat production, and the toll of milk production on the welfare state of animals in the dairy industry, may encourage a more sustainable and responsible food consumption.

© 2022 The Authors. Published by Elsevier B.V. on behalf of The Animal Consortium. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Implications

Abstaining from consuming meat, but not dairy products, is regarded by many to be a sign of compassion towards animals. Here, we show that food products derived from the most common dairy production systems (dairy or meat) are rated by cattle welfare experts as more harmful to the welfare of animals than consuming products derived from the most common beef production systems. With a projected 20% increase in global milk production by 2029, the highest among all livestock commodities, the toll of milk production on animals in the dairy industry may require a

thorough revision of our current societal, political and moral decisions.

Introduction

Abstaining from consuming meat, but not dairy products, is regarded by many to be a sign of compassion towards animals (Fox and Ward, 2008; Corrin and Papadopoulos, 2017). Vegans go a step further and reject the consumption of any animal-based product, but few, if any, advocate abstention from dairy while continuing to consume meat. The extent to which an independent analysis of animal welfare aligns with such consumer choices is unknown. Importantly, there are substantial reasons to predict a discrepancy, particularly when considering the role of the dairy industry in producing meat (Fig. 1A), and the higher degree of intervention in the lives of animals that are used “for more than their meat” (i.e. intervention associated with daily milking of the

* Corresponding author at: Section of Animal Welfare and Disease Control, Department of Veterinary and Animal Sciences, University of Copenhagen, 1870 Frederiksberg C, Denmark.

¹ Contributed equally.

<https://doi.org/10.1016/j.animal.2022.100622>

1751-7311/© 2022 The Authors. Published by Elsevier B.V. on behalf of The Animal Consortium.

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).




Fig. 1. (A) Diagram showing the origin (dairy/beef herd) and production goals, i.e. the general flow of calves from dairy herds and beef herds to red meat, veal and cow replacements [inspired by European Food Safety Authority Panel on Animal Health and Welfare (EFSA, 2012a; 2012b)]. Dashed arrows reflect production routes that are not common to all farms. (B) First principal component reflecting the overall welfare risk (likelihood of 12 welfare concerns, see text) as assessed by the experts as a function of the different origins and animal production goals. N = Number of experts that reported an assessment on all 12 areas of welfare concern. Boxplots show minimum, lower quartile, median, upper quartile and maximum values. Black lines: model estimates with 95% upper and lower confidence intervals.

dams, and the consequent management of their calves). Here, we obtained an independent assessment of the likelihood that dairy cattle and beef cattle would experience negative welfare, by surveying a panel of leading cattle welfare experts, each focusing on the most common beef and dairy production systems in their country of expertise. The comparison was carried out between dairy and beef calves when raised directly for their meat (i.e. dairy and beef calves raised for red meat and for veal, consisting mostly of male calves), and between dairy and beef cows raised for producing calves/milk.

Expert assessments can help to characterise uncertainty and fill data gaps where traditional scientific research is not possible or data are not yet accessible or available (EFSA, 2012a; 2012b). Such assessments have been widely used to review the impacts of housing, management or other anthropogenic challenges to domestic [e.g. cattle (Bertocchi et al., 2018); broilers (Bracke et al., 2019) and canine and felines (Dawson et al., 2016)] and wild animals (e.g. Nicol et al., 2020), using frameworks such as the Five Domains (e.g. McGreevy et al., 2018; Nicol et al., 2020), and more recently, in relation to the United Nations Sustainable Development Goals (United Nations, 2015; Keeling et al., 2019). The European Food and Safety Authority (EFSA) commonly uses expert opinion to inform debates around welfare topics, such as the use of perches for laying hens (EFSA, 2015), and for assessing welfare risks, such as those that relate to the farming of sheep for wool, meat and milk production (EFSA, 2014).

In the next section, we compare experts' assessment of the overall likelihood that bovines will experience a negative welfare state when raised in the most common housing systems in the dairy and beef sectors, with a focus on Europe and North America. We predicted that the higher degree of intervention in the lives of dairy cattle, which stems from the fact that they are used for "more than their meat" (i.e. milking them, which affects their management and the management of their calves), would result in dairy cattle being rated by experts as being at a higher welfare risk than beef cattle.

Material and methods

Data collection

Overall, 130 cattle welfare experts were invited to participate in our survey based on their 1. Number of publications on the topic of bovine welfare (peer-reviewed research manuscripts and review articles) that appear in the Web of Science database, when using the following keywords: "Welfare" + "bovine"/"cattle"/"dairy cow"/"beef cow") (selected experts were those with the highest number of publications) and 2. H-index (minimum of 10). In addition, we invited further researchers that were recommended to us by the selected experts. Overall, 83 experts agreed to participate in the survey, out of which 13 were omitted from the final analysis for the following reasons: 10 experts did not complete the survey, two experts felt that they could not provide accurate assessments, and one expert felt that his degree of expertise was not sufficient. Data were collected through an online survey built in Qualtrics (Qualtrics XM Platform™, Provo, Utah, USA), between October and December 2020.

The survey included four parts: (1) Consent including a short description of the aim of the experiment "to compare the welfare of cattle across food production systems". (2) General instructions. (3) Characterisation of the common production systems, likelihood ratings and confidence ratings. (4) Characterisation of experts (as summarised in [Supplementary Table S1](#)).

Characterisation of the common production systems

Before providing likelihood ratings, the experts were asked to describe the most common beef and dairy production systems in their country of expertise. The characterisation applied to each animal category separately, using 3–5 fixed criteria ([Supplementary Fig. S1](#); [Supplementary Table S2](#)), to which the experts could answer: "yes", "partial/part of the time", "no", or "I don't know".

Likelihood rating

The experts were asked to rate the likelihood of 12 statements on a scale of 1 (very low) to 7 (very high), and to notice that the statements were built in such a way that the higher the likelihood ratings are, the higher is the welfare risk for the animal (for illustration, see [Supplementary Fig. S1](#), for the exact questions and ratings, see [Table 1](#)). The 12 short welfare statements were inspired by the Welfare Quality protocol (Welfare Quality® 2009), a well-established protocol for assessing bovine welfare. The statements addressed the following core areas of potential welfare concern: (1) inadequate diet, (2) inadequate water supply, (3) thermal discomfort, (4) resting discomfort, (5) injuries, (6) disease, (7) pain resulting from management/handling/surgical procedures, (8) inability to move freely, (9) inability to perform social behaviour, (10) inability to perform other normal behaviours, (11) experiencing negative affective states, and (12) lack of experiencing positive affective states (for the order of presentation, see [Table 1](#)). Since management practices were expected to vary between countries for both beef and dairy sectors (for red meat, veal, replacement, and cows), animal categories were defined as follows: For red meat calves and veal, the evaluation period was from birth to slaughter (or up to 18 months of age). For replacement calves, it was from birth to first calving. For cows, it was from first calving to slaughter. In all cases, the evaluation did not include transportation to slaughter or the process of slaughter itself. To avoid order effects (where rating one animal origin first (e.g. beef) would affect the rating of the other (e.g. dairy), we used four complementary versions of the survey. In all four versions, the presentation of animal category was fixed: Red meat, replacement calves, cows and veal. However, the order of presenting the animal origin (beef/dairy) was counterbalanced (left/right * text colour blue/orange). In each of the four versions, the text colour and order of presentation in [Fig. 1A](#) were matched accordingly.

Confidence rating

The experts were asked to state their level of confidence in their likelihood ratings on a 5-point scale (1 star = low confidence to 5 stars = high confidence). Confidence ratings were reported separately for each of the animal origins in the four animal categories ([Supplementary Fig. S2](#), and [Table 1](#)).

Data processing

We omitted all answers from respondents who had provided an incomplete set of answers for a given combination of animal origin and category. This left the answers of 70 experts (out of 83). In total 1, 3, 34, 25, and seven experts gave complete answers for 3, 4, 6, 7, and 8 of the eight possible combinations of origin and animal category, respectively.

Calculating normally distributed values based on the likelihood ratings

The likelihood ratings were reported on a seven-point Likert-scale. A Likert-scale has a fixed lower and upper limit and is therefore comparable to a proportion. Accordingly, it can be expected that a logit transformation will lead to normally distributed data points. Two experts indicated an interval for the Likert-scale rang-

Table 1
Likelihood and confidence ratings for the 12 statements in the four beef and dairy cattle animal categories. Raw data: Mean (SD) and median [min – max].

Animal origin (herd)	Animal category							
	Red meat		Replacement calves		Cows		Veal	
	Beef	Dairy	Beef	Dairy	Beef	Dairy	Beef	Dairy
Q1: Receive inappropriate/inadequate diet (e.g. quantity of feed, number of feeding bouts, quality of feed, digestibility etc.).	2.4 (1.4) 2 [1–7]	4.2 (1.9) 4 [1–7]	2.2 (1.4) 2 [1–7]	3.4 (1.7) 3 [1–7]	2.8 (1.3) 3 [1–7]	2.7 (1.6) 2 [1–7]	3.5 (2.0) 4 [1–7]	4.7 (1.9) 5 [1–7]
Q2: Have inadequate water supply (e.g. quantity, quality, accessibility, high competition over access to water).	2.2 (1.3) 2 [1–7]	2.6 (1.5) 2 [1–7]	2.0 (1.2) 2 [1–7]	2.4 (1.5) 2 [1–7]	2.4 (1.3) 2 [1–7]	2.3 (1.4) 2 [1–7]	3.3 (2.3) 3 [1–7]	3.4 (1.8) 4 [1–7]
Q3: Experience discomfort when resting (e.g. when lying down, standing up etc.).	2.4 (1.6) 2 [1–7]	3.8 (1.7) 4 [1–7]	2.1 (1.3) 2 [1–7]	3.1 (1.5) 3 [1–7]	2.2 (1.3) 2 [1–6]	3.7 (1.5) 4 [1–7]	3.6 (2.3) 4 [1–7]	4.5 (1.6) 5 [1–7]
Q4: Experience thermal discomfort (i.e. being too warm (e.g. heat load) or too cold).	3.3 (1.4) 3 [1–7]	3.3 (1.6) 3 [1–7]	3.0 (1.4) 3 [1–6]	3.1 (1.4) 3 [1–6]	3.2 (1.4) 3 [1–6]	3.3 (1.5) 3 [1–7]	3.7 (1.6) 4 [2–6]	3.9 (1.5) 4 [1–6]
Q5: Experience restricted movement (e.g. limited ability to move around freely, e.g. being tied, kept in high stocking density).	2.4 (1.6) 2 [1–7]	4.3 (1.7) 4 [1–7]	2.0 (1.3) 2 [1–7]	3.6 (1.5) 3 [1–7]	2.0 (1.2) 2 [1–7]	3.6 (1.7) 4 [1–7]	3.5 (2.1) 4 [1–7]	5.0 (1.5) 5 [2–7]
Q6: Suffer from injuries (e.g. lameness, skin lesions etc.).	2.6 (1.4) 2 [1–7]	3.4 (1.5) 3 [1–7]	2.2 (1.1) 2 [1–7]	2.9 (1.3) 3 [1–7]	2.7 (1.3) 3 [1–7]	4.7 (1.4) 5 [2–7]	3.0 (1.8) 3 [1–6]	3.9 (1.5) 4 [1–7]
Q7: Suffer from disease	3.1 (1.4) 3 [1–6]	4.2 (1.6) 4 [1–7]	2.6 (1.1) 2 [1–7]	3.6 (1.4) 4 [1–7]	2.7 (1.1) 3 [1–6]	4.2 (1.6) 4 [1–7]	3.8 (1.8) 4 [2–6]	5.0 (1.4) 5 [1–7]
Q8: Suffer from pain induced by management, handling, or surgical procedures	4.1 (2.0) 4 [1–7]	4.4 (1.8) 4 [1–7]	3.2 (1.8) 3 [1–7]	4.1 (1.9) 4 [1–7]	2.7 (1.4) 2 [1–7]	3.4 (1.6) 3 [1–7]	3.5 (1.8) 3 [2–7]	3.8 (1.7) 3 [1–7]
Q9: Have limited opportunities to express normal, non-harmful, social behaviours (e.g. limited allo-grooming, limited social play etc.).	2.2 (1.5) 2 [1–7]	4.0 (1.9) 4 [1–7]	2.0 (1.2) 2 [1–7]	3.4 (1.7) 3 [1–7]	1.9 (1.1) 2 [1–6]	3.2 (1.4) 3 [1–6]	3.8 (2.2) 4 [1–7]	5.0 (1.5) 5 [2–7]
Q10: Have limited opportunities to express other normal behaviours (i.e. limited/restricted from opportunity to forage/graze).	2.7 (1.7) 2 [1–7]	4.9 (1.7) 6 [1–7]	1.9 (1.2) 2 [1–6]	3.9 (1.7) 4 [1–7]	1.8 (1.2) 2 [1–7]	3.9 (1.5) 4 [1–7]	4.1 (2.4) 4 [1–7]	6.0 (1.3) 6 [2–7]
Q11: Experience negative affective states (e.g. fear, distress, frustration or apathy; due to negative human-animal relationships, husbandry/management or the environment in which the animal is kept).	3.0 (1.5) 3 [1–7]	4.3 (1.7) 4 [1–7]	2.6 (1.2) 2 [1–7]	3.4 (1.5) 3 [1–7]	2.5 (1.3) 2 [1–7]	3.6 (1.6) 4 [1–7]	4.2 (2.3) 4 [1–7]	4.9 (1.7) 5 [1–7]
Q12: Have limited opportunities to experience positive emotions (e.g. limited opportunities to engage in rewarding activities).	2.7 (1.6) 2 [1–7]	4.5 (1.6) 5 [1–7]	2.2 (1.3) 2 [1–6]	3.6 (1.5) 4 [1–7]	2.2 (1.2) 2 [1–6]	3.6 (1.4) 4 [1–6]	3.6 (2.3) 3 [1–7]	5.1 (1.7) 5 [1–7]
Confidence in ratings	3.9 (0.9) 4 [1–5]	3.8 (0.8) 4 [2–5]	3.8 (1.1) 4 [1–5]	4.1 (0.8) 4 [2–5]	4.0 (1.0) 4 [1–5]	4.3 (0.7) 4 [2–5]	3.9 (1.0) 4 [2–5]	3.8 (1.0) 4 [1–5]

ing one scale point instead of a single number. In these cases, we used the mid-point of the interval (i.e. the average of the two indicated scale points), which resulted in non-integer scale points at half the original points. Consequently, the resulting Likert-scale had 13 scale points: the integers from 1 to 7 and all the midpoints between the integers (1.5, 2.5, ...). In a next step, we re-scaled the 13-point Likert-scale to a score on a scale from 0 to 1. We did this in a way that kept the steps between the points of the Likert-scale equal and shifted the minimum and maximum scale points half the distance between the points away from 0 and 1: proportion = $4/(2 * 13)$ Likert-scale – $3/(2 * 13)$ (Supplementary Fig. S3). We then logit transformed the numbers of this 0–1 scale (“normalised values”).

The likelihood ratings were dependent on the expert who rated the likelihood of the 12 welfare statements once for each of the two origins and for the four animal categories. To account for this dependency, we calculated the residuals of a mixed model with expert ID as the sole random effect and the intercept as the fixed effect using the 12 normalised values as an outcome variable each. In doing so, we adjusted the average likelihood rating of each expert to the overall average of all experts. Quantile-quantile plots of these 12 sets of residuals showed that they were very close to normally distributed. They thus provided the data for running a principal component analysis (PCA).

Principal component analysis, 12 animal welfare statements

The PCA on the residuals of the normalised values of the 12 welfare statements resulted in a first principal component (PC) that explained 50.3% of the overall variability, which was clearly more than any other PC (Supplementary Table S3, Supplementary Fig. S4). As all the original variables loaded positively on this first PC, it can be interpreted accordingly as the overall welfare risk summarising all the 12 original welfare statements. The highest loads were reached by the risk of restricted ability to express other

normal behaviours, of restricted movement, of restricted ability to express social behaviours, of discomfort when resting, of experiencing negative affective states and of restricted experience of positive emotions (Supplementary Table S3). The beef versus dairy origin was well separated along this first PC (Fig. 1 in main text; Supplementary Fig. S4). All additional PCs explained a much smaller proportion of the variance and were mostly difficult to interpret. Due to their limited contribution to the overall variability in the data, no attempt was made at their interpretation and they were not further considered for analyses.

Statistical analysis of the results

For statistical evaluations, R V 4.0.3 was used (R Core Team 2020). A PCA based on the correlation matrix (function princomp; base R) was run for each set of variables that reflected information on the most common housing systems of the different animal categories in the countries for which experts made an assessment. These PCAs were based on raw scorings of the answers (no = 1, partly = 2, yes = 3) and responses of experts with missing answers were omitted. Veal from beef origin was also omitted because information for this category was given by less than 15% of the experts (n = 10 of 70).

The first principal component of this PCA, which explained 50.3% of the total variance, was then used as the outcome variable in a linear mixed-effects model (function blmer, package blme; Chung et al., 2013, which uses features of the package lme4; Bates, 2015). The fixed effects in the model were the origin of the animals (two-level factor: dairy versus beef), the animal category (four-level factor: calves raised for red meat, calves raised for replacement, cows, and veal calves) and their interaction. To be able to interpret average main effects even in the presence of interactions, sum contrasts were used for the fixed effects. The random effect was animal category nested in expert and the confi-

dence ratings were used as weights. Residuals were checked graphically, using a QQ-plot of the raw residuals and the standard plots using simulated residuals in package DHARMA (Hartig, 2020). The model with the confidence weights showed a slight s-shape in the QQ-plot of DHARMA but the estimated effects were almost unchanged in comparison with the model without weights. We therefore report the model with weights. We calculated p-values by comparing the maximum model with one model each that omitted one of the fixed effects using a parametric bootstrap (package pbrktest; Halekoh and Højsgaard, 2014). A few warnings about non-convergence occurred in the process of evaluation, but these did not seem to influence the model estimates in any relevant way. Finally, we also used a parametric bootstrap to estimate confidence intervals of the model estimates, which fitted the raw data well (Fig. 1).

Results

Overall 70 cattle welfare experts participated in the survey. The experts, who had a median experience of at least 15 years, were recruited from Europe (35), North America (17), South America (8), Australia (5), and other regions of the world (5). Additional characteristics of the experts can be found in [Supplementary Table S1](#).

Welfare risk as a function of the different origins and animal production goals

Experts' likelihood and confidence ratings for each of the 12 statements are shown in [Table 1](#). The overall likelihood to experience a negative welfare state (i.e. welfare risk) was assessed by the experts as higher in animals from dairy in comparison to beef, for all animal categories (origin: $P = 0.001$, interaction: $P = 0.28$; [Fig. 1B](#)). The overall welfare risk increased from calves raised for replacement and cows, to calves raised for red meat and to veal calves ($P = 0.001$; [Fig. 1B](#)). These results indicate that, regardless of the production goal (calves raised for red meat, calves raised for veal, calves raised as replacement, and cows), animals born in dairy herds were considered to experience worse welfare than animals born in beef herds.

Principal component analyses, information on the housing systems

The analyses of the information on the housing systems showed that beef versus dairy origin can be well separated along the first PC in all the animal categories ([Table S3](#), [Fig. S5](#), a–d). For the calves raised for red meat, the housing systems could be described by two PCs, of which the first was composed of access to pasture, access to dam and suckling from dam during the first months of life and not being transported during the first 6 months of life (“extensivity” which may be closely related to available space), whereas the second concerned being slaughtered before or after 18 months of age ([Table S3](#)). Beef and dairy systems seemed to differ mostly by the beef systems being more extensive ([Fig. S5a](#)). For the calves raised for replacement, the results were very similar with the exception that the transportation loaded on the second PC ([Table S3](#), [Fig. S5b](#)) and that, due to their purpose, they were not slaughtered by default. The housing of beef and dairy cows seemed to vary mostly in the extent that access to the outdoors was provided ([Table S3](#), [Fig. S5c](#)). Here, beef cows seemed to have access to the outdoors more commonly than dairy cows and dairy cows ran a higher risk of being culled early. The housing of veal calves seemed to vary the most with respect to the type of flooring provided with bedding material and access to pasture versus housing on slatted floors loading heavily on the first PC ([Table S3](#), [Fig. S5d](#)). The sec-

ond PC was a contrast of the provision of (additional) solid feed and being slaughtered at a young age.

Discussion

The collection of bovine milk by humans has a long tradition (“secondary products revolution”; Sherratt, 1983), and it seems to have a promising future, based on recent projections by the Organisation for Economic Co-operation and Development (OECD) as well as the Food and Agriculture Organisation agency (FAO; OECD, 2020). For this reason, it is important to critically reflect on the welfare impact of this practice. Our expert survey showed that bovines raised in the main current food production systems are rated as more likely to experience negative welfare conditions if raised in dairy systems (regardless of whether they are used for their milk or meat) than in beef systems (i.e. when raised solely for their meat). This assessment was very similar in respect to the geographic region of experts (Europe, North America, others). Experts from different geographical regions only differed in their assessment of veal production (see supplementary analysis in [Supplementary Material S1](#)). The clear overall assessment of a higher welfare risk in dairy versus beef systems may be surprising in light of societal perceptions regarding ethical food choices (i.e. vegetarianism; Fox and Ward, 2008; Corrin and Papadopoulos, 2017) and deserves further consideration to identify likely underpinning reasons.

Our assessment protocol allowed us to cover a wide range of production methods (for the result of the PCAs for the information on the housing systems, see [Supplementary Fig. S5](#)) and to assess experts' ratings of the overall welfare risk of animals raised in these conditions during the majority of their lifetime (for the ‘raw’ ratings of experts, for each of the 12 areas of welfare concern, see [Table 1](#)). Nevertheless, it did not allow us to disentangle which of the welfare risks mentioned above formed the basis for the experts' ratings. In addition, we emphasise that the results do not necessarily mean that animals born in dairy herds are, at any given point of time and in every type of system, worse off than animals born in beef herds. For example, after spending several months on pasture with their dam, the welfare of beef suckler calves is expected to be substantially reduced once they are moved to feedlots, potentially below the levels experienced by many dairy calves raised for red meat (see [Fig. 1](#) in [Bracke et al., 2008](#); for a review of the factors affecting cattle welfare in feedlots, see [Tucker et al. \(2015\)](#) and [Salvin et al. \(2020\)](#); for cattle preferences to being on pasture over the feedlot area, see [Lee et al. \(2013\)](#)). It would be interesting to explore whether experts would consider that improvements to the welfare of bovines in the dairy industry e.g. by keeping dairy calves with their dams ([Meagher et al., 2019](#)), free access to pasture ([Mee and Boyle, 2020](#)) and the use of pain relief, e.g. in case of lameness ([Whay et al., 2005](#)) and during practices such as dehorning ([Stafford and Mellor, 2015](#)) would be sufficient to balance the overall welfare risk for animals raised in the two production systems. The answer cannot be predicted in advance because, for example, if such measures were similarly applied to bovines in the beef industry, then they may retain their perceived higher welfare status.

One possible reason why experts rated dairy systems as more harmful to welfare may be because raising bovines for their milk (and meat) involves a higher (negative) intervention in their lives compared to raising them solely for their meat. Raising bovines for their meat involves feeding and slaughtering them. Dairy cows are also bred for their milk, which is then collected 1–3 times per day, often for 305 days or more per lactation ([Dematawewa et al., 2007](#); [Sehested et al., 2019](#)), and this has implications for how these animals are raised and managed. Long-term genetic selection

for high milk yield in dairy cows has been recognised as a major factor causing poor welfare, in particular health problems, such as lameness, mastitis, reproductive disorders and metabolic disorders (EFSA, 2008). Adoption of high-yielding breeds such as the Holstein-Friesian without consideration for the animals' natural ability to cope with diseases (EFSA, 2008) and thermal challenges typical of extreme climates may lead to additional welfare compromises (Von Keyserlingk and Hötzel, 2015; Von Keyserlingk et al., 2013). Common housing and husbandry procedures characterising the management of dairy cattle increase the welfare risks further. Our results show that beef and dairy systems raising calves for red meat and as a replacement, differed mostly in relation to their "extensivity", indicated by their ability to access their dam and suckle from her during the first months of life, as well as from their ability to access pasture (Supplementary Table S2). Indeed, dairy calves, in contrast to beef calves, are commonly separated from their dam a few hours after birth to allow the collection of milk from their mothers (EFSA, 2008; Beaver et al., 2019). Upon separation, they are often kept in social isolation for several weeks (i.e. limited or no physical contact with their dam and conspecifics), a management practice that originated from the desire to reduce horizontal transmission of disease between calves yet has been repeatedly shown to inflict behavioural and developmental harm (Costa et al., 2016, 2019). The social and nutritional restrictions following the separation from the dam have been implicated as key animal welfare issues in commercially raised dairy calves (Costa et al., 2019), with the former being associated with cognitive and social impairments affecting the welfare of the animals also at a later age (for additional welfare risks placed on male dairy calves, that are often transported to a different farm at a young age, see Wilson et al., 2020). Later in life, dairy calves raised to produce milk ("replacement dairy calves") are commonly housed indoors for part of the year (i.e. winter) or continuously (i.e. zero-grazing system), in free or tie-stall systems (Barkema et al., 2015). Keeping dairy cattle indoors is associated with numerous behavioural restrictions (Mandel et al., 2016; Arnott et al., 2017) and health risks, such as higher incidence of lameness (Haskell, 2020), and increased risk for claw or foot problems, teat trampling, mastitis, metritis, dystocia, ketosis, retained placenta, and some bacterial infections compared to systems that allow cattle access outdoors (EFSA, 2008). Since dairy cattle are commonly raised indoors for at least part of the year, and since their management involves a high degree of intervention as described above (in the first weeks of life following the separation from their dam, and throughout the lactating periods, when milked by partially/fully automated milking systems), their management is expected to involve a higher risk to their welfare compared to beef cattle. A future study could specifically explore which of these factors most influence expert rankings of welfare.

A possible step towards minimising the welfare gap between the beef and the dairy sector would be to refine (or simply eliminate, when possible), husbandry practices that have long been recognised as compromising the welfare of both cows and their calves, such as, for example, early separation of the calves from their dams (Beaver et al., 2019). Dam rearing throughout the first months of life may certainly improve some welfare aspects for both the dam and the calf (Meagher et al., 2019), and may meet current demands/perception of welfare among consumers (Busch et al., 2017). Overcoming the challenges associated with existing dam-rearing systems (e.g. Wenker et al., 2022) may prove to be an important step in the process. A complementary approach, which goes beyond minimising the gap between the two sectors (dairy/beef), would be to elevate the overall positive welfare balance in both the dairy and the beef sectors (Rault et al., 2020). More access to pasture, in suitable climate conditions, can definitely be a step in the right direction, especially for dairy herds (Arnott et al.,

2017). Yet, it seems to contradict the general trend of a transition to indoor housing of cattle (zero-grazing systems). Another complementary direction would be to improve the training of animal handlers, especially when in daily contact with the animals (Rault et al., 2020). Unfortunately, in many regions of the world, the training of animal handlers is still not mandated by law. In countries that do demand training/certification (e.g. Switzerland), periodic training updates are not obligatory (or apply only to a limited set of professions, e.g., for animal transport personal, but not for farmers, BLV, 2022). Lastly, where high productivity has taken precedence over basic bodily strength and integrity, the use of healthier/more robust dairy genotypes would greatly improve welfare (Rauw et al., 1998; Webster, 2021).

Our study focused on the welfare risks of dairy and beef cattle, before being slaughtered/culled. It is important to note, however, that the period during which the animals are subjected to these risks varies between production goals, production stages and the management goals of the farm (Fig. 1A). Veal calves (from both dairy and beef herd origins) are usually slaughtered at the age of 6–11 months, depending on whether they are used to produce white or rose veal [but see culling of 'surplus'/bobby dairy calves within the first days of life (Haskell et al., 2006)]. Calves raised for red meat (from both dairy and beef herds) are usually slaughtered at 12–36 months, while replacement cows, which are used in both types of systems to produce both milk and calves, will commonly be slaughtered at an earlier age in dairy herds than beef herds. In the US, for example, dairy cows are slaughtered at about 5 years of age (after 2.5–3 lactations), while beef cows are slaughtered at 7–12 years of age (Moreira et al., 2021). The production of milk, like beef production, involves killing of animals, yet in many cases, this happens at a younger age in the dairy industry. Factoring the intensity of welfare compromise in the lifetime expectancy (i.e. duration) of these animals could deepen our understanding of the overall welfare impact. However, it is beyond the scope of this paper. Here and throughout the manuscript, we avoid making any ethical claims about the sanctity of life. Also, we do not aim to value the amount of suffering per animal in relation to the amount of animal protein (or calorie; Kolbe, 2018). Our aim was more straightforward, to assess, based on expert opinion, the welfare of animals born to common dairy and beef herds, until slaughtered/culled.

At a time of rising public concern for the welfare of animals (Miele et al., 2013) and awareness of the impact of our diets on the environment (Kearney, 2010), it is important that research priorities and dietary choices are aligned with the areas where welfare problems are most apparent. We call for further comparisons of bovine welfare in less common dairy and beef production systems that were not covered here (e.g. organic systems, dam-rearing systems, where dairy calves are kept together with their dam), preferably using animal-based welfare assessment on farm. It would also be valuable to obtain representative overview of the prevalence of these different systems across countries, and to explore the possible effect of animal welfare legislation (e.g. federal/union/state level) on the magnitude of welfare differences between the dairy and beef sectors. For a preliminary assessment of this idea (and its limitations), see S6. In addition, we encourage a similar expert comparison of other farm species that, like dairy cows, are used for "more than their meat"; e.g. laying hens could be compared with broilers to see if the findings from this study apply in other areas of animal agriculture.

Conclusion

Cattle welfare experts rated dairy cattle as substantially more likely to experience negative welfare than beef cattle in the most

common housing systems selected by the experts. The underpinning reasons for these evaluations were not explored with experts but are proposed to enable testable predictions for future research. Raising awareness about the linkage between dairy and meat production, and the toll of milk production on the welfare state of animals in the dairy industry, may encourage a more sustainable and responsible food consumption.

Supplementary material

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.animal.2022.100622>.

Ethics approval

An approval to conduct this study was granted by the Research Ethics Committee for SCIENCE and HEALTH of the university of Copenhagen, ref: SUSTA, case: 504-0272/21-5000. Approval and registration of processing of personal data in the project were provided by the legal department of the university of Copenhagen, ref PWI, case: 514-0273/21 – 5000.

Data and model availability statement

The data and models are deposited in an official repository (https://github.com/ku-awdc/dairy_vs_beef) and are freely available to the reader.

Author ORCIDs

Roi Mandel: <https://orcid.org/0000-0003-0714-5053>.
Marc B.M. Bracke: <https://orcid.org/0000-0003-0478-7771>.
Christine J. Nicol: <https://orcid.org/0000-0001-6734-2177>.
John A. Webster: NA.
Lorenz Gyax: <https://orcid.org/0000-0001-8546-2930>.

Author contributions

RM, LG, MBMB and **CJN** designed research; **RM** performed research; **LG** and **RM** analysed data; **RM, LG, MBMB, CJN** and **JAW** wrote the paper.

Declaration of interest

None.

Acknowledgements

A previous version of this manuscript was published as a preprint (Mandel et al., 2021).

Financial support statement

This research received no specific grant from any funding agency, commercial or not-for-profit section.

References

Arnott, G., Ferris, C.P., O'Connell, N.E., 2017. Welfare of dairy cows in continuously housed and pasture-based production systems. *Animal* 11, 261–273.
 Barkema, H.W., von Keyserlingk, M.A., Kastelic, J.P., Lam, T.J.G.M., Luby, C., Roy, J.P., LeBlanc, S.J., Keefe, G.P., Kelton, D.F., 2015. Invited review: Changes in the dairy industry affecting dairy cattle health and welfare. *Journal of Dairy Science* 98, 7426–7445.

Bates, D., 2015. Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software* 67, 1–48.
 Beaver, A., Meagher, R.K., von Keyserlingk, M.A., Weary, D.M., 2019. Invited review: A systematic review of the effects of early separation on dairy cow and calf health. *Journal of Dairy Science* 102, 5784–5810.
 Bertocchi, L., Fusi, F., Angelucci, A., Bolzoni, L., Pongolini, S., Strano, R.M., Ginestreti, J., Riuzzi, G., Moroni, P., Lorenzi, V., 2018. Characterization of hazards, welfare promoters and animal-based measures for the welfare assessment of dairy cows: Elicitation of expert opinion. *Preventive Veterinary Medicine* 150, 8–18.
 BLV, 2022. Training and further education in animal husbandry and in the handling of animals. Federal Food safety and veterinary office, Switzerland. Information retrieved on 10th June 2022 from <https://www.blv.admin.ch/blv/de/home/tiere/tierschutz/aus-und-weiterbildung.html>.
 Bracke, M.B.M., Edwards, S.A., Engel, B., Buist, W.G., Algers, B., 2008. Expert opinion as 'validation' of risk assessment applied to calf welfare. *Acta Veterinaria Scandinavica* 50, 29. <https://doi.org/10.1186/1751-0147-50-29>. Accessed 22-10-2019.
 Bracke, M.B., Koene, P., Estevez, I., Butterworth, A., de Jong, I.C., 2019. Broiler welfare trade-off: A semi-quantitative welfare assessment for optimised welfare improvement based on an expert survey. *PLoS One* 14, e0222955.
 Busch, G., Weary, D.M., Spiller, A., von Keyserlingk, M.A., 2017. American and German attitudes towards cow-calf separation on dairy farms. *PLoS One* 12, e0174013.
 Chung, Y., Rabe-Hesketh, S., Dorie, V., Gelman, A., Liu, J., 2013. A nondegenerate penalized likelihood estimator for variance parameters in multilevel models. *Psychometrika* 78, 685–709.
 Corrin, T., Papadopoulos, A., 2017. Understanding the attitudes and perceptions of vegetarian and plant-based diets to shape future health promotion programs. *Appetite* 109, 40–47.
 Costa, J.H., Cantor, M.C., Adderley, N.A., Neave, H.W., 2019. Key animal welfare issues in commercially raised dairy calves: social environment, nutrition, and painful procedures. *Canadian Journal of Animal Science* 99, 649–660.
 Costa, J.H.C., Von Keyserlingk, M.A.G., Weary, D.M., 2016. Invited review: Effects of group housing of dairy calves on behavior, cognition, performance, and health. *Journal of Dairy Science* 99, 2453–2467.
 Dawson, L.C., Dewey, C.E., Stone, E.A., Guerin, M.T., Niel, L., 2016. A survey of animal welfare experts and practicing veterinarians to identify and explore key factors thought to influence canine and feline welfare in relation to veterinary care. *Animal Welfare* 25, 125–134.
 Dematawewa, C.M.B., Pearson, R.E., VanRaden, P.M., 2007. Modeling extended lactations of Holsteins. *Journal of Dairy Science* 90, 3924–3936.
 EFSA Panel on Animal Health and Animal Welfare (AHAW), 2008. Scientific Opinion on the overall effects of farming systems on dairy cow welfare and disease. *EFSA Journal* 1143, 1–38.
 EFSA Panel on Animal Health and Animal Welfare (AHAW), 2015. Scientific Opinion on welfare aspects of the use of perches for laying hens. *EFSA Journal* 13, 4131.
 EFSA Panel on Animal Health and Welfare (AHAW), 2012a. Guidance on risk assessment for animal welfare. *EFSA Journal* 10, 2513.
 EFSA Panel on Animal Health and Welfare (AHAW), 2012b. Scientific Opinion on the welfare of cattle kept for beef production and the welfare in intensive calf farming systems. *EFSA Journal* 10, 2669.
 EFSA Panel on Animal Health and Welfare (AHAW), 2014. Scientific opinion on the welfare risks related to the farming of sheep for wool, meat and milk production. *EFSA Journal* 12, 3933.
 Fox, N., Ward, K., 2008. Health, ethics and environment: A qualitative study of vegetarian motivations. *Appetite* 50, 422–429.
 Halekoh, U., Højsgaard, S., 2014. A kenward-roger approximation and parametric bootstrap methods for tests in linear mixed models—the R package pbkrtest. *Journal of Statistical Software* 59, 1–30.
 Hartig, F., 2020. DHARMA: Residual Diagnostics for Hierarchical (Multi-Level/Mixed) Regression Models. R package version 0.3.3.0. Retrieved on 25.11.2020 from: <https://CRAN.R-project.org/package=DHARMA>.
 Haskell, M.J., 2020. What to do with surplus dairy calves? Welfare, economic and ethical considerations. *Landbauforschung* 70, 45–48.
 Haskell, M.J., Rennie, L.J., Bowell, V.A., Bell, M.J., Lawrence, A.B., 2006. Housing system, milk production, and zero-grazing effects on lameness and leg injury in dairy cows. *Journal of Dairy Science* 89, 4259–4266.
 Kearney, J., 2010. Food consumption trends and drivers. *Philosophical Transactions of the Royal Society B: Biological Sciences* 365, 2793–2807.
 Keeling, L., Tunón, H., Olmos Antillón, G., Berg, C., Jones, M., Stuardo, L., Swanson, J., Wallenbeck, A., Winckler, C., Blokhuis, H., 2019. Animal welfare and the United Nations sustainable development goals. *Frontiers in Veterinary Science* 6, 336.
 Kolbe, K., 2018. Why milk consumption is the bigger problem: Ethical implications and deaths per calorie created of milk compared to meat production. *Journal of Agricultural and Environmental Ethics* 31, 467–481.
 Lee, C., Fisher, A.D., Colditz, I.G., Lea, J.M., Ferguson, D.M., 2013. Preference of beef cattle for feedlot or pasture environments. *Applied Animal Behaviour Science* 145, 53–59.
 Mandel, R., Bracke, M.B., Nicol, C.J., Webster, J., Gyax, L., 2021. Dairy vs. beef production—expert views on welfare. *bioRxiv*. <https://doi.org/10.1101/2021.12.06.471462>.
 Mandel, R., Whay, H.R., Klement, E., Nicol, C.J., 2016. Invited review: Environmental enrichment of dairy cows and calves in indoor housing. *Journal of Dairy Science* 99, 1695–1715.
 McGreevy, P., Berger, J., De Brauwere, N., Doherty, O., Harrison, A., Fiedler, J., Jones, C., McDonnell, S., McLean, A., Naconechny, L., Nicol, C., Preshaw, L., Thomson, P.,

- Tzioumis, Webster, J., Wolfensohn, S., Yeates, J., Jones, B., 2020. Using the five domains model to assess the adverse impacts of husbandry, veterinary, and equitation interventions on horse welfare. *Animals* 8, 41.
- Meagher, R.K., Beaver, A., Weary, D.M., von Keyserlingk, M.A.G., 2019. Invited review: A systematic review of the effects of prolonged cow-calf contact on behavior, welfare, and productivity. *Journal of Dairy Science* 102, 5765–5783.
- Mee, J.F., Boyle, L.A., 2020. Assessing whether dairy cow welfare is “better” in pasture-based than in confinement-based management systems. *New Zealand Veterinary Journal* 68, 168–177.
- Miele, M., Blokhuis, H., Bennett, R., Bock, B., 2013. Changes in farming and in stakeholder concern for animal welfare. In: *Improving farm animal welfare*. Wageningen Academic Publishers, Wageningen, The Netherlands, pp. 19–47.
- Moreira, L.C., Rosa, G.J.M., Schaefer, D.M., 2021. Beef production from cull dairy cows: a review from culling to consumption. *Journal of Animal Science* 99, 1–18.
- Nicol, C., Bejder, L., Green, L., Johnson, C., Keeling, L., Noren, D., Van der Hoop, J., Simmonds, M., 2020. Anthropogenic threats to wild cetacean welfare and a tool to inform policy in this area. *Frontiers in Veterinary Science* 7, 57.
- OECD, 2020. *OECD-FAO Agricultural forecast 2020–2029* (OECD Publishing, Paris, 2020). Retrieved on 23 November 2021 from <https://www.fao.org/documents/card/en/c/ca8861en>.
- R Core Team, 2020. *R: A language and environment for statistical computing* [<https://www.r-project.org/>]. R Foundation for Statistical Computing, Vienna, Austria.
- Rault, J.L., Hintze, S., Camerlink, I., Yee, J.R., 2020. Positive welfare and the like: Distinct views and a proposed framework. *Frontiers in Veterinary Science* 7, 370.
- Rauw, W.M., Kanis, E., Noordhuizen-Stassen, E.M., Grommers, F.J., 1998. Undesirable side effects of selection for high production efficiency in farm animals: a review. *Livestock Production Science* 56, 15–33.
- Salvin, H.E., Lees, A.M., Cafe, L.M., Colditz, I.G., Lee, C., 2020. Welfare of beef cattle in Australian feedlots: A review of the risks and measures. *Animal Production Science* 60, 1569–1590.
- Sehested, J., Gaillard, C., Lehmann, J.O., Maciel, G.M., Vestergaard, M., Weisbjerg, M. R., Mogensen, L., Larsen, L.B., Poulsen, N.A., Kristensen, T., 2019. Review: extended lactation in dairy cattle. *Animal* 13, 65–74.
- Sherratt, A., 1983. The secondary exploitation of animals in the Old World. *World Archaeology* 15, 90–104.
- Stafford, K.J., Mellor, D.J., 2015. Addressing the pain associated with disbudding and dehorning in cattle. *Applied Animal Behaviour Science* 135, 226–231.
- Tucker, C.B., Coetzee, J.F., Stookey, J.M., Thomson, D.U., Grandin, T., Schwartzkopf-Genswein, K.S., 2015. Beef cattle welfare in the USA: identification of priorities for future research. *Animal Health Research Reviews* 16, 107–124.
- United Nations (UN), 2015. *Transforming our world: The 2030 agenda for sustainable development*. UN, New York, NY, USA.
- Von Keyserlingk, M.A., Hötzel, M.J., 2015. The ticking clock: Addressing farm animal welfare in emerging countries. *Journal of Agricultural and Environmental Ethics* 28, 179–195.
- Von Keyserlingk, M.A.G., Martin, N.P., Kebreab, E., Knowlton, K.F., Grant, R.J., Stephenson, M., Sniffen, C.J., Harner III, J.P., Wright, A.D., Smith, S.I., 2013. Invited review: Sustainability of the US dairy industry. *Journal of Dairy Science* 96, 5405–5425.
- Webster, J., 2021. Green Milk From Contented Cows: Is It Possible? *Frontiers in Animal Science* 2, 667196.
- Welfare Quality®, 2009. *Assessment protocol for cattle*. Welfare Quality® Consortium, Lelystad, Netherlands. Retrieved on 01 August 2020 from <http://www.welfarequalitynetwork.net/en-us/reports/assessment-protocols>.
- Wenker, M.L., Verwer, C.M., Bokkers, E.A., Te Beest, D.E., Gort, G., De Oliveira, D., Koets, A., Bruckmaier, R.M., Gross, J.J., Van Reenen, C.G., 2022. Effect of Type of Cow-Calf Contact on Health, Blood Parameters, and Performance of Dairy Cows and Calves. *Frontiers in Veterinary Science* 9, 855086.
- Whay, H.R., Webster, A.J.F., Waterman-Pearson, A.E., 2005. Role of ketoprofen in the modulation of hyperalgesia associated with lameness in dairy cattle. *Veterinary Record* 157, 729–733.
- Wilson, D.J., Canning, D., Giacomazzi, T., Keels, K., Lothrop, R., Renaud, D.L., Sillett, N., Taylor, D., Van Huigenbos, H., Wynands, B., Zuest, D., 2020. Hot topic: health and welfare challenges in the marketing of male dairy calves—Findings and consensus of an expert consultation. *Journal of Dairy Science* 103, 11628–11635.